Large field versus small field expansions and Sobolev inequalities

Pirmin Lemberger

Centre de Physique Théorique,
Ecole Polytechnique, 91128 Palaiseau, France

April 25, 1997

Abstract

We study a model for a two dimensional random interface $\phi(x)$, $x \in {}^2$ described by a massless gaussian measure perturbed by a weak potential $V(\phi) = \frac{\varepsilon^2}{2}(\mathrm{e}^{-\alpha\phi} - 1)^2$. Such a model occurs for instance in a phenomenological description of the wetting transition. We prove that, provided α is small enough, the two-point function decreases exponentially with a rate of order $m \equiv \varepsilon\alpha$ which is just the mean field value. The large field region problem due to the fact that $V(\phi)$ remains bounded when $\phi \to +\infty$ is treated by means of a large field versus small field expansion combined with elementary Sobolev inequalities. The paper is intended to be accessible to non-experts.

PACS numbers: **02.90.+p**, 68.45.Gd
KEY WORDS: Cluster-expansions, large field/small field decomposition, Sobolev inequalities, wetting transition.

1 Introduction

A fundamental difficulty in statistical mechanics and constructive field theory is to perform the thermodynamic limit in a mathematicaly rigorous way. In statistical mechanics the existence of such a limit simply corresponds to the extensivity of thermodynamic potentials like the pressure, the free energy and so on. In field theory, it proves in the nontrivial cases the existence of non-gaussian translation invariant measures on some distribution spaces. One of the methods to control this limit is the cluster expansion. It is basically perturbative but allows for very general types of interactions and gives detailed information about the limiting theory in contrast with non-perturbative methods, like correlation inequalities. Let us briefly recall the setting in field theory when no renormalisation is needed. One is given a gaussian measure $\mathrm{d}\mu_C(\phi)$ with covariance C on a space of distributions, e.g. $\mathcal{S}'({}^n)$, and a functional $F_\Lambda[\phi]$ depending only on the field in the volume $\Lambda \subset {}^n$. F_Λ is usually assumed to have some factorisation properties, for example $F_\Lambda = \prod_{\Delta\subset\Lambda} F_\Delta$, where the Δ's are squares covering Λ. The problem then is to control the $|\Lambda| \to \infty$ limit of the free energy $f_\Lambda \equiv |\Lambda|^{-1} \log Z(\Lambda)$ where $Z(\Lambda) \equiv \int \mathrm{d}\mu_C(\phi) F_\Lambda[\phi]$. The cluster expansion is an expansion in the volume for $Z(\Lambda)$. More precisely $Z(\Lambda)$ is written as a finite sum of contributions associated to each partition of the volume Λ into disjoint subsets X_j, each X_j being a collection of squares $\Delta \subset \Lambda$. The contribution to a given partition factorises over the subsets X_j. The sets X_j are called clusters or polymers. The factors $z(X_j)$ associated to a given polymer is its activity. So basically the cluster expansion is a rewriting of the partition function $Z(\Lambda)$ as a system of polymers X_j interacting through a hard core exclusion. The standard algebraic machinery of Mayer series [1], therefore allows to write a formal power series in the activities $z(X_j)$ for the free energy $f \equiv \lim_{\Lambda \to \infty} f_\Lambda$ in the thermodynamic limit. The hard work really begins with showing that $|z(X_j)|$ is small enough for this series to converge towards f. To be a little bit more specific consider the case where $F_\Delta = \exp(-\lambda V_\Delta[\phi])$, where $V_\Delta[\phi]$ is some potential. There are two ways in which $|z(X_j)|$ should be small. First $|z(X_j)|$ has to become exponentially small with the volume $|X_j|$ occupied by the polymer X_j. One necessary condition for this is that $|\lambda|$ be small enough. Second, $|z(X_j)|$ must become small whenever the squares Δ comprising X_j are far apart from each other. From the explicit formula for the activity one can show that a sufficient condition for this is that the propagator $C(x, y)$ decays sufficiently fast at infinity. Actually, integrability at infinity is enough. An example satisfying this condition is the free field of mass $m > 0$ for which $C(x, y) \leq \mathbf{C} \exp(-m|x - y|)$, $\mathbf{C} > 0$ as $|x - y| \to \infty$.

In this paper we want to study a model in which a two-dimensional massless gaussian measure is perturbed by a weak potential. We choose $V_\Lambda[\phi] \equiv \int_\Lambda V(\phi(x))\mathrm{d}x$ with $V(\phi) \equiv \frac{\varepsilon^2}{2}(\mathrm{e}^{-\alpha\phi} - 1)^2$. Notice that for $|\phi| \ll \alpha^{-1}$ we have the quadratic approximation $V(\phi) \simeq \frac{m^2\phi^2}{2}$ where $m \equiv \varepsilon\alpha$. The main point about $V(\phi)$ is that it remains bounded when $\phi \to +\infty$. Hence the interacting measure will be effectively massless in the regions where the field is large and positive. This is actually the major source of difficulty and interest for this model. Recall that the massless propagator is not integrable, therefore a naive cluster expansion will not work. Actually the kernel of the covariance is not even defined in two dimensions in the infinite volume limit. There are three reasons for choosing this particular V.

- The V above appears in a phenomenological description of the wetting transition [2, 3]. The field $\phi(x)$ corresponds for example to the height of a liquid/vapor interface above a wall. $V(\phi)$ is a weak potential which tends to localise the interface near the wall. If we put Dirichlet boundary conditions on $\partial\Lambda$, the massless gaussian measure is defined by the covariance $C_{0,\Lambda} \equiv (-\Delta_D)^{-1}$ and just penalizes high curvatures of the interface ϕ. A direct computation shows that $C_{0,\Lambda}(x, x) \sim \log|\Lambda|$, hence we expect that in the thermodynamic limit $\langle\phi(x)^2\rangle_V \to \infty$ as $\varepsilon \to 0$. This phenomenon is supposed to describe how a liquid/vapor interface delocalises as the temperature T approaches from below the critical wetting temperature T_w. One can identify ε with $T_w - T > 0$ see [4].

- The particular form of the V chosen above seems to be well suited for a multiscale analysis, which would allow to control the model outside the mean-field regime where the mass is equal to $\alpha\varepsilon$. We shall not deal with these questions here, see however section 2.

- At last $V(\phi)$ is a simple example for which a large field versus small field decomposition is needed. Many of the most interesting models in constructive field theory (non-Abelian gauge theories [5], two-dimensional Gross-Neuveu models [6]) or many body theory (BCS theory of supraconductivity) require a similar treatment.

Our aim is to prove that, at least in some regime of the parameters α and ε, the potential $V(\phi)$ generates a mass $m' = O(m)$, in other words we want to prove that the connected two-point functions verifies $|\langle\phi(x)\phi(y)\rangle_V - \langle\phi(x)\rangle_V\langle\phi(y)\rangle_V| \leq \mathbf{C}\mathrm{e}^{-m'|x-y|}$.

To show this we first have to find good definitions for what is meant by small and large field regions. Then we must show that big large-field regions, in which there is no exponential decay of the correlations, are rare. That this is true can be understood intuitively as follows. Take a square $\Delta \subset \Lambda$ (whose size will have to be chosen properly), and suppose ϕ is large in mean on Δ. There are just two possibilities: either ϕ is more or less constant on Δ in which case $\exp(-V_\Delta[\phi])$ is small, or this last factor is not small but in that case ϕ necessarily has a large mean curvature and thus $\exp[-\frac{1}{2}\int_\Delta(\nabla\phi)^2]\mathrm{d}x$ is small. This is where the Sobolev inequalities enter. They allow to treat this alternative in a natural and efficient way. These inequalities replaces a substantial part of the very heavy formalism of [7, 8]. As an input for the Sobolev inequalities we need to extract a factor $\exp[-\frac{1}{2}\int_\Delta(\nabla\phi)^2]\mathrm{d}x$ per square in the large field regions from a "locally massless" non-translation invariant gaussian measure. This is a technically non-trivial problem which could have applications in other contexts. It is solved by means of some operator inequalities.

The paper is organized as follows. In section 2 we describe mathematically the model we want to study and state our main result. Section 3 shows how to generate a cluster expansion with small versus large field decomposition. To generate this expansion we make an easy adaptation from Brydges formula in [9]. Its main advantage is its compactness and that it avoids the pain of going through many recursive formulas. A polymer X in this new cluster expansion will be a collection of squares $\{\Delta\}_{j=1}^{s(X)}$ *and* large field regions $\{\gamma\}_{j=1}^{l(X)}$. In section 4 we shall bound the activities $z(X)$ and prove the convergence of the cluster expansion. The last section contains the proof of two crucial propositions. The first uses Sobolev inequalities to show that in the large field region the product $\exp(-V_\Delta[\phi])\exp[-\frac{1}{2}\int_\Delta(\nabla\phi)^2\mathrm{d}x]$ is very small if α is small. The second shows how to extract the factor $\exp[-\frac{1}{2}\int_\Delta(\nabla\phi)^2\mathrm{d}x]$ from the locally massless gaussian measure which appears in the expression for the activities. Proofs are given in details, even when the arguments are rather standard. This makes the paper a little bit longer but hopefully more readable for people who are, like me, beginners in the cluster expansion business.