

# ASYMPTOTIC BEHAVIOR OF UNSTABLE PERTURBATIONS OF THE FUBINI–STUDY METRIC IN RICCI FLOW

DAVID GARFINKLE, JAMES ISENBERG, DAN KNOPF, AND HAOTIAN WU\*

ABSTRACT. Kröncke has shown that the Fubini–Study metric is an unstable generalized stationary solution of Ricci flow [Krö20]. In this paper, we carry out numerical simulations which indicate that Ricci flow solutions originating at unstable perturbations of the Fubini–Study metric develop local singularities modeled by the blowdown soliton discovered in [FIK03].

## CONTENTS

1. Introduction	1
2. An unstable conformal perturbation	3
3. Cohomogeneity-one metrics	4
4. Initial data	5
5. Ricci–DeTurck flow	6
6. A system better suited to simulation	7
7. The blowdown soliton	9
8. Results of simulations	10
9. Directions for future work	13
References	13

## 1. INTRODUCTION

In studying dynamical systems, it is always important to identify and classify fixed points. For Ricci flow, regarded as a dynamical system on the infinite-dimensional space of Riemannian metrics, one must consider generalized fixed points. These are the *Ricci solitons*, specified by the data  $(\mathcal{M}^n, g, \lambda, X)$  for which

$$-2 \operatorname{Rc}[g] = 2\lambda g + \mathcal{L}_X g,$$

where  $\operatorname{Rc}[g]$  is the Ricci curvature of the metric  $g$ ,  $\lambda \in \mathbb{R}$ , and  $\mathcal{L}_X g$  is the Lie derivative of  $g$  with respect to the vector field  $X$ .

---

*Key words and phrases.* Ricci flow, blowdown soliton, unstable perturbations, Fubini–Study metric, asymptotic analysis, numerical method.

\*Author to whom any correspondence should be addressed.

DG thanks the NSF for support in PHY-1806219 and PHY-2102914. JI thanks the NSF for support in PHY-1707427. DK thanks the Simons Foundation for support in Award 635293. HW thanks the Australian Research Council for support in DE180101348.

A prominent example of a Ricci soliton is the Fubini–Study metric on complex projective space, an Einstein metric corresponding to  $(\mathbb{C}\mathbb{P}^N, G_{\text{FS}}, -2(N+1), 0)$ .<sup>1</sup> In §10 of [Ham95], Hamilton conjectures that it is a stable fixed point. We refer the reader to Section 4 for an explicit description of the Fubini–Study metric. In Example 2.3 of [CHI04], Cao, Hamilton, and Ilmanen note that the Fubini–Study metric is neutrally linearly stable with respect to the second variation of Perelman’s shrinker entropy, crediting this observation to unpublished work of Goldschmidt. However, in his dissertation, Kröncke computes the third variation of Perelman’s entropy to prove the surprising result that Fubini–Study is in fact unstable. This calculation has appeared as [Krö20] and has been independently verified in [KŠ19]. The instability is conformal and hence not Kähler.

The instability of the Fubini–Study raises an interesting question. What happens to Ricci flow solutions that start at arbitrarily small but unstable perturbations of  $G_{\text{FS}}$ ? According to the conjectural hierarchy of 4-dimensional solutions given in [CHI04], one expects the Ricci flow of unstable perturbations of  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$  to develop local singularities modeled by the blowdown gradient shrinking soliton  $\mathcal{L}_{-1}^2$  discovered in [FIK03]. These solutions would start arbitrarily close to (but *not* in) the subset of Kähler metrics and, in finite time, develop local singularities that are asymptotically Kähler (but with the complex structure reversed — see the analysis of Figure 2 in Section 8). A closely related type of singularity formation, an asymptotic approach to the space of Kähler metrics in the blowup limit, has been partially investigated in [IKŠ19]. It should be noted that, while the blowdown soliton has been shown to occur as a model of singularity formation on compact manifolds by Máximo [Máx14], the solutions considered in that work are Kähler, whereas the unstable perturbations studied here all lie outside the space of Kähler initial data. On the other hand, the blowdown soliton is a smooth, complete, asymptotically conical, gradient shrinking Ricci soliton, and therefore arises as a Type-I singularity model of a *non-Kähler* Ricci flow on *some* closed manifold by recent work of Stolarski [Sto22]. The numerical findings in this paper suggest that the blowdown soliton models a Type-I singularity in the Ricci flow starting from an unstable perturbation of  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$ , which is consistent with Stolarski’s result.

In this paper, we describe numerical simulations that provide evidence in favor of the conjecture that there exists an unstable Ricci flow “orbit” starting arbitrarily close to  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$  and ending at the blowdown soliton  $\mathcal{L}_{-1}^2$  in real dimension  $n = 4$ . This is not a true orbit, because convergence to the blowdown soliton happens only locally and after parabolic dilation of the developing singularity. Nonetheless, it provides evidence in favor of the hierarchy of shrinking Ricci solitons outlined in [CHI04, Section 4]. In particular, the monotonicity of shrinker entropy  $\nu$  under Ricci flow implies that if the Ricci flow starting from an unstable perturbation of  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$  were to develop a different singularity, then this singularity would have higher density  $\Theta$  than  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$  [CHI04]. Since the blowdown soliton  $\mathcal{L}_{-1}^2$  is known to be unique among Kähler solitons [FIK03], our results have not detected any non-Kähler gradient shrinking soliton with the topology of  $\mathbb{C}^2$  blown up at a point, at least none with a density value above that of  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$ .

Because unperturbed  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$  is a positively-curved Einstein manifold, under Ricci flow it vanishes in a point. In particular, the area of the distinguished  $\mathbb{C}\mathbb{P}^1$  goes to zero at the same

---

<sup>1</sup> $N$  is the complex dimension, so that the real dimension is  $n = 2N$ .

time that the diameter of the manifold goes to zero. In each of our numerical simulations, we begin with initial data that are small non-Kähler conformal perturbations  $(\mathbb{C}\mathbb{P}^2, \tilde{G})$ . The conformal factor lies in the nullspace of the second variation of Perelman’s shrinker entropy  $\nu$  and is constructed in Section 2. Our numerics indicate that evolution of this initial data by Ricci–DeTurck flow crushes the  $\mathbb{C}\mathbb{P}^1$  fiber before the remainder of the manifold vanishes, that is, before the diameter goes to zero. Performing a parabolic rescaling of the solution fixes the area of the  $\mathbb{C}\mathbb{P}^1$  fiber. Our numerics indicate that the developing singularity is Type-I, see the analysis of Figure 1 in Section 8; that is, it develops at the natural parabolic rate. Because of this, one expects by [EMT11] to see local convergence of the parabolic blowups to a nonflat gradient shrinking soliton. How do we recognize the soliton numerically? We do so in two ways.

First, using a measure of local closeness to Kähler derived in Section 7 below, we find that the blowups are becoming asymptotically Kähler near the distinguished fiber, but with the opposite orientation of the complex structure on unperturbed  $(\mathbb{C}\mathbb{P}^2, G_{\text{FS}})$ . This is significant because we know by [FIK03] that the blowdown soliton is the unique gradient shrinking Kähler soliton with the topology of  $\mathbb{C}^2$  blown up at the origin, which corresponds in the original solution to the distinguished  $\mathbb{C}\mathbb{P}^1$ .

Second, we use the fact that the blowdown soliton is asymptotically conical. By [KW15], there is at most one shrinking gradient soliton asymptotic to any cone. So, as further verification, we determine that the parabolic blowup is converging numerically to the correct cone near the  $\mathbb{C}\mathbb{P}^1$ , in a neighborhood suitable to the rescaling. Thus, while numerical simulations cannot provide proof, we regard these two observations as compelling evidence in favor of the formation of the [FIK03] soliton in solutions emerging from unstable perturbations of the Fubini–Study metric.

## 2. AN UNSTABLE CONFORMAL PERTURBATION

We seek a conformal factor  $\psi$  so that  $(1 + \delta\psi)G_{\text{FS}}$  is an unstable perturbation of the Fubini–Study metric for small  $\delta > 0$ .

As shown in [KŠ19], the function  $\psi$  must satisfy three conditions:

$$\left(\Delta + \frac{1}{\tau}\right)\psi = 0, \quad \int_{\mathbb{C}\mathbb{P}^2} \psi \, dV = 0, \quad \int_{\mathbb{C}\mathbb{P}^2} \psi^3 \, dV > 0.$$

These are, respectively: the condition that  $\psi$  lie in the nullspace of the second variation of Perelman’s entropy  $\nu$ , a normalization necessary for that functional, and a sufficient condition for the perturbation generated by  $\psi$  to be unstable. Here,  $1/(2\tau)$  is the Einstein constant of the metric, which in our case is  $2(N + 1) = 6$ .

Because  $\text{Rc}(G_{\text{FS}}) = 6G_{\text{FS}}$  and the Laplacian of a rotationally-symmetric function  $\psi$  is

$$\Delta\psi = \psi_{\theta\theta} + \{3 \cot(\theta) - \tan(\theta)\}\psi_{\theta},$$

to specify rotationally symmetric unstable perturbations of the Fubini–Study metric, we seek solutions  $\psi$  of the second-order linear ODE

$$\begin{aligned} 0 &= (\Delta + 12)\psi(\theta) \\ &= \psi''(\theta) + \{3 \cot(\theta) - \tan(\theta)\}\psi'(\theta) + 12\psi(\theta). \end{aligned}$$

Only one fundamental solution of this ODE is smooth for  $\theta \in [0, \pi/2]$ . Up to an arbitrary multiplicative constant, it is

$$\psi = \csc^2(\theta) \{2 - 8 \cos(2\theta) + 6 \cos(4\theta)\} = 1 + 3 \cos(2\theta).$$

Using  $\det(G_{\text{FS}}) = \sin^6(\theta) \cos^2(\theta)$ , one readily verifies that this solution satisfies

$$\begin{aligned} \int_{\mathbb{CP}^2} \psi \, dV &= \int_0^{\pi/2} \{1 + 3 \cos(2\theta)\} \sin^3(\theta) \cos(\theta) \, d\theta = 0, \\ \int_{\mathbb{CP}^2} \psi^3 \, dV &= \int_0^{\pi/2} \{1 + 3 \cos(2\theta)\}^3 \sin^3(\theta) \cos(\theta) \, d\theta = \frac{2}{5}. \end{aligned}$$

Hence, given  $\delta > 0$ , we consider Ricci flow starting from  $(1 + \delta\psi)G_{\text{FS}}$ . After taking a suitable parabolic dilation of the flow, it suffices to consider initial data

$$\tilde{G} = \frac{1 + \delta\psi}{1 + \delta} G_{\text{FS}} = \left(1 + 3 \frac{\delta}{1 + \delta} \cos(2\theta)\right) G_{\text{FS}} = h G_{\text{FS}},$$

where  $h$  is defined in equation (7) below, with  $\varepsilon = \frac{\delta}{1 + \delta}$ .

### 3. COHOMOGENEITY-ONE METRICS

On  $[0, \pi/2] \times \text{SU}(2)$ , we consider cohomogeneity-one metrics of the form

$$(1) \quad G = \rho^2 \, d\theta^2 + \sum_{i=1}^3 f_i^2 \omega^i \otimes \omega^i,$$

where  $\{\omega^i \otimes \omega^i\}_{i=1}^3$  is a Milnor coframe for  $\text{SU}(2)$ . The cohomogeneity-one condition means that the fiber over a generic  $\theta$  is diffeomorphic to  $\mathbb{S}^3$ , except for  $\theta \in \{0, \pi/2\}$ . We are interested in initial data that are unstable conformal perturbations of the Fubini–Study metric on  $\mathbb{CP}^2$ . Accordingly, we impose an *Ansatz* of  $\text{U}(2)$  symmetry in the form

$$(2) \quad f_1 = f \quad \text{and} \quad f_2 = f_3 = g.$$

From the viewpoint of analysis, it is convenient to fix a gauge and let  $s(\theta, t)$  represent arclength with respect to  $G$ . Then  $s$  and  $\theta$  satisfy the infinitesimal spatial relation

$$ds = \rho \, d\theta.$$

In this gauge, the metric takes the form

$$G = ds^2 + \sum_{i=1}^3 f_i^2 \omega^i \otimes \omega^i.$$

Then the sectional curvatures of  $G$  are convex linear combinations of

$$(3a) \quad \kappa_{12} = \kappa_{31} = \frac{f^2}{g^4} - \frac{f_s g_s}{f g},$$

$$(3b) \quad \kappa_{23} = \frac{4g^2 - 3f^2}{g^4} - \frac{g_s^2}{g^2},$$

$$(3c) \quad \kappa_{01} = -\frac{f_{ss}}{f},$$

$$(3d) \quad \kappa_{02} = \kappa_{03} = -\frac{g_{ss}}{g},$$

where  $e_0 = \frac{d}{ds}$  and  $(e_1, e_2, e_3)$  is a Milnor frame for  $SU(2)$ .

The variables  $(s, t)$  do not commute, but this approach yields a strictly parabolic system for Ricci flow on these geometries,

$$(4a) \quad f_t = f_{ss} + 2\frac{g_s}{g}f_s - 2\frac{f^3}{g^4},$$

$$(4b) \quad g_t = g_{ss} + \left(\frac{f_s}{f} + \frac{g_s}{g}\right)g_s + 2\frac{f^2 - 2g^2}{g^3},$$

where all time derivatives on the LHS are taken at fixed  $\theta$ . This system has been studied in [IKŠ19]. One can recover the behavior of  $\rho$  under Ricci flow using its evolution equation

$$\rho_t = \left(\frac{f_{ss}}{f} + 2\frac{g_{ss}}{g}\right)\rho.$$

From the viewpoint of numerical simulation, on the other hand, it is preferable to use fixed commuting variables. Using the fact that any smooth  $\zeta(s)$  must satisfy

$$\zeta_s = \frac{\zeta_\theta}{\rho} \quad \text{and} \quad \zeta_{ss} = \frac{\zeta_{\theta\theta}}{\rho^2} - \frac{\rho_\theta \zeta_\theta}{\rho^3},$$

we rewrite system (4) in terms of the fixed commuting variables  $(\theta, t)$  as

$$(5a) \quad \rho_t = \frac{f_{\theta\theta}}{\rho f} - \frac{\rho_\theta f_\theta}{\rho^2 f} + 2\frac{g_{\theta\theta}}{\rho g} - 2\frac{\rho_\theta g_\theta}{\rho^2 g},$$

$$(5b) \quad f_t = \frac{f_{\theta\theta}}{\rho^2} - \frac{\rho_\theta f_\theta}{\rho^3} + 2\frac{f_\theta g_\theta}{\rho^2 g} - 2\frac{f^3}{g^4},$$

$$(5c) \quad g_t = \frac{g_{\theta\theta}}{\rho^2} - \frac{\rho_\theta g_\theta}{\rho^3} + \left(\frac{f_\theta}{f} + \frac{g_\theta}{g}\right)\frac{g_\theta}{\rho^2} + 2\frac{f^2 - 2g^2}{g^3}.$$

The evolution equation (5a) for  $\rho$  prevents this system from being parabolic. To remedy this, we implement DeTurck's trick below.

#### 4. INITIAL DATA

Our reference (unperturbed) Fubini–Study metric  $G_{\text{FS}}$  is determined by three functions  $\rho_{\text{FS}}, f_{\text{FS}}, g_{\text{FS}}: [0, \pi/2] \rightarrow \mathbb{R}$ , given by

$$\rho_{\text{FS}}(\theta) = 1, \quad f_{\text{FS}}(\theta) = \frac{1}{2}\sin(2\theta) = \sin(\theta)\cos(\theta), \quad \text{and} \quad g_{\text{FS}}(\theta) = \sin(\theta).$$

That is, using (1), we have

$$G_{\text{FS}} = d\theta^2 + \sin^2(\theta)\cos^2(\theta)\omega^1 \otimes \omega^1 + \sin^2(\theta)\omega^2 \otimes \omega^2 + \sin^2(\theta)\omega^3 \otimes \omega^3.$$

It is readily verified that these choices have the correct boundary behavior to induce a smooth metric on  $\mathbb{C}\mathbb{P}^2$ , with the geometry of an asymptotically round 4-ball near  $\theta = 0$  and with a distinguished  $\mathbb{C}\mathbb{P}^1$  fiber at  $\theta = \pi/2$ . Furthermore, it follows easily from (3) that  $\text{Rc}[G_{\text{FS}}] = 6G_{\text{FS}}$ .

For use in specifying a Ricci-DeTurck flow below, which shadows the Ricci flow and is parabolic, unlike the Ricci flow itself, we note that the Fubini–Study connection is determined by

$$(6) \quad (\Gamma_{\text{FS}})_{00}^0 = 0, \quad (\Gamma_{\text{FS}})_{11}^0 = -\frac{1}{4} \sin(4\theta), \quad (\Gamma_{\text{FS}})_{22}^0 = (\Gamma_{\text{FS}})_{33}^0 = -\frac{1}{2} \sin(2\theta).$$

Given any  $\varepsilon > 0$ , we recall that the conformal factor determining an unstable perturbation of the Fubini–Study metric is given by

$$(7) \quad h(\theta) := (1 + 3\varepsilon \cos(2\theta)),$$

as shown in Section 2. As initial data for our perturbed Ricci flow, we take  $G(0) = \tilde{G} = h G_{\text{FS}}$ , for which  $(\rho, f, g) = (h\rho_{\text{FS}}, hf_{\text{FS}}, hg_{\text{FS}})$ , namely

$$\begin{aligned} \rho(\theta) &= (1 + 3\varepsilon \cos(2\theta)), \\ f(\theta) &= \frac{1}{4} (2 \sin(2\theta) + 3\varepsilon \sin(4\theta)), \\ g(\theta) &= \sin(\theta) (1 + 3\varepsilon \cos(2\theta)) \end{aligned}$$

at time  $t = 0$ . The Levi–Civita connection of  $G(0)$  is determined by

$$\begin{aligned} \Gamma_{00}^0 &= -\frac{6\varepsilon \sin(2\theta)}{1 + 3\varepsilon \cos(2\theta)}, \\ \Gamma_{11}^0 &= -\frac{\sin(2\theta) (\cos(2\theta) + 3\varepsilon \cos(4\theta))}{2(1 + 3\varepsilon \cos(2\theta))}, \\ \Gamma_{22}^0 = \Gamma_{33}^0 &= -\frac{\sin(\theta) \cos(\theta) (1 - 6\varepsilon + 9\varepsilon \cos(2\theta))}{1 + 3\varepsilon \cos(2\theta)}, \end{aligned}$$

with the unlisted Levi-Civita coefficients vanishing.

## 5. RICCI-DETURCK FLOW

The formulas below that suppress time apply at all  $t \geq 0$  such that a solution exists.

We consider the PDE system

$$(8) \quad \partial_t G = -2 \text{Rc}[G] + \mathcal{L}_V G$$

with initial data  $G(0) = h G_{\text{FS}}$ , where our DeTurck vector field  $V$  is defined by

$$(9a) \quad V^\beta = G^{\alpha\alpha} \left( \Gamma_{\alpha\alpha}^\beta - (\Gamma_{\text{FS}})_{\alpha\alpha}^\beta \right),$$

in which the implicit summation ranges over  $\alpha \in \{0, 1, 2, 3\}$ . For the metrics under consideration, this reduces to  $V = V^0 \frac{d}{d\theta}$ , where

$$(9b) \quad V^0 = G^{\alpha\alpha} \left( \Gamma_{\alpha\alpha}^0 - (\Gamma_{\text{FS}})_{\alpha\alpha}^0 \right).$$

*Remark.* We note that at  $t = 0$ , one has

$$V^0(\theta, 0) = \frac{12\varepsilon \sin(2\theta)}{\left(1 + 3\varepsilon \cos(2\theta)\right)^3},$$

which is smooth for all sufficiently small values of  $\varepsilon \geq 0$ .

In working with the DeTurck vector field  $V = V^0 \frac{d}{d\theta}$  and the 1-form  $V^\flat = V_0 d\theta$ , where  $V_0 = \rho^2 V^0$ , it is convenient to relabel  $V^0$  as the function  $v(\theta, t)$  defined by

$$(10) \quad v = \frac{\rho_\theta}{\rho^3} + \frac{\frac{1}{4}\rho^2 \sin(4\theta) - f f_\theta}{\rho^2 f^2} + \frac{\rho^2 \sin(2\theta) - 2g g_\theta}{\rho^2 g^2}.$$

One finds that  $(\mathcal{L}_V G)_{\alpha\alpha} = 2\nabla_\alpha V_\alpha$ , where the only nonzero components are

$$\begin{aligned} \nabla_0 V_0 &= \rho^2 v_\theta + v \rho \rho_\theta = \rho(\rho v)_\theta, \\ \nabla_1 V_1 &= v f f_\theta, \\ \nabla_2 V_2 &= \nabla_3 V_3 = v g g_\theta. \end{aligned}$$

Thus from system (5) above, we obtain a manifestly parabolic system corresponding to the Ricci-DeTurck flow,

$$(11a) \quad \begin{aligned} \rho_t &= \frac{\rho_{\theta\theta}}{\rho^2} - 3\frac{\rho_\theta^2}{\rho^3} + \left(\frac{f_\theta}{\rho^2 f} + 2\frac{g_\theta}{\rho^2 g} + v\right)\rho_\theta \\ &\quad + \left(\left[\frac{\sin(4\theta)}{4f^2}\right]_\theta + \left[\frac{\sin(2\theta)}{g^2}\right]_\theta\right)\rho + \frac{f_\theta^2}{\rho f^2} + 2\frac{g_\theta^2}{\rho g^2}, \end{aligned}$$

$$(11b) \quad f_t = \frac{f_{\theta\theta}}{\rho^2} + \left(v - \frac{\rho_\theta}{\rho^3}\right)f_\theta + 2\frac{f_\theta g_\theta}{\rho^2 g} - 2\frac{f^3}{g^4},$$

$$(11c) \quad g_t = \frac{g_{\theta\theta}}{\rho^2} + \left(v - \frac{\rho_\theta}{\rho^3}\right)g_\theta + \left(\frac{f_\theta}{f} + \frac{g_\theta}{g}\right)\frac{g_\theta}{\rho^2} + 2\frac{f^2 - 2g^2}{g^3}.$$

Below, we convert this into an equivalent form for which suitable boundary behaviors as  $\theta \searrow 0$  and  $\theta \nearrow \pi/2$  can be robustly enforced in numerical simulations.

## 6. A SYSTEM BETTER SUITED TO SIMULATION

Here, we convert system (11) into an equivalent one appropriate for numerical simulation of the evolution. The issue we must address is this: for the evolving functions  $(\rho, f, g)$  to induce smooth metrics on  $\mathbb{C}\mathbb{P}^2$  at each time requires that we enforce the identities

$$f = 0, \quad f_s = 1, \quad \text{and} \quad g = 0, \quad g_s = 1 \quad \text{at} \quad \theta = 0,$$

and the identities

$$f = 0, \quad f_s = -1, \quad \text{and} \quad g > 0, \quad g_s = 0 \quad \text{at} \quad \theta = \frac{\pi}{2}.$$

However, we may impose only one Dirichlet or Neumann condition at each boundary for the functions that we evolve numerically. We proceed to resolve this issue in two steps, as follows:

We begin by defining

$$\tilde{\rho} := \log \rho, \quad \tilde{f} := \log f, \quad \tilde{g} := \log g,$$

and

$$Q := \rho^2 \sin(\theta) \cos(\theta) \left( \frac{\cos(2\theta)}{f^2} + \frac{2}{g^2} \right).$$

Then using (11) and diligently doing some algebra, we obtain the system:

$$(12a) \quad \rho^2 \tilde{\rho}_t = \tilde{\rho}_{\theta\theta} - Q\tilde{\rho}_\theta - \tilde{\rho}_\theta^2 + \tilde{f}_\theta^2 + 2\tilde{g}_\theta^2 + Q_\theta,$$

$$(12b) \quad \rho^2 \tilde{f}_t = \tilde{f}_{\theta\theta} + Q\tilde{f}_\theta - 2\frac{\rho^2}{g^4}f^2,$$

$$(12c) \quad \rho^2 \tilde{g}_t = \tilde{g}_{\theta\theta} + Q\tilde{g}_\theta + \frac{\rho^2}{g^4}(2f^2 - 4g^2).$$

Next we define  $A, Y, Z$  via the relations

$$\rho =: e^{A+Y+Z}, \quad f =: e^Y \cos(\theta) g, \quad g =: e^A \sin(\theta),$$

noting that

$$Q = 2e^{2Z} (\cot(2\theta) + e^{2Y} \cot(\theta)).$$

Then we obtain this system:

$$(13a) \quad \rho^2 A_t = A_{\theta\theta} + QA_\theta + \csc^2(\theta) (2e^{4Y+2Z} \cos^2(\theta) - 4e^{2Y+2Z} - 1) \\ + 2e^{2Z} \cot(\theta) (\cot(2\theta) + e^{2Y} \cot(\theta)),$$

$$(13b) \quad \rho^2 Y_t = Y_{\theta\theta} + QY_\theta - 2e^{2Z} \tan(\theta) \cot(2\theta) - 2e^{2Y+2Z} - \sec^2(\theta) \\ + 4e^{2Y+2Z} \csc^2(\theta) (1 - e^{2Y} \cos^2(\theta)),$$

$$(13c) \quad \rho^2 Z_t = Z_{\theta\theta} - Q(2A_\theta + 2Y_\theta + Z_\theta) + Q_\theta - (A_\theta + Y_\theta + Z_\theta)^2 + 3(A_\theta + \cot(\theta))^2 \\ + (Y_\theta - \tan(\theta))^2 + 2(A_\theta + \cot(\theta))(Y_\theta - \tan(\theta)) + 2e^{2Y+2Z} \\ + 2e^{2Y+2Z} \cot^2(\theta)(e^{2Y} - 1) + \sec^2(\theta) + \csc^2(\theta) + 2e^{2Z} \cot(2\theta) (\tan(\theta) - \cot(\theta)).$$

Now consider what boundary conditions we need to impose smoothness. We need  $A_\theta = 0$  at  $\theta = 0$  and  $\theta = \pi/2$ . We need  $Y = 0$  and  $Y_\theta = 0$  at  $\theta = 0$  and  $Y_\theta = 0$  at  $\theta = \pi/2$ . We need  $Z = 0$  and  $Z_\theta = 0$  at both  $\theta = 0$  and  $\theta = \pi/2$ . Thus,  $A$  is the sort of variable we need, but  $Y$  and  $Z$  are not. However, defining  $B$  and  $C$  by

$$Y =: B\sin^2(\theta), \quad Z =: C\sin^2(2\theta),$$

we see that the conditions needed for smoothness can be imposed as long as  $B_\theta = C_\theta = 0$  at  $\theta = \pi/2$ .

After some tedious algebra we find that equation (13b) yields

$$(14) \quad \partial_t B = \rho^{-2} \left[ B_{\theta\theta} + (Q + 4\cot(\theta)) B_\theta + B \left( \frac{2 + 2e^{2Z} - 4e^{2Y+2Z}}{\sin^2(\theta)} \right) \right. \\ \left. - 4B(1 + e^{2Z} + e^{2Y+2Z}) + e^{2Z} \left( \frac{-2 - 2e^{2Y} + 4e^{4Y}}{\sin^2(\theta)} \right) \right. \\ \left. + 4 \left( \frac{e^{2Z} - 1}{\sin^2(2\theta)} \right) + 4e^{2Y+2Z} \left( \frac{1 + 2Y - e^{2Y}}{\sin^4(\theta)} \right) \right].$$



Similarly some tedious algebra applied to equation (13c) yields the following:

$$\begin{aligned}
(15) \quad \partial_t C = & \rho^{-2} \left[ C_{\theta\theta} + (Q + 8 \cot(2\theta)) C_\theta - 8C(2 + e^{2Z} + e^{2Y+2Z}) \right. \\
& + \left( \frac{1 - e^{2Y+2Z}}{\sin^2(\theta)} \right) \left( \frac{2A_\theta}{\sin(2\theta)} - 4C \right) \\
& + \left( \frac{1 - e^{2Z}}{\sin^2(2\theta)} \right) (-8C + 4 \cot(2\theta)A_\theta + 2 \cos(2\theta) (\tan(\theta)B_\theta + 2B) - 6) \\
& + e^{2Y+2Z} \left( \frac{\cosh(2Y) - 1}{\sin^4(\theta)} \right) + 8(1 + \cos^2(\theta)) \left( \frac{1 + 2Z - e^{2Z}}{\sin^4(2\theta)} \right) + 2 \left( \frac{A_\theta}{\sin(2\theta)} \right)^2 \\
& \left. - (\sin(2\theta)C_\theta + 4 \cos(2\theta)C) \left( \sin(2\theta)C_\theta + 4 \cos(2\theta)C + \tan(\theta)B_\theta + 2B + \frac{2A_\theta}{\sin(2\theta)} \right) \right].
\end{aligned}$$

In summary, we evolve the variables  $A$ ,  $B$  and  $C$ , and their evolution equations are equations (13a), (14), and (15), respectively. The Neumann boundary conditions are that  $A_\theta$ ,  $B_\theta$  and  $C_\theta$  vanish at  $\theta = 0$  and  $\theta = \pi/2$ .

## 7. THE BLOWDOWN SOLITON

We first derive a necessary and sufficient condition for a metric of the form (1) to be Kähler, under the symmetry assumptions  $f_1 = f$  and  $f_2 = f_3 = g$ . Calabi observed [Cal82] that any  $U(2)$ -invariant Kähler metric on  $\mathbb{C}^2 \setminus (0,0)$  may be written in complex form as

$$(16) \quad G_{\mathbb{C}} = \left\{ e^{-r} \varphi \delta_{\alpha\beta} + e^{-2r} (\varphi_r - \varphi) \bar{z}_\alpha z_\beta \right\} dz^\alpha \otimes d\bar{z}^\beta,$$

with respect to his coordinate  $r := \log(|z_1|^2 + |z_2|^2)$ . Written in real coordinates, the same metric becomes

$$(17) \quad G_{\mathbb{R}} = \varphi_r \left( \frac{1}{4} dr \otimes dr + \omega^1 \otimes \omega^1 \right) + \varphi \left( \omega^2 \otimes \omega^2 + \omega^3 \otimes \omega^3 \right),$$

which shows that  $\varphi > 0$  and  $\varphi_r > 0$  are necessary conditions for this to be a Kähler metric.

A comparison of equations (1) and (17) shows that a coordinate transformation is needed to write a Kähler metric with respect to arclength  $s$ , where  $ds = \rho d\theta$ . To see this, we observe that if  $s$  and  $r$  are related by the ODE

$$\frac{dr}{ds} = \frac{2}{f},$$

then equation (1) takes the form

$$(18) \quad G = f^2 \left( \frac{1}{4} dr \otimes dr + \omega^1 \otimes \omega^1 \right) + g^2 \left( \omega^2 \otimes \omega^2 + \omega^3 \otimes \omega^3 \right),$$

which matches (17) if and only if  $f^2 = \varphi_r$  and  $g^2 = \varphi$  are related by

$$(19) \quad f = gg_s.$$

It follows that a cohomogeneity-one metric with the  $U(2)$  symmetries we have imposed is Kähler if and only if condition (19) holds. So, we may use the ratio  $f/(gg_s)$  to provide a local measurement of the closeness of a metric to the Kähler subspace.

Now, it follows from Lemma 6.1 and equation (27) of [FIK03] with  $\lambda = -1$ ,  $\mu = \sqrt{2}$ , and  $\nu = 0$ , that the metric on the blowdown soliton  $\mathcal{L}_{-1}^2$  is determined by a function  $\phi(r)$  that solves the separable first-order ODE

$$(20) \quad \phi_r = \frac{1}{\sqrt{2}}\phi - (\sqrt{2} - 1) - \left(1 - \frac{1}{\sqrt{2}}\right)\phi^{-1}.$$

One can solve (20) implicitly up to an arbitrary constant  $\eta$ , obtaining

$$(21) \quad e^{r+\eta} = \frac{\phi - 1}{(\phi + \sqrt{2} - 1)^{\sqrt{2}-1}}.$$

As shown in [FIK03], the  $\mathcal{L}_{-1}^2$  soliton is complete and exists for all  $r \in \mathbb{R}$ . Using this fact and the positivity of  $\phi$  and  $\phi_r$ , it is not difficult to see from (20) and (21) that  $\phi \nearrow \infty$  as  $r \nearrow \infty$  and that the metric is asymptotically conical in the precise sense that

$$(22) \quad \frac{\phi_r}{\phi} \rightarrow \frac{1}{\sqrt{2}} \quad \text{as} \quad \phi \nearrow \infty.$$

Specifically, if we define  $\gamma := 2^{-1/4}$ , then the asymptotic cone of the blowdown soliton corresponds to

$$(23) \quad f = \gamma^2 s \quad \text{and} \quad g = \gamma s \quad \Rightarrow \quad \frac{f^2}{g^2} = \gamma^2 = \frac{1}{\sqrt{2}}.$$

All sectional curvatures of the cone vanish except  $\kappa_{23} = \frac{4(\sqrt{2}-1)}{s^2}$ .

## 8. RESULTS OF SIMULATIONS

The simulations are performed using standard numerical methods for parabolic equations: centered differences for spatial derivatives and Euler's method for time evolution. We describe the method here in more detail.

Any function  $F(t, \theta)$  is represented by the values  $F_i^k$  that the function takes at points  $\theta_i$  equally spaced with spacing  $\Delta\theta$  and with times  $t_k$  equally spaced with spacing  $\Delta t$ . We use standard centered finite differences, so that  $F_\theta$  and  $F_{\theta\theta}$  are approximated by

$$(24) \quad F_\theta = \frac{F_{i+1}^k - F_{i-1}^k}{2\Delta\theta}$$

and

$$(25) \quad F_{\theta\theta} = \frac{F_{i+1}^k + F_{i-1}^k - 2F_i^k}{(\Delta\theta)^2},$$

respectively.

Time evolution is carried out using the Euler method, so that

$$(26) \quad F_i^{k+1} = F_i^k + \Delta t \partial_t F,$$

where  $\partial_t F$  is the finite difference version of the right hand side of equation (13a), (14) or (15) with the spatial derivatives evaluated using equations (24) and (25). The standard von Neumann stability analysis of equation (26) reveals that the time step must satisfy the Courant condition

$$(27) \quad \Delta t < \frac{1}{2}(\rho_{\min}\Delta\theta)^2.$$

where  $\rho_{\min}$  is the minimum value of  $\rho$ .

We now show the results of a simulation with  $\varepsilon = 0.1$  in equation (7). To get an idea of how the evolution proceeds we plot  $1/\kappa_{23}$  at  $\theta = \pi/2$  as a function of time. This curve goes to zero (i.e.  $\kappa_{23} \rightarrow \infty$ ) at the final time. We note that the graph is linear near the final time, which suggests that the developing singularity is Type-I. Thus  $1/\kappa_{23}$  is a good proxy (up to some overall scale) for time remaining until the singularity.




FIGURE 1.  $1/\kappa_{23}$  plotted vs. time

We now want to examine the extent to which the metric becomes Kähler as the singularity is approached. Recall that a Kähler metric has  $f = gg_s$ . We define the quantity  $K$  by  $K = gg_s/f$ . Then a metric is Kähler if  $K = 1$ , but it is also Kähler (but with the opposite orientation of  $s$ ) if  $K = -1$ . In Figure 2, we plot  $K$  vs.  $\theta$  at a time near the final time. Note that near  $\theta = \pi/2$  the quantity  $K$  is approaching  $-1$ , thus indicating that the metric in this region is becoming Kähler.




FIGURE 2.  $K = gg_s/f$  plotted vs.  $\theta$  near the final time

We would now like to know the form of the Kähler metric that is being approached. Here we take a small region near  $\theta = \pi/2$  and examine the behavior of the cone angle  $\gamma^2 = f^2/g^2$ . In Figure 3, we plot  $\gamma^2 = f^2/g^2$  vs. the length from  $\pi/2$  for three times near the final time. We see that  $\gamma^2$  approaches  $0.707 \approx 1/\sqrt{2}$  numerically, indicating local convergence of the metric cone angle  $\gamma^2$  to that of the blowdown soliton; *cf.* equation (23). What is perhaps surprising is how close one needs to get to the singularity to obtain this asymptotic behavior. In figure (3) the bottom curve corresponds to  $1/\kappa_{23} = 4.7 \times 10^{-8}$  and has an asymptotic value for  $f^2/g^2$  of 0.666. The middle curve has  $1/\kappa_{23} = 9.4 \times 10^{-9}$  and has an asymptotic value for  $f^2/g^2$  of 0.700. The top curve has  $1/\kappa_{23} = 1.9 \times 10^{-9}$  and has an asymptotic value for  $f^2/g^2$  of 0.707.




FIGURE 3.  $\gamma^2 = f^2/g^2$  plotted vs. length near the final time

## 9. DIRECTIONS FOR FUTURE WORK

Our results here provide evidence in favor of the conjecture that unstable perturbations of the Fubini–Study Einstein metric — perturbations which are conformal and not Kähler — develop finite-time local singularities modeled by the Kähler blowdown soliton  $\mathcal{L}_{-1}^2$  discovered in [FIK03], but with the opposite complex structure.

Short of proving the full conjecture, a useful next step would be to derive formal matched asymptotics that describe how parabolic dilations at these singularities approach  $\mathcal{L}_{-1}^2$ , analogous to the matched asymptotics formally derived in [AIK11] for solutions that approach the Bryant soliton. However, because  $\mathcal{L}_{-1}^2$  is only known in the implicit form (21), a somewhat more approachable next step would be to develop formal matched asymptotics for parabolic dilations that approach its asymptotic cone  $(f, g) = (\gamma^2 s, \gamma s)$ , where  $\gamma = 2^{-1/4}$ , as described in (23).

## REFERENCES

- [AIK11] Sigurd B. Angenent, James Isenberg, and Dan Knopf, *Formal matched asymptotics for degenerate Ricci flow neckpinches*, *Nonlinearity* **24** (2011), no. 8, 2265–2280.
- [Cal82] Eugenio Calabi, *Extremal Kähler metrics*, *Seminar on Differential Geometry*, 1982, pp. 259–290.
- [CHI04] Huai-Dong Cao, Richard S. Hamilton, and Tom Ilmanen, *Gaussian densities and stability for some Ricci solitons* (2004), available at [arXiv:0404165v1](https://arxiv.org/abs/0404165v1).
- [EMT11] Joerg Enders, Reto Müller, and Peter M. Topping, *On type-I singularities in Ricci flow*, *Comm. Anal. Geom.* **19** (2011), no. 5, 905–922.

- [FIK03] Mikhail Feldman, Tom Ilmanen, and Dan Knopf, *Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons*, J. Differential Geom. **65** (2003), no. 2, 169–209.
- [Ham95] Richard S. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in differential geometry, Vol. II (Cambridge, MA, 1993), 1995, pp. 7–136.
- [IKŠ19] James Isenberg, Dan Knopf, and Nataša Šešum, *Non-Kähler Ricci flow singularities modeled on Kähler-Ricci solitons*, Pure Appl. Math. Q. **15** (2019), no. 2, 749–784.
- [Krö20] Klaus Kröncke, *Stability of Einstein metrics under Ricci flow*, Comm. Anal. Geom. **28** (2020), no. 2, 351–394.
- [KŠ19] Dan Knopf and Nataša Šešum, *Dynamic instability of  $\mathbb{C}\mathbb{P}^N$  under Ricci flow*, J. Geom. Anal. **29** (2019), no. 1, 902–916.
- [KW15] Brett Kotschwar and Lu Wang, *Rigidity of asymptotically conical shrinking gradient Ricci solitons*, J. Differential Geom. **100** (2015), no. 1, 55–108.
- [Máx14] Davi Máximo, *On the blow-up of four-dimensional Ricci flow singularities*, J. Reine Angew. Math. **692** (2014), 153–171.
- [Sto22] Maxwell Stolarski, *Closed Ricci flows with singularities modeled on asymptotically conical shrinkers* (2022), available at [arXiv:2202.03386v2](https://arxiv.org/abs/2202.03386v2).

(David Garfinkle) OAKLAND UNIVERSITY

*Email address:* [garfinkl@oakland.edu](mailto:garfinkl@oakland.edu)

(James Isenberg) UNIVERSITY OF OREGON

*Email address:* [isenberg@uoregon.edu](mailto:isenberg@uoregon.edu)

*URL:* <http://www.uoregon.edu/~isenberg/>

(Dan Knopf) UNIVERSITY OF TEXAS AT AUSTIN

*Email address:* [danknopf@math.utexas.edu](mailto:danknopf@math.utexas.edu)

*URL:* <http://www.ma.utexas.edu/users/danknopf>

(Haotian Wu) THE UNIVERSITY OF SYDNEY

*Email address:* [haotian.wu@sydney.edu.au](mailto:haotian.wu@sydney.edu.au)