

Differential Equations with Linear Algebra (M427J)

Some answers to homework # 4

Fall 2025

Prof. Hector E. Lomeli

§2.5 # 2. There are two cases: $\alpha = -2$ and $\alpha \neq 2$.

- If $\alpha = -2$, then $\Psi(t) = \frac{1}{6}e^{-2t}t^3$.
- If $\alpha \neq 2$, then

$$\Psi(t) = \frac{e^{\alpha t}(\alpha t + 2t - 2)}{(\alpha + 2)^3}.$$

§2.5 # 4. $\Psi(t) = t^2 - t$.

§2.5 # 6. $\Psi(t) = \frac{(273 - 1672t + 3872t^2)e^{7t}}{340736}$.

§2.5 # 7. The characteristic equation is $r^2 + 4 = 0$. We have two complex roots: $2i$ and $-2i$. The right hand side of the non-homogeneous equation is of the form

$$g(t) = t \sin(2t) = P(t)e^{\alpha t} \cos(\beta t) + Q(t)e^{\alpha t} \sin(\beta t).$$

We will use $\alpha = 0$ and $\beta = 2$, and we have that $\alpha + i\beta$ is a single root, so we will use $s = 1$. Also, $P(t)$ is a polynomial of degree 0 and $Q(t)$ is a polynomial of degree 1. We will use $n = 1$.

The considerations above imply that the form of the function $\Psi(t)$ is

$$\begin{aligned}\Psi(t) &= t^s [(A_0 + A_1 t) e^{\alpha t} \cos(\beta t) + (B_0 + B_1 t) e^{\alpha t} \sin(\beta t)] \\ &= (A_0 t + A_1 t^2) \cos(2t) + (B_0 t + B_1 t^2) \sin(2t).\end{aligned}$$

Computing the linear operator, we find that

$$L[\Psi] = \Psi''(t) + 4\Psi(t) = 2(A_1 + 2B_0) \cos(2t) + (8B_1)t \cos(2t) + 2(B_1 - 2A_0) \sin(2t) + (-8A_1)t \sin(2t).$$

Matching coefficients, we get the equations

$$A_1 + 2B_0 = 0, \quad 8B_1 = 0, \quad B_1 - 2A_0 = 0, \quad -8A_1 = 1.$$

Solving these equations we find that

$$A_0 = 0, \quad A_1 = -\frac{1}{8}, \quad B_0 = \frac{1}{16}, \quad B_1 = 0.$$

We conclude that $\Psi(t) = \frac{1}{16}t \sin(2t) - \frac{1}{8}t^2 \cos(2t)$.

§2.5 # 14. $\Psi(t) = \frac{t^3}{6} - \frac{t^2}{4} + \frac{3t}{4} - \frac{1}{2}t e^{-2t}$.