

MORE HYPOTHESIS TESTING FOR TWO-WAY ANOVA

What do we do after testing for interaction?

This depends on whether or not interaction is significant (statistically or otherwise) *and* on what the original questions were in designing the experiment *and* on whether or not the analyzer wishes to engage in data-snooping *and* on the context of the experiment. We will spend a while discussing this.

I. If we *reject* H_0^{AB} (i.e., assume there *is* interaction) then it is usually inappropriate to test for main effects (that is, the contributions of the two factors A and B separately), since the question of what a “main effect” is in the presence of interaction is unclear. (How can you “separate out” the effect of A from the interaction if there is interaction?) Instead, it is usually preferable to use the equivalent cell-means model to examine contrasts in the treatment combinations.

II. If we *do not reject* H_0^{AB} (i.e., decide there is *no* interaction), then we are usually interested in main effects. These can be tested within the complete model. *Staying with this model is advisable rather than switching to the inequivalent main-effects model.*

Testing the contribution of each factor in the complete model (equal sample sizes)

Note: We are still assuming equal sample sizes.

We wish to test whether or not the factor A is needed in the model. Recall that the model states

$$Y_{ijt} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + \varepsilon_{ijt}$$

A occurs in the model through the terms α_i and $(\alpha\beta)_{ij}$. So “A is not needed in the model” means that the contribution of these two terms is independent of the level of A. That is,

$$\alpha_i + (\alpha\beta)_{ij} = \alpha_s + (\alpha\beta)_{sj} \text{ for all } i, s, \text{ and } j.$$

Thus, the null hypothesis “A is not needed in the model” could be stated as

$$H_0: \alpha_i + (\alpha\beta)_{ij} = \alpha_s + (\alpha\beta)_{sj} \text{ for all } i, s, \text{ and } j$$

with alternate hypothesis

$$H_a: \alpha_i + (\alpha\beta)_{ij} \neq \alpha_s + (\alpha\beta)_{sj} \text{ for at least one combination } i, s, \text{ and } j$$

The textbook does not explicitly mention this H_0 . Instead, it lists two possible null hypotheses. The first is

$$H_0^A: \alpha_1^* = \alpha_2^* = \dots = \alpha_a^*$$

(with H_a^A : At least two of the α_i^* 's are different),

where $\alpha_i^* = \alpha_i + (\overline{\alpha\beta})_{i\cdot}$. That is, the test is whether or not the levels of A, averaged over the levels of B, have the same average effect on the response. (Recall that the α_i^* 's occurred previously in the notes Analysis of Variance for the Two-Way Complete Model.)

The second null hypothesis mentioned in the textbook is

$$H_0^{A+AB}: H_0^A \text{ and } H_0^{AB} \text{ are both true.}$$

What are the connections between these three possible null hypotheses?

i) Clearly, H_0^{A+AB} implies H_0^A .

ii) The following calculations show that H_0 implies H_0^{A+AB} :

If H_0 is true, then $\alpha_i + (\alpha\beta)_{ij} = \alpha_s + (\alpha\beta)_{sj}$ for all i, s , and j . Averaging over the subscript j gives

$$\alpha_i + (\overline{\alpha\beta})_{i\cdot} = \alpha_s + (\overline{\alpha\beta})_{s\cdot} \text{ for all } i \text{ and } s,$$

which says H^A is true.

Subtracting this from the original equation,

$$(\alpha\beta)_{ij} - (\overline{\alpha\beta})_{i\cdot} = (\alpha\beta)_{sj} - (\overline{\alpha\beta})_{s\cdot} \text{ for all } i, j, \text{ and } s.$$

Rearranging,

$$(\alpha\beta)_{ij} - (\alpha\beta)_{sj} = (\overline{\alpha\beta})_{i\cdot} - (\overline{\alpha\beta})_{s\cdot}.$$

But the right side is independent of j , so we conclude

$$(\alpha\beta)_{ij} - (\alpha\beta)_{sj} = (\alpha\beta)_{iq} - (\alpha\beta)_{sq} \text{ for all } i, s, j, \text{ and } q,$$

which says there is no interaction.

iii) The following shows that H_0^{A+AB} implies H_0 :

If H_0^{A+AB} is true, then so is H_0^{AB} , so

$$(\alpha\beta)_{ij} - (\alpha\beta)_{sj} = (\alpha\beta)_{iq} - (\alpha\beta)_{sq} \text{ for all } i, s, j, \text{ and } q.$$

Averaging over q and rearranging,

$$(\alpha\beta)_{ij} - (\overline{\alpha\beta})_{i\cdot} = (\alpha\beta)_{sj} - (\overline{\alpha\beta})_{s\cdot} \text{ for all } i, j, \text{ and } s.$$

Add this to the equation for H^A to get

$$\alpha_i + (\alpha\beta)_{ij} = \alpha_s + (\alpha\beta)_{sj} \text{ for all } i, j, \text{ and } s,$$

which says H_0 is true.

iv) Does H_0^A imply H_0^{A+AB} (equivalently, H_0)?

No. Consider the example where $\alpha_1 = \alpha_2 = \beta_1 = \beta_2 = 0$, $(\alpha\beta)_{11} = (\alpha\beta)_{22} = 0$, $(\alpha\beta)_{12} = (\alpha\beta)_{21} = 1$. Thus

$$Y_{11} = \varepsilon_{11}, \quad \overline{Y}_{12} = 1 + \varepsilon_{12}, \quad Y_{21} = 1 + \varepsilon_{21}, \quad Y_{22} = \varepsilon_{22}$$

Then $\alpha_1^* = \alpha_1 + (\overline{\alpha\beta})_{1\cdot} = 0 + (0 + 1)/2 = 1/2$ and $\alpha_2^* = \alpha_2 + (\overline{\alpha\beta})_{2\cdot} = 0 + (1+0)/2 = 1/2$, so H^A is true. But H^{AB} is not true. (Draw an interaction plot!)

The test for H_0^A is the one that is automatic in most software. *We will take the perspective that it does not make sense to test for a main effect of A unless there is no interaction*, so using this test will not cause problems. (But if you ever see a paper that tests for “main effects” when there is interaction, be cautious in the interpretation. Do *not* interpret the null hypothesis as saying “A has no effect;” it just means that “the levels of A, averaged over the levels of B, have the same average effect on the response.”)

To test H_0^A , we will again use an F test comparing the full model with a reduced model: the one where H_0^A is true. If sample sizes are equal, it can be shown that the least squares estimate of $E[Y_{ijt}]$ under this new reduced model (i.e, under H_0^A) is

$$\overline{y}_{ij\cdot} - \overline{y}_{i\cdot\cdot} + \overline{y}_{\cdot\cdot\cdot},$$

giving sum of squares for the reduced model

$$ssE_0^A = \sum_i \sum_j \sum_t (y_{ijt} - \bar{y}_{ij.} + \bar{y}_{i..} - \bar{y}_{...})^2,$$

which by appropriate algebraic manipulations becomes

$$\begin{aligned} ssE_0^A &= \sum_i \sum_j \sum_t (y_{ijt} - \bar{y}_{ij.})^2 + br \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2 \\ &= ssE + br \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2, \end{aligned}$$

so the *sum of squares for treatment factor A* is

$$\begin{aligned} ssA &= ssE_0^A - ssE \\ &= br \sum_{i=1}^a (\bar{y}_{i..} - \bar{y}_{...})^2 \\ &= (1/br) \sum_{i=1}^a (y_{i..})^2 - (y_{...})^2/abr, \end{aligned}$$

which resembles the formula for ssT used to test equality of effects in one-way analysis of variance. The reasoning behind the test used is: If H_0^A is true, then ssA should be small compared to ssE , so we will have evidence lending doubt to H_0^A if ssA/ssE is unusually large.

If SSA is the random variable corresponding to ssA , it can be shown that when H_0^A is true and sample sizes are equal,

- i) $SSA/\sigma^2 \sim \chi^2(a-1)$
- ii) SSA and SSE are independent.

Thus, when sample sizes are equal and H_0^A is true,

$$\frac{SSA/(a-1)\sigma^2}{SSE/(n-ab)\sigma^2} = \frac{MSA}{MSE} \sim F(a-1, n-ab)$$

Since msA/msE is just a scalar multiple of the ratio ssA/ssE , we can use msA/msE as a test statistic, rejecting for large values.

Similarly, we can form the *sum of squares for treatment factor B* and obtain an F-test based on

$$\frac{SSB/(b-1)\sigma^2}{SSE/(n-ab)\sigma^2} = \frac{MSB}{MSE} \sim F(b-1, n-ab)$$

for

- H_0^B : Every β_j and every $(\alpha\beta)_{ij} = 0$
 against the alternate hypothesis
 H_a^B : At least one of the β_j 's or $(\alpha\beta)_{ij}$'s is not zero.

Analysis of Variance Table

For each of the three tests (for interaction, effect of A and effect of B), we have a corresponding sum of squares, $ssAB$, ssA , and ssB . We also have the error sum of squares, ssE . If we add up the formulas for these three sums of squares and do appropriate algebraic manipulations, we will get

$$ssA + ssB + ssAB + ssE = \sum_i \sum_j \sum_t (y_{ijt} - \bar{y}_{...})^2.$$

This last sum of squares is called the *total sum of squares*, denoted ssT . It can be seen as a measure of the total variability of the data without taking into account either A or B. Similarly, ssE is a measure of the variability taking into account A, B and their interaction; ssA is a measure of the variability taking B into account but not A, and ssB is a measure of the variability taking A into account but not B.

The sums of squares and the additional information used in the tests for A, B and AB are traditionally summarized in an *Analysis of Variance Table* with one line each for A, B, AB, error, and "total sum of squares"

$$sstot = ssA + ssB + ssAB + ssE$$

Interpreting ANOVA tests

Interpretation requires thought -- we need to take into account the purpose of the study, the context, multiple comparisons, and whether or not we are willing to do data snooping. Interpretation can sometimes be frustrating -- for example, what if the test for interaction is significant, but the test for one of the factors is not?

Examples: Battery and reaction time.

Note: When sample sizes are unequal, the formulae for the sums of squares are more complicated, and the corresponding random variables are not independent. More on this later.