

## INFERENCE FOR SIMPLE OLS

### *Model Assumptions ("The" Simple Linear Regression Model Version IV):*

(We consider  $x_1, \dots, x_n$  as fixed.)

1.  $E(Y|x) = \eta_0 + \eta_1 x$  (linear mean function)
2.  $\text{Var}(Y|x) = \sigma^2$  (Equivalently,  $\text{Var}(e|x) = \sigma^2$ ) (constant variance)
3.  $y_1, \dots, y_n$  are independent observations. (independence)
4. (NEW)  $Y|x$  is normal for each  $x$  (normality)

(1) + (2) + (4) can be summarized as:

$$Y|x \sim N(\eta_0 + \eta_1 x, \sigma^2)$$

Recall:  $e|x = Y|x - E(Y|x)$

So:  $e|x \sim N(0, \sigma^2)$

i.e., all errors have the same distribution -- so we just say  $e$  instead of  $e|x$ .

Since  $\hat{\eta}_0$  and  $\hat{\eta}_1$  are linear combinations of the  $Y|x_i$ 's, (3) + (4) imply that  $\hat{\eta}_0$  and  $\hat{\eta}_1$  are normally distributed random variables (that is, their sampling distributions are normal).

Recalling that

$$E(\hat{\eta}_1) = \eta_1 \quad \text{Var}(\hat{\eta}_1) = \frac{\sigma^2}{SXX} \quad E(\hat{\eta}_0) = \eta_0 \quad \text{Var}(\hat{\eta}_0) = \sigma^2 \left( \frac{1}{n} + \frac{\bar{x}^2}{SXX} \right),$$

We have

$$\hat{\eta}_1 \sim \hat{\eta}_0 \sim$$

Look more at  $\hat{\eta}_1$ : We can standardize to get

$$\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\sigma^2/SXX}} \sim N(0,1)$$

But we don't know  $\sigma^2$ , so need to approximate it by  $\hat{\sigma}^2$  -- in other words approximate

$\text{Var}(\hat{\eta}_1)$  by  $\hat{\text{Var}}(\hat{\eta}_1) = [\text{s.e.}(\hat{\eta}_1)]^2 = \frac{\hat{\sigma}^2}{SXX}$ . Thus we want to use  $\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2/SXX}}$ . But we can't

expect this to be normal, too. However,

$$\begin{aligned}
 & \frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2 / SXX}} = \\
 (*) & \frac{\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\sigma^2 / SXX}}}{\sqrt{\hat{\sigma}^2 / \sigma^2}}
 \end{aligned}$$

The numerator of the last fraction *is* normal (in fact, standard normal), as noted above.

Facts: (Proofs omitted)

a.  $(n-2) \frac{\hat{\sigma}^2}{\sigma^2}$  has a  $\chi^2$  distribution with  $n-2$  degrees of freedom

$$\text{Notation: } (n-2) \frac{\hat{\sigma}^2}{\sigma^2} \sim \chi^2(n-2)$$

b.  $(n-2) \frac{\hat{\sigma}^2}{\sigma^2}$  is independent of  $\hat{\eta}_1 - \eta_1$  (hence independent of the numerator in (\*))

### Comments on distributions:

1. A  $\chi^2(k)$  distribution is defined as the distribution of a random variable which is a sum of squares of  $k$  independent standard normal random variables.

[Comment: Recall that  $\hat{\sigma}^2 = \frac{1}{n-2} RSS$ , so  $(n-2) \frac{\hat{\sigma}^2}{\sigma^2} = \frac{RSS}{\sigma^2} = \sum \left( \frac{\hat{\epsilon}_i}{\sigma} \right)^2$  is a sum of  $n$

squares; the fact quoted above says that it can also be expressed as a sum of  $n-2$  squares of *independent standard normal* random variables.]

2. A  $t$ -distribution with  $k$  degrees of freedom is defined as the distribution of a random variable of the form  $\frac{Z}{\sqrt{U/k}}$  where

- $Z \sim N(0,1)$
- $U \sim \chi^2(k)$
- $Z$  and  $U$  are independent.

In the fraction (\*) above, take

$$\begin{aligned}
 U &= (n-2) \frac{\hat{\sigma}^2}{\sigma^2} \sim \chi^2(n-2) \\
 Z &= \frac{\hat{\eta}_1 - \eta_1}{\sqrt{\sigma^2 / SXX}} \sim N(0,1)
 \end{aligned}$$

Thus: 
$$\frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2 / SXX}} \sim t(n-2),$$

so we can do inference on  $\eta_1$ , using  $t = \frac{\hat{\eta}_1 - \eta_1}{\sqrt{\hat{\sigma}^2 / SXX}}$  as our test statistic.

### **Inference on $\eta_0$**

With the same assumptions, it can be shown in an analogous manner (details omitted) that

$$\frac{\hat{\eta}_0 - \eta_0}{s.e.(\hat{\eta}_0)} \sim t(n-2),$$

so we can use this statistic to do inference on  $\eta_0$ .