

NOTES FOR SUMMER STATISTICS INSTITUTE COURSE

**COMMON MISTAKES IN STATISTICS –
SPOTTING THEM AND AVOIDING THEM**

Day 2: Important Details about Statistical Inference

“The devil is in the details” (anonymous)

MAY 20 – 23, 2013

Instructor: Martha K. Smith

CONTENTS OF DAY 2

I. More Precise Definition of Simple Random Sample	3
Connection with independent random variables	4
Problems with small populations	9
II. Why Random Sampling is Important	10
A myth, an urban legend, and the real reason	
III. Overview of Frequentist Hypothesis Testing	11
Basic elements of most frequentist hypothesis tests	11
Illustration: Large sample z-test	13
Common confusions in terminology and concepts	14
Sampling distribution	16
Role of model assumptions	20
Importance of model assumptions in general	23
Robustness	24
IV. Frequentist Confidence Intervals	25
Point vs interval estimates	25
Illustration: Large sample z-procedure	27 -
Cautions and common confusions	33, 34, 36, 40
Importance of model assumptions	26, 27, 29, 30, 32, 35, 37
Robustness	37
Variations and trade-offs	38, 39
V. More on Frequentist Hypothesis Tests	41
Illustration: One-sided t-test for a sample mean	42
p-values	45
VI. Misinterpretations and Misuses of p-values	49
VII. Type I error and significance levels	55
VIII. Pros and cons of setting a significance level	58

I. MORE PRECISE DEFINITION OF SIMPLE RANDOM SAMPLE

In practice in applying statistical techniques, we're interested in *random variables* defined on the population under study.

Recall the examples mentioned yesterday:

1. In a medical study, the population might be all adults over age 50 who have high blood pressure.
2. In another study, the population might be all hospitals in the U.S. that perform heart bypass surgery.
3. If we're studying whether a certain die is fair or weighted, the population is all possible tosses of the die.

In these examples, we might be interested in the following random variables:

Example 1: The difference in blood pressure with and without taking a certain drug.

Example 2: The number of heart bypass surgeries performed in a particular year, or the number of such surgeries that are successful, or the number in which the patient has complications of surgery, etc.

Example 3: The number that comes up on the die.

Connection with Independent Random Variables:

If we take a *sample of units from the population*, we have a corresponding *sample of values of the random variable*.

In Example 1:

- The *random variable* is “difference in blood pressure with and without taking the drug.”
 - Call this random variable Y (upper case Y)
- The sample of *units from the population* is a sample of adults over age 50 who have high blood pressure.
 - Call them person 1, person 2, etc.
- The corresponding sample of values of the random variable will consist of values we will call y_1, y_2, \dots, y_n (lower case y 's), where
 - n = number of people in the sample
 - y_1 = the difference in blood pressures (that is, the value of Y) for the first person in the sample,
 - y_2 = the difference in blood pressures (that is, the value of Y) for the second person in the sample
 - etc.

We can look at this another way, in terms of n random variables Y_1, Y_2, \dots, Y_n , described as follows:

- The random process for Y_1 is “pick the first person in the sample”; the value of Y_1 is the value of Y for that person – i.e., y_1 .
- The random process for Y_2 is “pick the second person in the sample”; the value of Y_2 is the value of Y for that person – i.e., y_2 .
- etc.

The difference between using the small y 's and the large Y 's is that when we use the small y 's we are thinking of a fixed sample of size n from the population, but when we use the large Y 's, we are thinking of letting the sample vary (but always with size n).

Note: The Y_i 's are sometimes called *identically distributed*, because they have the same probability distribution (in this example, the distribution of Y).

Precise definition of simple random sample of a random variable:

"The sample y_1, y_2, \dots, y_n is a simple random sample" means that the associated random variables Y_1, Y_2, \dots, Y_n are independent.

Intuitively speaking, "*independent*" means that *the values of any subset of the random variables Y_1, Y_2, \dots, Y_n do not influence the values of the other random variables in the list.*

Recall: We defined a random sample as one that is chosen by a random process.

- Where is the random process in the precise definition?

Note: To emphasize that the Y_i 's all have the same distribution, the precise definition is sometimes stated as, “ Y_1, Y_2, \dots, Y_n are independent, identically distributed,” sometimes abbreviated as *iid*.

Connection with the initial definition of simple random sample

Recall the preliminary definition (from Moore and McCabe, *Introduction to the Practice of Statistics*) given in Simple Random Samples, Part 1:

"A simple random sample (SRS) of size n consists of n individuals from the population chosen in such a way that every set of n individuals has an equal chance to be the sample actually selected."

Recall Example 3 above: We toss a die; the number that comes up on the die is the value of our random variable Y .

- In terms of the preliminary definition:
 - The population is all possible tosses of the die.
 - A simple random sample is n different tosses.
- The different tosses of the die are *independent events* (i.e., what happens in some tosses has no influence on the other tosses), which means that in the precise definition above, the random variables Y_1, Y_2, \dots, Y_n are indeed independent: The numbers that come up in some tosses in no way influence the numbers that come up in other tosses.

Compare this with example 2: The population is all hospitals in the U.S. that perform heart bypass surgery.

- Using the preliminary definition of simple random sample of size n , we end up with n *distinct* hospitals.
- This means that when we have chosen the first hospital in our simple random sample, *we cannot choose it again to be in our simple random sample*.
- Thus the events "Choose the first hospital in the sample; choose the second hospital in the sample; ...," are *not* independent events: The choice of first hospital restricts the choice of the second and subsequent hospitals in the sample.
- If we now consider the random variable Y = the number of heart bypass surgeries performed in 2008, then it follows that the random variables Y_1, Y_2, \dots, Y_n are *not* independent.

The Bottom Line: In many cases, the preliminary definition does *not* coincide with the more precise definition.

More specifically, the preliminary definition allows sampling *without* replacement, whereas the more precise definition requires sampling *with* replacement.

The Bad News: *The precise definition is the one used in the mathematical theorems that justify many of the procedures of statistical inference. (More detail later.)*

The Good News:

- *If the population is large enough, the preliminary definition is close enough for all practical purposes.*
- *In many cases where the population is not “large enough,” there are alternate theorems giving rise to alternate procedures using a “finite population correction factor” that will work.*
 - Unfortunately, the question, "How large is large enough?" does not have a simple answer.
- *In many cases, even if the population is not large enough, there are alternate procedures (known as permutation or randomization or resampling tests) that are applicable.*

Consequent Problems with Small Populations:

- 1) Using a “large population” procedure with a “small population” is a **common mistake**.
- 2) One more difficulty in selecting an appropriate sample, which leads to one more source of uncertainty.

II. WHY RANDOM SAMPLING IS IMPORTANT

Recall the Myth:

"A random sample will be representative of the population".

A slightly better explanation (partly true but partly Urban Legend):

"Random sampling prevents bias by giving all individuals an equal chance to be chosen."

- The element of truth: Random sampling will eliminate *systematic* bias.
- A practical rationale: This explanation is often the best plausible explanation that is acceptable to someone with little mathematical background.
- However, this statement could easily be misinterpreted as the myth above.

An additional, very important, reason why random sampling is important, at least in frequentist statistical procedures, which are those most often taught (especially in introductory classes) and used:

The Real Reason: *The mathematical theorems that justify most parametric frequentist statistical procedures apply only to truly random samples.*

The next section elaborates.

III. OVERVIEW OF FREQUENTIST HYPOTHESIS TESTING

Basic Elements of Most Frequentist Hypothesis Tests:

Most commonly-used (“parametric”), frequentist hypothesis tests involve the following four elements:

- i. *Model assumptions*
- ii. *Null and alternative hypotheses*
- iii. *A test statistic*
 - A *test statistic* is something calculated by a rule from a sample.
 - It needs to have the property that *extreme values of the test statistic are rare, and hence cast doubt on the null hypothesis.*
- iv. *A mathematical theorem* saying,

"If the model assumptions and the null hypothesis are both true, then the *sampling distribution* of the test statistic has a certain particular form."

Note:

- The *sampling distribution* is the probability distribution of the test statistic, when considering *all* possible suitably random samples of the same size. (More later.)
- The exact details of these four elements will depend on the particular hypothesis test.
- In particular, the form of the sampling distribution will depend on the hypothesis test.

Illustration: Large Sample z-Test for the mean, with two-sided alternative

The above elements for this test are:

1. *Model assumptions*: We are working with simple random samples of a random variable Y that has a normal distribution.

2. *Null hypothesis*: "The mean of the random variable Y is a certain value μ_0 ."

Alternative hypothesis: "The mean of the random variable Y is not μ_0 ." (This is called the *two-sided alternative*.)

3. *Test statistic*: \bar{y} (the *sample mean* of a simple random sample of size n from the random variable Y).

Before discussing item 4 (the mathematical theorem), we first need to to:

1. Clarify terminology
2. Discuss sampling distributions

1. Terminology and common confusions:

- The **mean of the random variable** Y is also called the *expected value* or the *expectation* of Y .
 - It's denoted $E(Y)$.
 - It's also called the *population mean*, often denoted as μ .
 - It's what we do *not* know in this example.
- A **sample mean** is typically denoted \bar{y} (read "y-bar").
 - It's calculated from a sample y_1, y_2, \dots, y_n of values of Y by the familiar formula $\bar{y} = (y_1 + y_2 + \dots + y_n)/n$.
- The sample mean \bar{y} is an *estimate* of the population mean μ , but they are usually *not* the same.
 - Confusing them is a **common mistake**.
- Note the articles: "*the* population mean" but "*a* sample mean".
 - There is only one *population* mean associated with the random variable Y .
 - However, a *sample* mean depends on the sample chosen.
 - Since there are many possible samples, there are many possible sample means.

Illustration:

http://wise.cgu.edu/sdmmod/sdm_applet.asp

Putting this in a more general framework:

- A *parameter* is a constant associated with a population.
 - In our example: The population mean μ is a parameter.
 - When applying statistics, parameters are usually unknown.
 - However, the goal may be to gain some information about parameters.
- To help gain information about (unknown) parameters, we use *estimates* that are calculated from a sample.
 - In our example: We calculate the sample mean \bar{y} as an estimate of the population mean (parameter) μ .

“Variance” is another common example where parameter and estimate might be confused:

- The *population variance* (or “the variance of Y”) is a parameter, usually called σ^2 (or $\text{Var}(Y)$).
- If we have a sample from Y, we can calculate the *sample variance*, usually called s^2 .
- A sample variance is an estimate of the population variance.
- Different samples may give different estimates.

2. Sampling Distribution:

Although we apply a hypothesis test using a single sample, we need to step back and consider *all* possible simple random samples of Y of size n, in order to understand the test.

- For each simple random sample of Y of size n, we get a value of \bar{y} .
- We thus have a *new* random variable \bar{Y}_n :
 - The associated random process is “pick a simple random sample of size n”
 - The value of \bar{Y}_n is the sample mean \bar{y} for this sample
- Note that
 - \bar{Y}_n stands for the new random variable
 - \bar{y} stands for the value of \bar{Y}_n , for a particular sample of size n.
 - \bar{y} (the value of \bar{Y}_n) depends on the sample, and typically varies from sample to sample.
- The distribution of the new random variable \bar{Y}_n is called the *sampling distribution of \bar{Y}_n* (or the *sampling distribution of the mean*).
- Note: \bar{Y}_n is an example of an *estimator*: a random variable whose values are estimates.

Rice Simulation of sampling distribution:

http://onlinestatbook.com/stat_sim/sampling_dist/index.html

Now we can state the theorem that the large sample z-test for the mean relies on:

4. The *theorem* states: *If* the model assumptions are all true (i.e., *if* Y is normal and all samples considered are simple random samples), and *if in addition* the mean of Y is indeed μ_0 (i.e., *if* the null hypothesis is true), then

- i. The sampling distribution of \bar{Y}_n is normal
- ii. The sampling distribution of \bar{Y}_n has mean μ_0
- iii. The sampling distribution of \bar{Y}_n has standard deviation $\frac{\sigma}{\sqrt{n}}$, where σ is the standard deviation of the original random variable Y.

Check that this is consistent with what the simulation shows.

Also note:


- $\frac{\sigma}{\sqrt{n}}$ is smaller than σ (if n is larger than 1)
- The larger n is, the smaller $\frac{\sigma}{\sqrt{n}}$ is.
- Why is this nice?

More Terminology: σ is called the population standard deviation of Y; it is *not* the same as the sample standard deviation s, although s is an estimate of σ .

The following chart and picture summarize the conclusion of the theorem and related information:

	Considering the population ↓	Considering <i>one</i> sample ↓	Considering <i>all</i> suitable samples ↓
	Random variable Y (population distribution)	Related quantity calculated from a sample y_1, y_2, \dots, y_n	Random variable \bar{Y}_n (sampling distribution)
Type of Distribution	Y has a normal distribution	The sample is <i>from</i> the (normal) distribution of Y	\bar{Y}_n has a normal distribution
Mean	<u>Population mean</u> μ ($\mu = \mu_0$ <u>if</u> null hypothesis true)	<u>Sample mean</u> $\bar{y} = (y_1 + y_2 + \dots + y_n)/n$ \bar{y} is an <u>estimate</u> of the population mean μ	<u>Mean of the sampling distribution of \bar{y}</u> -- it's also μ . ($\mu = \mu_0$ <u>if</u> null hypothesis true)
Standard deviation	<u>Population standard deviation</u> σ	<u>Sample standard deviation</u> $s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (\bar{y} - y_i)^2}$ s is an <u>estimate</u> of the population standard deviation σ	<u>Sampling distribution standard deviation</u> $\frac{\sigma}{\sqrt{n}}$

↑
From the Theorem



- Which column of the chart corresponds to the blue distribution?
- Which column of the chart corresponds to the red distribution?
- How could you tell without looking at the legend?

The roles of the model assumptions for this hypothesis test
(large sample z-test for the mean):

Recall:

The theorem has *three assumptions*:

Assumption 1: Y has a normal distribution (*a model assumption*).

Assumption 2: All samples considered are simple random samples (*also a model assumption*).

Assumption 3: The null hypothesis is true (*assumption for the theorem, but not a model assumption*).

The theorem also has *three conclusions*:

Conclusion 1: The sampling distribution of \bar{Y}_n is normal

Conclusion 2: The sampling distribution of \bar{Y}_n has mean μ_0

Conclusion 3: The sampling distribution of \bar{Y}_n has standard deviation $\frac{\sigma}{\sqrt{n}}$, where σ is the standard deviation of the original random variable Y.

The following chart shows *which conclusion depends on which assumption(s)*:

	Assumptions of Theorem ↓		
Conclusions <i>about Sampling Distribution</i> (Distribution of \bar{Y}_n) ↓	1: Y normal (model assumption)	2: Simple random samples (model assumption)	3: Null hypothesis true (not a model assumption)
1: Normal	√	√	
2: Mean μ_0			√
3: Standard deviation $\frac{\sigma}{\sqrt{n}}$		√	

↑
Corresponds to third column in table on p. 18

Note that the model assumption that the sample is a simple random sample (in particular, that the Y_i 's as defined earlier are independent) is used to prove

1. that the sampling distribution is normal *and*
2. (even more importantly) that the standard deviation of the sampling distribution is $\frac{\sigma}{\sqrt{n}}$.

(This illustrates a general phenomenon that *independence conditions are usually very important in statistical inference.*)

Consequences (More detail later):

1. *If* the conclusion of the theorem is true, the sampling distribution of \bar{Y}_n is narrower than the original distribution of Y

- In fact, conclusion 3 of the theorem gives us an idea of just how narrow it is, depending on n .
- *This will allow us to construct a useful hypothesis test.*

2. The only way we know the conclusion is true is if we know the hypotheses of the theorem (the model assumptions and the null hypothesis) are true.

3. Thus: *If the model assumptions are not true, then we do not know that the theorem is true, so we do not know that the hypothesis test is valid.*

In the example (large sample z-test for a mean), this translates to:

If the sample is not a simple random sample, or if the random variable is not normal, then the reasoning establishing the validity of the test breaks down.

Importance of Model Assumptions in General:

Different hypothesis tests have different model assumptions.

- Some tests apply to random samples that are not simple.
- For many tests, the model assumptions consist of *several* assumptions.
- If *any one* of these model assumptions is not true, we do not know that the test is valid. (*The bad news*)

Robustness:

Many techniques are **robust** to some departures from at least some model assumptions. (*The good news*)

- “Robustness” means that if the particular assumption is *not too far* from true, then the technique is still approximately valid.
- For example, the large sample z-test for the mean is *somewhat* robust to departures from normality. In particular, for *large enough* sample sizes, the test is very close to accurate.
 - Unfortunately, how large is large enough depends on the distribution of the random variable Y. (*More bad news*)
- See the sampling distribution simulation for an illustration.
 - Try a distribution with a full height spike at the left, a half height spike at the right, and nothing in between, with $n=25$; or three equally spaced spikes, descending in height, with nothing in between.

A very common mistake in using statistics: *Using a hypothesis test without paying attention to whether or not the model assumptions are true and whether or not the technique is robust to possible departures from model assumptions is a.* (More later.)

Caution re terminology “Robust” is used in other ways – for example, “a finding is robust” means that the finding appears to be true in a wide variety of situations.

IV: FREQUENTIST CONFIDENCE INTERVALS

Before continuing the discussion of hypothesis tests, it will be helpful first to discuss the related concept of confidence intervals.

The General Situation:

- We're considering a *random variable* Y .
- We are interested in a certain *parameter* (e.g., a proportion, or mean, or regression coefficient, or variance) associated with the random variable Y (i.e., associated with the population)
- We *don't* know the value of the parameter.
- *Goal 1*: We'd like to estimate the unknown parameter, using data from a sample. (*A point estimate*)
- *Goal 2*: We'd also like to get some sense of how good our estimate is. (Typically achieved by an *interval estimate*)

The first goal is usually easier than the second.

Example: If the parameter we are interested in estimating is the *mean of the random variable* (i.e., the population mean, which we call ____), we can estimate it using a *sample mean* (which we call ____).

The idea for achieving the second goal (getting some sense of how good our estimate is) has a lot in common with hypothesis testing:

- It involves the sampling distribution
- It depends on model assumptions.

A little more specifically:

- Although we typically have just one sample at hand when we do statistics, *the reasoning used in classical frequentist inference depends on thinking about all possible suitable samples of the same size n .*
- Which samples are considered "suitable" will depend on the particular statistical procedure to be used.
- Each statistical procedure has *model assumptions* that are needed to ensure that the reasoning behind the procedure is sound.
- The model assumptions determine which samples are "suitable."

Illustration: Large Sample z-Procedure for a Confidence Interval for a Mean

- The parameter we want to estimate is the population mean $\mu = E(Y)$ of the random variable Y .
- The model assumptions for this procedure are: The random variable is normal, and samples are simple random samples.
 - So in this case, "suitable sample" means "simple random sample".
 - We'll also assume Y is normal, so both model assumptions are satisfied.

Notation and terminology:

- We'll use σ to denote the (population) standard deviation of Y .
- We have a simple random sample, say of size n , consisting of observations y_1, y_2, \dots, y_n .
 - For example, if Y is "height of an adult American male", we take a simple random sample of n adult American males; y_1, y_2, \dots, y_n are their heights.
- We use the sample mean $\bar{y} = (y_1 + y_2 + \dots + y_n)/n$ as our estimate of the population mean μ .
 - This is an example of a **point estimate** -- a numerical estimate with no indication of how good the estimate is.

More details:

- To get an idea of how good our estimate is, we use the concept of *confidence interval*.
 - This is an example of an **interval estimate**.
- To understand the concept of "confidence interval", we need to consider all possible simple random samples of size n from Y .
 - In the specific example, we consider all possible simple random samples of n adult American males.
 - For each such sample, the heights of the men in the sample of people constitute our simple random sample of size n from Y .
- We consider the *sample means \bar{y} for all possible simple random samples of size n from Y* .
 - This amounts to defining a new random variable, which we will call \bar{Y}_n (read "Y-bar sub n ", or "Y-sub- n -bar").
 - We can describe the random variable \bar{Y}_n briefly as "sample mean of a simple random sample of size n from Y ", or more explicitly as: "pick a simple random sample of size n from Y and calculate its sample mean".
 - Note *that each value of \bar{Y}_n is an estimate of the population mean μ* .
 - e.g. each simple random sample of n adult American males gives us an estimate of the population mean μ .

- This new random variable \bar{Y}_n has a distribution, called a *sampling distribution* (since it arises from considering varying samples).
 - The values of \bar{Y}_n are all the possible values of sample means \bar{y} of simple random samples of size n of Y – i.e., the values of our *estimates* of μ .
 - *The sampling distribution (distribution of \bar{Y}_n) gives us information about the variability (as samples vary) of our estimates of the population mean μ .*
 - A mathematical theorem tells us that *if* the model assumptions are true, then:
 1. The sampling distribution is normal
 2. The mean of the sampling distribution is also μ .
 3. The sampling distribution has standard deviation $\frac{\sigma}{\sqrt{n}}$
 - Use these conclusions to compare and contrast the shapes of the distribution of Y and the distribution of \bar{Y}_n
 - What is the same?
 - What is different?
 - How do the widths compare?
 - The chart (best read one column at a time) and picture below summarize some of this information.