

M346 (92153), Homework #13

Due: 10:00am, Thursday, Jul. 05

**Infinite-dimensional inner product spaces (6.8)**

A)

- i. Let  $\mathbf{v} = (a_1, a_2, a_3, \dots)$  with  $a_n = (-1)^n / \sqrt{n}$ . Is  $\mathbf{v}$  in  $l_2(\mathbb{R})$ ?
- ii. Consider the function  $f(x) = 1/x^p$ . For what  $p \geq 0$  is  $f$  in  $L_2([1, \infty))$ ? For what  $p \geq 0$  is  $f$  in  $L_2((0, 1])$ ?

**Fourier series (6.9, 8.5, 8.7)**

A) Use integration by parts to evaluate the following integrals with constant  $k \neq 0$ :

- i.  $\int_0^1 x \sin(kx) dx$
- ii.  $\int_0^1 x \cos(kx) dx$
- iii.  $\int_0^1 x \exp(ikx) dx$

[Hint: Use Euler's formula and your answers to (i) and (ii).]

B) Let  $f(x) = x$  for  $x \in [0, 1]$ . Use your solutions from problem (A) for the following parts:

- i. Write the Fourier sine series for  $f$ —that is, write

$$f(x) = \sum_{n=1}^{\infty} c_n \sin(n\pi x)$$

by finding the coefficients

$$c_n = 2 \int_0^1 f(x) \sin(n\pi x) dx, \quad n \in \{1, 2, 3, \dots\}.$$

How fast do the coefficients  $c_n$  decay?

- ii. Plot  $f(x)$  and the approximations  $f_N(x) = \sum_{n=1}^N c_n \sin(n\pi x)$  for  $N = 1, 2, 3, 4$ .

C)

- i. Derive a solution  $u(x, t)$  to the partial differential equation (PDE)

$$\begin{aligned} \partial_t u &= -\partial_{xx} u \\ u(0, t) &= 0, \quad u(a, t) = 0 & x \in [0, a], \quad t \geq 0 \\ u(x, 0) &= \begin{cases} x & \text{if } x < a/2 \\ a - x & \text{if } x \geq a/2 \end{cases} \end{aligned}$$

using Fourier sine series. How does the  $n^{\text{th}}$  Fourier coefficient evolve, and what does this imply about the solution  $u(x, t)$  for arbitrarily small times  $t > 0$ ? Contrast this behavior to that of the solution to the ordinary heat equation  $\partial_t u = \partial_{xx} u$  discussed in class.

[Note: The equation above is called the *backward* heat equation because it arises from the ordinary heat equation under the time change  $t \rightarrow -t$ . It is *ill-posed* in that it behaves extremely badly for almost all initial conditions  $u(x, 0)$ .]

ii. Instead, solve the PDE

$$\begin{aligned} \partial_t u &= -\partial_{xx} u - \partial_{xxxx} u \\ u(0, t) &= 0, \quad u(a, t) = 0 & x \in [0, a], \quad t \geq 0 \\ u(x, 0) &= \begin{cases} x & \text{if } x < a/2 \\ a - x & \text{if } x \geq a/2 \end{cases} \end{aligned}$$

using Fourier sine series. Now how does the  $n^{\text{th}}$  Fourier coefficient evolve and what does this imply about the solution  $u(x, t)$  for small times  $t > 0$ ? What happens as  $t \rightarrow \infty$ ? Again, compare this to the ordinary heat equation.

[Note: By adding the term  $-\partial_{xxxx} u$  to the equation, we have dramatically changed its behavior. This term, called a fourth-order regularization, overcomes the ill-posed nature of the term  $-\partial_{xx} u$ .]