



Nonarchimedean geometry, tropicalization, and metrics on curves

Matthew Baker, Sam Payne and Joseph Rabinoff

ABSTRACT

We develop a number of general techniques for comparing analytifications and tropicalizations of algebraic varieties. Our basic results include a projection formula for tropical multiplicities and a generalization of the Sturmfels–Tevelev multiplicity formula in tropical elimination theory to the case of a nontrivial valuation. For curves, we explore in detail the relationship between skeletal metrics and lattice lengths on tropicalizations and show that the maps from the analytification of a curve to the tropicalizations of its toric embeddings stabilize to isometries on finite subgraphs. Other applications include generalizations of Speyer’s well-spacedness condition and the Katz–Markwig–Markwig results on tropical j -invariants.

1. Introduction

The recent work of Gubler [Gub07a, Gub07b], in addition to earlier work of Bieri–Groves [BG84], Berkovich [Ber90, Ber99, Ber04], and others, has revealed close connections between nonarchimedean analytic spaces (in Berkovich’s sense) and tropical geometry. One such connection is given by the second author’s theorem that ‘analytification is the inverse limit of all tropicalizations’ (see Theorem 1.2). This result is purely topological, providing a natural homeomorphism between the nonarchimedean analytification X^{an} of a quasiprojective variety X and the inverse limit of all ‘extended tropicalizations’ of X coming from closed immersions of X into quasiprojective toric varieties that meet the dense torus. In this paper, we develop a number of general techniques for comparing finer properties of analytifications and tropicalizations of algebraic varieties and apply these techniques to explore in detail the relationship between the natural metrics on analytifications and tropicalizations of curves. The proofs of our main results rely on the geometry of formal models and initial degenerations as well as Berkovich’s theory of nonarchimedean analytic spaces.

Let K be an algebraically closed field that is complete with respect to a nontrivial nonarchimedean valuation $\text{val}: K \rightarrow \mathbb{R} \cup \{\infty\}$. Let X be a nonsingular curve defined over K . The underlying topological space of X^{an} can be endowed with a ‘polyhedral’ structure locally modeled on an

Received 24 October 2014, accepted in final form 25 May 2015.

2010 Mathematics Subject Classification 14T05 (primary) 14G22, 14M25 (secondary).

Keywords: algebraic curves, tropical curves, tropical geometry, faithful tropicalization, extended tropicalization, initial degeneration, tropical multiplicities, nonarchimedean geometry, analytification, Berkovich spaces, skeletons, well-spacedness.

This journal is © Foundation Compositio Mathematica 2016. This article is distributed with Open Access under the terms of the Creative Commons Attribution Non-Commercial License, which permits non-commercial reuse, distribution, and reproduction in any medium, provided that the original work is properly cited. For commercial re-use, please contact the Foundation Compositio Mathematica.

The first and second authors were supported in part by NSF Research Grants DMS-0901487 and DMS-1068689, respectively.

\mathbb{R} -tree. The leaves of X^{an} are the K -points, together with the ‘type 4 points’ in Berkovich’s classification (see Section 3.5). The nonleaves are exactly those points that are contained in an embedded open segment, and the space $\mathbf{H}_o(X^{\text{an}})$ of nonleaves carries a canonical metric which, like the polyhedral structure, is defined using semistable models for X . Our primary reference for these results is [BPR13]; see also [Ber90, § 4], [Thu05], and [Bak08, § 5].

Suppose that X is embedded in a toric variety Y_Δ and meets the dense torus \mathbf{T} . The tropicalization $\text{Trop}(X \cap \mathbf{T})$ is a 1-dimensional polyhedral complex with no leaves in the real vector space spanned by the lattice of 1-parameter subgroups of \mathbf{T} . All edges of $\text{Trop}(X \cap \mathbf{T})$ have slopes that are rational with respect to the lattice of 1-parameter subgroups, so there is a natural metric on $\text{Trop}(X \cap \mathbf{T})$ given locally by lattice length on each edge, and globally by shortest paths. The metric space $\mathbf{H}_o(X^{\text{an}})$ naturally surjects onto $\text{Trop}(X \cap \mathbf{T})$, but this map is far from being an isometry since infinitely many embedded segments in $\mathbf{H}_o(X^{\text{an}})$ are contracted. Furthermore, even when an edge of $\mathbf{H}_o(X^{\text{an}})$ maps homeomorphically onto an edge of $\text{Trop}(X \cap \mathbf{T})$, this homeomorphism need not be an isometry; see Section 2.5. Nevertheless, each embedded subgraph in $\mathbf{H}_o(X^{\text{an}})$ maps isometrically onto its image in all ‘sufficiently large’ tropicalizations.

THEOREM 1.1. *Let Γ be a finite embedded subgraph in $\mathbf{H}_o(X^{\text{an}})$. Then there is a closed embedding of X into a quasiprojective toric variety such that X meets the dense torus and Γ maps isometrically onto its image in $\text{Trop}(X \cap \mathbf{T})$. Furthermore, the set of all such embeddings is stable and hence cofinal in the system of all embeddings of X into quasiprojective toric varieties whose images meet \mathbf{T} .*

Here if $\iota: X \hookrightarrow Y_\Delta$ and $\iota': X \hookrightarrow Y_{\Delta'}$ are closed embeddings into quasiprojective toric varieties such that X meets the dense tori \mathbf{T} and \mathbf{T}' , then we say that ι' *dominates* ι and we write $\iota' \geq \iota$ if there exists an equivariant morphism of toric varieties $\psi: Y_{\Delta'} \rightarrow Y_\Delta$ such that $\psi \circ \iota' = \iota$ (see Section 5.15). In this case we have an induced map $\text{Trop}(X \cap \mathbf{T}') \rightarrow \text{Trop}(X \cap \mathbf{T})$; the above theorem says in particular that if Γ maps isometrically onto its image in $\text{Trop}(X \cap \mathbf{T})$, then the same is true for $\Gamma \rightarrow \text{Trop}(X \cap \mathbf{T}')$. In other words, the maps from $\mathbf{H}_o(X^{\text{an}})$ to the tropicalizations of toric embeddings of X stabilize to isometries on every finite subgraph.

Both the analytification and the tropicalization constructions described above for subvarieties of tori globalize in natural ways. The analytification functor extends to arbitrary finite-type K -schemes (see [Ber90, Chapters 2 and 3] or [Ber93]), and tropicalization extends to closed subvarieties of toric varieties as follows. If Δ is a fan in $N_{\mathbb{R}}$ and Y_Δ is the associated toric variety, then there is a natural ‘partial compactification’ $N_{\mathbb{R}}(\Delta)$ of $N_{\mathbb{R}}$ which is, set theoretically, the disjoint union of the tropicalizations of all torus orbits in Y_Δ . The topology on $N_{\mathbb{R}}(\Delta)$ is such that the natural map from $Y_\Delta(K)$ extends to a continuous, proper, and surjective map $\text{trop}: Y_\Delta^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta)$. As in the case where Y_Δ is the torus \mathbf{T} , the tropicalization $\text{Trop}(X)$ of a closed subvariety X in Y_Δ is the closure of $\text{trop}(X(K))$ in $N_{\mathbb{R}}(\Delta)$, and the extended tropicalization map extends to a continuous, proper, surjective map from X^{an} onto $\text{Trop}(X)$. See [Pay09a, Rab12] and Section 4.2 for further details.

THEOREM 1.2 (Payne). *Let X be an irreducible quasiprojective variety over K . Then the inverse limit of the extended tropicalizations $\text{Trop}(\iota(X))$ over all closed immersions $\iota: X \hookrightarrow Y_\Delta$ into quasiprojective toric varieties is canonically homeomorphic to the analytification X^{an} .*

The inverse limit in Theorem 1.2 can be restricted to those closed immersions ι whose images meet the dense torus T_ι , and then the homeomorphism maps $X^{\text{an}} \setminus X(K)$ homeomorphically onto the inverse limit of the ordinary tropicalizations $\text{Trop}(\iota(X) \cap T_\iota)$.

When X is a curve, our Theorem 1.1 says that the metric structures on $\text{trop}(\iota(X) \cap T_\ell)$ stabilize to a metric on the subset $\mathbf{H}_\circ(X^{\text{an}})$ of the inverse limit, and the restriction of this homeomorphism is an *isometry*. In general, each sufficiently small segment e in $\mathbf{H}_\circ(X^{\text{an}})$ is mapped via an affine linear transformation with integer slope onto a (possibly degenerate) segment e' in $\text{Trop}(X)$. We write $m_{\text{rel}}(e)$ for the absolute value of the slope of this map, so if e has length ℓ , then its image e' has lattice length $m_{\text{rel}}(e) \cdot \ell$. In Corollary 5.9, we relate these ‘expansion factors’ to *tropical multiplicities* of edges in $\text{Trop}(X)$. The notation is meant to suggest that $m_{\text{rel}}(e)$ may be thought of in this context as the relative multiplicity of e over e' . By definition, the tropical multiplicity $m_{\text{Trop}}(e')$ of an edge e' in a suitable polyhedral structure on $\text{Trop}(X)$ is the number of irreducible components, counted with multiplicities, in the *initial degeneration* $\text{in}_w(X \cap \mathbf{T})$ for any w in the relative interior of e' . These tropical multiplicities are fundamental invariants in tropical geometry and play a key role in the *balancing formula*. See Section 2.1 for a definition of the initial degeneration $\text{in}_w(X \cap \mathbf{T})$ and further discussion of tropical multiplicities.

THEOREM 1.3. *There is a polyhedral structure on $\text{Trop}(X \cap \mathbf{T})$ with the following properties:*

- (i) *For each edge e' in $\text{Trop}(X \cap \mathbf{T})$, there are finitely many embedded segments e_1, \dots, e_r in $\mathbf{H}_\circ(X^{\text{an}})$ mapping homeomorphically onto e' .*
- (ii) *Any embedded segment in the preimage of e' that is disjoint from $e_1 \cup \dots \cup e_r$ is contracted to a point.*
- (iii) *The tropical multiplicity of e' is the sum of the corresponding expansion factors:*

$$m_{\text{Trop}}(e') = m_{\text{rel}}(e_1) + \dots + m_{\text{rel}}(e_r).$$

The properties above are preserved by subdivision, so they hold for any sufficiently fine polyhedral structure on $\text{Trop}(X \cap \mathbf{T})$. See Proposition 5.4 and Corollary 5.9.

The tropical multiplicity formula in the above theorem gives an important connection to nonarchimedean analytic spaces that is not visible from the definitions. The formula shows, for example, that if e' is a small segment in $\text{Trop}(X \cap \mathbf{T})$ whose tropical multiplicity is equal to 1, then there is a unique segment e in $\mathbf{H}_\circ(X^{\text{an}})$ mapping homeomorphically onto e' , and the length of e is equal to the tropical length of e' . It is well known that the skeleton of the analytification of an elliptic curve with bad reduction is a loop of length equal to minus the valuation of the j -invariant (see for example [BPR13, Remark 4.24]), so these formulas explain earlier results of Katz, Markwig, and Markwig on tropical j -invariants of genus 1 curves in toric surfaces [KMM08, KMM09]. See, for instance, Example 2.8. The following theorem also provides natural generalizations for genus 1 curves in higher-dimensional toric varieties, as well as curves of arbitrary genus.

THEOREM 1.4. *Let Γ' be a finite embedded subgraph of $\text{Trop}(X \cap \mathbf{T})$, and suppose that $\text{in}_w(X \cap \mathbf{T})$ is irreducible and generically reduced for every w in Γ' . Then there is a unique embedded subgraph Γ in $\mathbf{H}_\circ(X^{\text{an}})$ mapping homeomorphically onto Γ' , and this homeomorphism is an isometry.*

See Section 6 for details on deducing the tropical j -invariant results of Katz, Markwig, and Markwig from this theorem.

The expansion factors $m_{\text{rel}}(e)$ in our tropical multiplicity formula are often computable in practice. If X is an affine curve embedded in the torus \mathbf{G}_m^n via an n -tuple of invertible regular functions f_1, \dots, f_n , then

$$m_{\text{rel}}(e) = \gcd(s_1(e), \dots, s_n(e)),$$

where $s_i(e)$ is the absolute value of the slope of the integer-affine function $\log |f_i|$ along the edge e . See Remark 5.6. The quantities $s_i(e)$ are easily calculated from the divisors of f_1, \dots, f_n using the ‘slope formula’ of [BPR13, Theorem 5.15].

In concrete situations, it is useful to be able to certify that a given tropicalization map faithfully represents a large piece of the nonarchimedean analytification X^{an} (for example, the ‘minimal skeleton’ Σ of X^{an} in the sense of Berkovich [Ber90] or [BPR13, Corollary 4.23]) using only ‘tropical’ computations (for example, Gröbner complex computations which have been implemented in computer algebra packages such as SINGULAR or Macaulay2), as opposed to calculations with formal models that have not been implemented in a systematic way in any existing software package. We prove that a tropicalization map represents Σ faithfully, meaning that the map is an isometry on Σ , provided that certain combinatorial and topological conditions are satisfied. Our results on faithful representations are presented in conjunction with some observations about initial degenerations which help explain the special role played by trivalent graphs in the literature on tropical curves (cf. Theorem 5.25 and Remark 5.27).

We explore tropicalizations of elliptic curves in detail as a concrete illustration of our methods and results. We are able to say some rather precise things in this case; for example, we show that every elliptic curve E/K with multiplicative reduction admits a closed embedding in \mathbb{P}^2 whose tropicalization faithfully (and certifiably) represents the minimal skeleton of E^{an} . Furthermore, we interpret Speyer’s ‘well-spacedness condition’ for trivalent tropicalizations of totally degenerate genus 1 curves [Spe14] as a statement about rational functions on the analytification of the curve, and prove generalizations of this condition for nontrivalent tropicalizations, and for genus 1 curves with good reduction.

The paper concludes with a generalization of the important Sturmfels–Tevelev multiplicity formula, which calculates $\text{Trop}(\alpha(X))$ (as a weighted polyhedral complex) in terms of $\text{Trop}(X)$ when $\alpha: \mathbf{T} \rightarrow \mathbf{T}'$ is a homomorphism of tori which induces a generically finite map from a subvariety X in \mathbf{T} onto its image. The multiplicity formula in [ST08] is formulated and proved in the ‘constant coefficient’ setting, where K is the field of Puiseux series over an algebraically closed coefficient field k of characteristic 0 and X is defined over k . We use the methods of this paper to generalize the Sturmfels–Tevelev formula to the case where X is any closed subvariety of a torus \mathbf{T} defined over a complete and algebraically closed nonarchimedean field K .

Philosophically speaking, there are at least two long-term goals to this paper. On the one hand, we believe that the systematic use of modern tools from nonarchimedean geometry is extremely useful for understanding and proving theorems in tropical geometry. This paper takes several steps in that direction, establishing some new results in tropical geometry via Berkovich’s theory and the Bosch–Lütkebohmert–Raynaud theory of admissible formal schemes. On the other hand, much of this paper can be viewed as a comparison between two different ways of approximating nonarchimedean analytic spaces. Nonarchimedean analytic spaces have proved to be useful in many different contexts, but the topological spaces underlying them are wildly branching infinite complexes which are difficult to study directly, so one usually approximates them with finite polyhedral complexes. One such approximation goes through skeleta of nice (for example, semistable) formal models (cf. [BPR13, Theorem 5.2]), another through (extended) tropicalizations (cf. Theorem 1.2). Our Theorem 5.21 shows that, in the case of curves, these two approximations have the same metric structure in the limit (though the metrics may be different at any given finite level).

For further details and examples, we refer the reader to the expanded, earlier preprint version of this paper, [arxiv:1104.0320v2](https://arxiv.org/abs/1104.0320v2). This final version differs from that one in several respects. The

numbering (of equations, paragraphs, sections, theorems, etc.) has changed. The former Section 5, *The structure theory of analytic curves*, was extracted and published separately [BPR13]. Furthermore, much expository material and many examples in the remaining sections have been omitted. The earlier preprint version remains available on the arXiv.

Since this paper was written, there have been a large number of follow-up articles: the reader may also want to read [GRW14], in which many of the results in this article are extended to higher dimensions; [CHW14, DP14, CS13], in which several interesting examples of faithful tropicalizations are given; and [ABBR15a, ABBR15b], in which ‘relative’ versions of some of the results in this paper are used to prove tropical lifting theorems.

2. Basic notions and examples

Here we give a brief overview of the basic notions necessary to understand the theorems stated in the introduction, followed by a few key examples illustrating these results. Throughout this paper, K is an algebraically closed field that is complete with respect to a nontrivial nonarchimedean valuation

$$\text{val}: K \rightarrow \mathbb{R} \cup \{\infty\}.$$

We let $G = \text{val}(K^\times)$ be its value group, $R = \text{val}^{-1}([0, \infty])$ its valuation ring, $\mathfrak{m} \subset R$ the maximal ideal, and $k = R/\mathfrak{m}$ its residue field (which is algebraically closed by [Rob00, § 2.1, Proposition 3]). Let $|\cdot| = \exp(-\text{val}(\cdot))$ be the absolute value on K associated with the valuation.

2.1 Tropicalization

Let M be a free abelian group of rank n , let $\mathbf{T} = \text{Spec}(K[M])$ be the K -torus with character group M , and let $N = \text{Hom}(M, \mathbb{Z})$ be the dual lattice. If X is a closed subscheme of \mathbf{T} , there is a natural *tropicalization map*

$$\text{trop}: X(K) \rightarrow N_{\mathbb{R}},$$

where $N_{\mathbb{R}} = \text{Hom}(M, \mathbb{R})$. The image of a point x in $X(K)$ is the linear function taking $u \in M$ to the valuation of the corresponding character evaluated at x . Then $\text{Trop}(X)$ is the closure of $\text{trop}(X(K))$ in the Euclidean topology on $N_{\mathbb{R}}$. Note that the choice of an isomorphism $M \cong \mathbb{Z}^n$ induces an identification of \mathbf{T} with \mathbf{G}_m^n . In such coordinates, the tropicalization map sends a point (x_1, \dots, x_n) in $X(K)$ to $(\text{val}(x_1), \dots, \text{val}(x_n))$ in \mathbb{R}^n .

One of the basic results in tropical geometry says that if X is an integral subscheme of \mathbf{T} of dimension d , then $\text{Trop}(X)$ is the underlying set of a connected ‘balanced weighted integral G -affine polyhedral complex’ of pure dimension d . We do not define all of these terms here, but briefly recall how one gets a polyhedral complex and defines weights on the maximal faces of this complex. Let w be a point in $N_G = \text{Hom}(M, G)$. The ‘tilted group ring’ $R[M]^w$ is the subring of $K[M]$ consisting of Laurent polynomials $a_1 x^{u_1} + \dots + a_r x^{u_r}$ such that

$$\text{val}(a_i) + \langle u_i, w \rangle \geq 0$$

for all i . The R -scheme $\mathbf{T}^w = \text{Spec } R[M]^w$ is a torsor for the torus $\text{Spec } R[M]$, and its generic fiber is canonically isomorphic to \mathbf{T} . If X is a closed subscheme of \mathbf{T} defined by an ideal $\mathfrak{a} \subset K[M]$, then

$$X^w = \text{Spec} (R[M]^w / (\mathfrak{a} \cap R[M]^w))$$

is a flat R -scheme with generic fiber X , which we call the *tropical integral model* associated with w . It is exactly the closure of X in \mathbf{T}^w . The special fiber $\text{in}_w(X)$ of X^w is called the *initial*

degeneration of X with respect to w and is the subscheme of the special fiber of \mathbf{T}^w cut out by the w -initial forms of Laurent polynomials in \mathfrak{a} , in the sense of generalized Gröbner theory.

The scheme \mathbf{T}^w is not proper, so points in $X(K)$ may fail to have limits in the special fiber. Indeed, the special fiber $\text{in}_w(X)$ is often empty. One of the fundamental theorems in tropical geometry says that w is in $\text{Trop}(X)$ if and only if $\text{in}_w(X)$ is not empty.¹ Moreover, $\text{Trop}(X)$ can be given the structure of a finite polyhedral complex in such a way that whenever w and w' belong to the relative interior of the same face, the corresponding initial degenerations $\text{in}_w(X)$ and $\text{in}_{w'}(X)$ are \mathbf{T} -affinely equivalent.

We define the *multiplicity* $m_{\text{Trop}}(w)$ of a point w in $\text{Trop}(X)$ to be the number of irreducible components of $\text{in}_w(X)$, counted with multiplicities. In particular, $m_{\text{Trop}}(w) = 1$ if and only if $\text{in}_w(X)$ is irreducible and generically reduced. These tropical multiplicities are constant on the relative interior of each face F of $\text{Trop}(X)$, and we define the multiplicity $m_{\text{Trop}}(F)$ to be $m_{\text{Trop}}(w)$ for any w in the relative interior of F . The multiplicities for maximal faces are the ‘weights’ mentioned above that appear in the balancing condition. These weights have the following simple interpretation for hypersurfaces.

Remark 2.2. If $X = V(f)$ is a hypersurface, then $\text{Trop}(X)$ is the corner locus of the convex piecewise-linear function associated with a defining equation f [EKL06, §2.1]. In this case, $\text{Trop}(X)$ has a unique minimal polyhedral structure, and the initial degenerations are essentially constant on the relative interior of each face. There is a natural inclusion reversing bijection between the faces of $\text{Trop}(X)$ in this minimal polyhedral structure and the positive-dimensional faces of the *Newton polytopal complex* (or *Newton complex*) of f : a face of $\text{Trop}(X)$ corresponds to the convex hull of the monomials whose associated affine linear function is minimal on that face. In particular, the maximal faces of $\text{Trop}(X)$ correspond to the edges of this Newton complex. In this special case, the multiplicity of a maximal face is the lattice length of the corresponding edge.² The relationship between the tropical hypersurface and the Newton complex is also explained in more detail in [Rab12, §8].

2.3 Analytification

Let A be a finite-type K -algebra. The *Berkovich spectrum* of A , denoted $\mathcal{M}(A)$, is defined to be the set of multiplicative seminorms $\|\cdot\|$ on A extending the absolute value $|\cdot|$ on K . The Berkovich spectrum $\mathcal{M}(A)$ is the underlying set of the nonarchimedean analytification X^{an} of $X = \text{Spec}(A)$. The topology on X^{an} is the coarsest such that the map $\|\cdot\| \mapsto \|f\|$ is continuous for every $f \in A$; this coincides with the subspace topology induced by the inclusion of X^{an} in \mathbb{R}^A .

Remark. We will often write \mathbb{A}_{an}^1 for $\mathbb{A}^{1,\text{an}}$ and \mathbb{P}_{an}^1 for $\mathbb{P}^{1,\text{an}}$, etc.

If X is connected, then X^{an} is a path-connected locally compact Hausdorff space that naturally contains $X(K)$ as a dense subset; a point $x \in X(K)$ corresponds to the seminorm $\|\cdot\|_x$ given by $\|f\|_x = |f(x)|$. The analytification procedure $X \mapsto X^{\text{an}}$ gives a covariant functor from the category of locally finite-type K -schemes to the category of topological spaces.³

¹In the special case where \mathbf{T} has dimension 1 and X is the zero locus of a Laurent polynomial f , this is equivalent to the statement that f has a root with valuation s if and only if $-s$ is a slope of the Newton polygon of f .

²This is explained in [ST08, Example 3.16] in the special case where X is irreducible, K is the field of Puiseux series over k , and X is defined over k . The arguments given there work in full generality.

³The analytification X^{an} has additional structure, including a structure sheaf, and $X \mapsto X^{\text{an}}$ may also be seen as a functor from locally finite-type K -schemes to locally ringed spaces. See [Ber90, §§2.3, 3.1, 3.4] for more details.

If X is a closed subvariety of \mathbf{T} , then the tropicalization map described in the previous section extends from $X(K)$ to a continuous and proper map

$$\text{trop}: X^{\text{an}} \rightarrow N_{\mathbb{R}}$$

taking a seminorm $\|\cdot\|$ to the linear function $u \mapsto -\log \|x^u\|$, and the image of this map is exactly $\text{Trop}(X)$. In other words, $\text{trop}(X^{\text{an}})$ is the closure of the image of $X(K)$ in $N_{\mathbb{R}}$.

2.4 Metric structure of analytic curves

There is a natural metric on $\mathbb{P}_{\text{an}}^1 \setminus \mathbb{P}^1(K)$; see [BPR13] for details. We write $\mathbf{H}(\mathbb{P}_{\text{an}}^1)$ to denote $\mathbb{P}_{\text{an}}^1 \setminus \mathbb{P}^1(K)$ with this metric structure. It is important to note that the metric topology on $\mathbf{H}(\mathbb{P}_{\text{an}}^1)$ is much finer than the subspace topology on $\mathbb{P}_{\text{an}}^1 \setminus \mathbb{P}^1(K)$. Our notation follows [BR10] and reflects the fact that the metric on $\mathbf{H}(\mathbb{P}_{\text{an}}^1)$ is 0-hyperbolic in the sense of Gromov.

The metric on $\mathbf{H}(\mathbb{P}_{\text{an}}^1)$ has the important property that, roughly speaking, $\log |f|$ is piecewise affine with integer slopes for any nonzero rational function $f \in K(T)$. More precisely, suppose that f is nonconstant and that $\widehat{\Sigma}$ is the minimal closed connected subset of \mathbb{P}_{an}^1 containing the set S of zeros and poles of f . Let $\Sigma = \widehat{\Sigma} \setminus S$. Then:

- (i) The subspace Σ of $\mathbf{H}(\mathbb{P}_{\text{an}}^1)$ is a metric graph with finitely many edges, in which the edges whose closures meet K have infinite length.
- (ii) The restriction of $\log |f|$ to Σ is piecewise affine with integer slopes.
- (iii) There is a natural retraction map from \mathbb{P}_{an}^1 onto $\widehat{\Sigma}$.
- (iv) The function $\log |f|$ from \mathbb{P}_{an}^1 to $\mathbb{R} \cup \{\pm\infty\}$ factors through the retraction onto $\widehat{\Sigma}$, and hence is determined by its restriction to Σ .

The metric on the complement of the set of K -points in the analytification of an arbitrary algebraic curve is induced by the metric on $\mathbb{P}_{\text{an}}^1 \setminus \mathbb{P}^1(K)$ via semistable decomposition; see [BPR13] for details.

There is also a notion of a skeleton of a smooth and connected but not necessarily complete curve X . Let \widehat{X} be the smooth compactification of X , and let $D = \widehat{X} \setminus X$ be the set of ‘punctures’. Choose a semistable model \mathcal{X} of \widehat{X} such that the punctures reduce to distinct smooth points of the special fiber $\overline{\mathcal{X}}$. Then there is unique minimal closed connected subset Σ of X^{an} which contains the skeleton $\Sigma_{\mathcal{X}}$ of \widehat{X} and whose closure in \widehat{X}^{an} contains D . We call Σ the *skeleton* of X associated with \mathcal{X} . As above, there is a canonical retraction map $\tau_{\Sigma}: X^{\text{an}} \rightarrow \Sigma$. If $X \subset \mathbf{T}$, then the tropicalization map $\text{trop}: X \rightarrow N_{\mathbb{R}}$ factors through τ_{Σ} . There is a skeleton which is minimal over all models \mathcal{X} if $2 - 2g(\widehat{X}) - \#D \leq 0$. See [BPR13] for a complete discussion of the skeleta of a curve.

2.5 Examples

To illustrate our main results concerning the relationship between analytification and tropicalization in the case of curves, taking into account the metric structure on both sides, we present the following examples. In each example, we fix a specific coefficient field for concreteness.

Our first example shows how a loop in the analytification of a genus 1 curve can be collapsed onto a segment of multiplicity greater than 1.

EXAMPLE 2.6. Let K be the completion of the field of Puiseux series $\mathbb{C}\{\{t\}\}$. Consider the genus 1 curve $\widehat{E} \subset \mathbb{P}^2$ over K defined by the Weierstrass equation $y^2 = x^3 + x^2 + t^4$, and let $E = \widehat{E} \cap \mathbf{G}_m^2$. The j -invariant of \widehat{E} has valuation -4 , so \widehat{E} has multiplicative reduction and the minimal skeleton




FIGURE 1. The minimal skeleton $\Gamma \subset E^{\text{an}}$ and the tropicalization $\text{Trop}(E)$ from Example 2.6. The points P_i, Q_j are defined as follows: the rational function x on \widehat{E} has divisor $(Q_1) + (Q_2) - 2(\infty)$, where $Q_1 = (0, t^2)$ and $Q_2 = (0, -t^2)$, and y has divisor $(P_1) + (P_2) + (P_3) - 3(\infty)$, where $\text{val}(x(P_1)) = \text{val}(x(P_2)) = 2$ and $\text{val}(x(P_3)) = 0$.

Σ of \widehat{E} is isometric to a circle of circumference 4. In this example, $\text{Trop}(E)$ does not have a cycle even though \widehat{E}^{an} does; it is interesting to examine exactly what tropicalization is doing to \widehat{E}^{an} . Let Γ be the minimal skeleton of E ; as above, trop factors through the retraction of E^{an} onto Γ . Figure 1 shows the restriction of trop to Γ .

The tropicalization map sends Σ two-to-one onto its image in $\text{Trop}(E)$, which is a segment of tropical length 2 and tropical multiplicity 2. Locally on Σ the tropicalization map is an isometry. Each of the rays of Γ emanating from Σ maps isometrically onto its image. The two rays in $\text{Trop}(E)$ with multiplicity 1 have unique preimages in Γ , while there are two distinct rays in Γ mapping onto each of the two rays in $\text{Trop}(E)$ of multiplicity 2.

The following example illustrates a different kind of collapse, where a segment e in the minimal skeleton of the analytification is collapsed to a point, that is, the relative multiplicity $m_{\text{rel}}(e)$ is 0.

EXAMPLE 2.7. Let $p \geq 5$ be a prime, let $K = \mathbb{C}_p$, and let $k \cong \overline{\mathbb{F}}_p$ be its residue field. Let $X \subset \mathbf{G}_m^2$ be the affine curve over K defined by the equation $f(x, y) = x^3y - x^2y^2 - 2xy^3 - 3x^2y + 2xy - p = 0$. The curve $\widehat{X} \subset \mathbb{P}^2$ defined by the homogenization $\widehat{f}(x, y, z) = x^3y - x^2y^2 - 2xy^3 - 3x^2yz + 2xyz^2 - pz^4 = 0$ is a smooth plane quartic of genus 3, and the given equation $\widehat{f}(x, y, z) = 0$ defines the minimal regular proper semistable model \mathcal{X} for \widehat{X} over \mathbb{Q}_p . The special fiber $\overline{\mathcal{X}}$ of \mathcal{X} consists of four (reduced) lines in general position in \mathbb{P}_k^2 , since $\widehat{f} \bmod p$ factors as $xy(x+y-z)(x-2y-2z)$. The tropicalization $\text{Trop}(X) \subset \mathbb{R}^2$ consists of a triangle with vertices $(0, 0)$, $(1, 0)$, $(0, 1)$ together with three rays emanating from these three vertices in the directions of $(-1, -1)$, $(3, -1)$, $(-1, 3)$ respectively. The three bounded edges/rays incident to $(0, 0)$ all have tropical multiplicity 2, and all other bounded edges/rays in $\text{Trop}(X)$ have tropical multiplicity 1. Let Σ be the skeleton $\Sigma_{\mathcal{X}}$ of \widehat{X} , and let Γ be the minimal skeleton of X . Then Σ is a tetrahedron (with vertices corresponding to the four irreducible components of $\overline{\mathcal{X}}$) with six edges of length 1, because this is the dual graph of a regular semistable model defined over \mathbb{Z}_p , and Γ is obtained from Σ by adding a ray emanating from each vertex of Σ toward the zeros and poles of x and y , namely toward the points $(0 : 1 : 0)$, $(1 : 0 : 0)$, $(2 : 1 : 0)$, $(-1 : 1 : 0)$. The tropicalization map $\text{trop}: X^{\text{an}} \rightarrow \text{Trop}(X) \subset \mathbb{R}^2$ factors through the retraction map $X^{\text{an}} \rightarrow \Gamma$. See Figure 2.

The points of Σ corresponding to the irreducible components $x + y = z$ and $x - 2y = 2z$ of $\overline{\mathcal{X}}$,




FIGURE 2. The skeleton $\Gamma \subset X^{\text{an}}$ and the tropicalization $\text{Trop}(X)$, where X is the curve from Example 2.7. Here x has divisor $3(P_1) - (P_2) - (Q_1) - (Q_2)$ and y has divisor $3(P_2) - (P_1) - (Q_1) - (Q_2)$ on \widehat{X} , where $P_1 = (0 : 1 : 0)$, $P_2 = (1 : 0 : 0)$, $Q_1 = (2 : 1 : 0)$, and $Q_2 = (-1 : 1 : 0)$. The points $A, B, C, D \in \Gamma$ correspond to the irreducible components $x = 0$, $x + y = z$, $y = 0$, and $x - 2y = 2z$, respectively, of \overline{X} . The collapsed segment is \overline{BD} .

as well as the entire edge of Σ connecting these two points, get mapped by trop to the point $(0, 0) \in \text{Trop}(X)$. This edge therefore has expansion factor 0 with respect to trop . The other five bounded edges of $\Sigma \subset \Gamma$ map isometrically (that is, with expansion factor 1) onto their images in $\text{Trop}(X)$. In fact, the tropicalization map is a local isometry everywhere on Γ except along the bounded edge which is contracted to the origin.

Our final example, which is meant to illustrate Theorem 1.4, is a genus 1 curve with multiplicative reduction for which the tropicalization map takes the minimal skeleton isometrically onto its image. This example is also discussed in [KMM09, Example 5.2].

EXAMPLE 2.8. Let K be the completion of the field $\mathbb{C}\{\{t\}\}$ of Puiseux series. Consider the curve E' in \mathbf{G}_m^2 cut out by the equation $f(x, y) = x^2y + xy^2 + \frac{1}{t}xy + x + y$. Its closure in \mathbb{P}^2 is the smooth projective genus 1 curve \widehat{E}' defined by $\widehat{f}(x, y, z) = x^2y + xy^2 + \frac{1}{t}xyz + xz^2 + yz^2$.

Using the description of $\text{Trop}(E')$ as the corner locus of the convex piecewise-linear function associated with f , one sees that $\text{Trop}(E')$ consists of a square with side length 2 plus one ray emanating from each corner of the square; see Figure 3. By restricting f to faces of the Newton complex (see Remark 2.2), one checks that $\text{in}_w(E')$ is reduced and irreducible for every w in $\text{Trop}(E')$. Therefore, by Theorem 1.4 there is a unique graph Γ in the analytification of E' mapping isometrically onto $\text{Trop}(X)$.

In particular, the analytification of E' contains a loop of length 8. One can check by an explicit computation that $\text{val}(j(\widehat{E}')) = -8$, which is consistent with the fact that the analytification of a smooth projective genus 1 curve is either contractible (if the curve has good reduction) or else contains a unique loop of length $-\text{val}(j)$ (if the curve has multiplicative reduction).

3. Admissible algebras and nonarchimedean analytic spaces

Recall that K is an algebraically closed field that is complete with respect to a nontrivial nonarchimedean valuation $\text{val}: K \rightarrow \mathbb{R} \cup \{\infty\}$. Let $|\cdot| = \exp(-\text{val}(\cdot))$ be the associated absolute value.




FIGURE 3. The tropicalization of the elliptic curve \widehat{E}' from Example 2.8. The edges in the square each have lattice length 2.

Let R be the valuation ring of K , let $\mathfrak{m} \subset R$ be its maximal ideal, and let $k = R/\mathfrak{m}$ be its residue field. Choose a nonzero element $\varpi \in \mathfrak{m}$ (as R is not noetherian, it has no uniformizer), so R is ϖ -adically complete. Let $G = \text{val}(K^\times) \subset \mathbb{R}$ be the value group, which is divisible.

In this section we define some notation and collect some results about admissible R -algebras and Raynaud's generic fiber functor that will be needed in the sequel. We refer the reader to [BL93] and [Bos14] for a detailed discussion of admissible formal schemes and the Raynaud generic fiber functor.

3.1 Admissible formal schemes

An *admissible R -algebra* is a topological R -algebra A which is flat and topologically of finite presentation. A formal scheme \mathfrak{X} over $\text{Spf}(R)$ which is locally isomorphic to the formal spectrum of an admissible R -algebra (in its ϖ -adic topology) is called an *admissible formal scheme*. All admissible formal schemes appearing in this paper will be assumed to be quasi-compact and separated. If $\mathfrak{X} = \text{Spf}(A)$, then we write \mathfrak{X}^{an} for the K -affinoid Berkovich analytic space $\mathcal{M}(A \otimes_R K)$. This construction globalizes to give a functor from admissible formal schemes to Berkovich analytic spaces that is compatible with fiber products. The image of a formal scheme under this functor is called the *Raynaud generic fiber*.

An admissible formal scheme \mathfrak{X} with reduced special fiber is called a *formal analytic variety*. If $\text{Spf}(A)$ is a formal affine open subset of a formal analytic variety, then A is reduced and A is equal to the full ring of power-bounded elements in $A_K := A \otimes_R K$ by Proposition 3.13. The canonical reduction \widetilde{A}_K of A_K therefore coincides with $A \otimes_R k$, so if $\text{Spf}(B)$ is a formal affine open subset of $\text{Spf}(A)$, then $\text{Spec}(\widetilde{B}_K)$ is an affine open subset of $\text{Spec}(\widetilde{A}_K)$. (Our definition of a formal analytic variety differs from the original one given in [BL85], but the above argument shows that the two definitions are equivalent in our situation.)

NOTATION. Let \mathfrak{X} be an R -scheme or a formal R -scheme. We denote the special fiber $\mathfrak{X} \otimes_R k$ of \mathfrak{X} by $\overline{\mathfrak{X}}$.

3.2 Reductions of analytic spaces

Let \mathcal{A} be a K -affinoid algebra. We denote by $\mathring{\mathcal{A}}$, $\check{\mathcal{A}}$, and $\widetilde{\mathcal{A}} = \mathring{\mathcal{A}}/\check{\mathcal{A}}$ the subring of power-bounded elements, the ideal of topologically nilpotent elements, and the canonical reduction, respectively. Setting $\mathcal{X} = \mathcal{M}(\mathcal{A})$, there is a *reduction map* $\text{red}: \mathcal{X} \rightarrow \widetilde{\mathcal{X}} := \text{Spec}(\widetilde{\mathcal{A}})$. By [Ber90, Corollary 2.4.2 and Proposition 2.4.4] this map is surjective and anti-continuous, in the sense that

the inverse image of an open subset is closed. Similarly, if \mathfrak{X} is an admissible formal scheme over $\mathrm{Spf}(R)$, then there is a canonical surjective and anti-continuous *reduction map* $\mathrm{red}: \mathfrak{X}^{\mathrm{an}} \rightarrow \overline{\mathfrak{X}}$ which coincides with the map defined above when $\mathfrak{X} = \mathrm{Spf}(\mathring{\mathcal{A}})$. In particular, $\overline{\mathfrak{X}}$ is connected if $\mathfrak{X}^{\mathrm{an}}$ is connected. The inverse image of a closed point of $\overline{\mathfrak{X}}$ under the reduction map is called a *formal fiber*.

3.3 Shilov points

The *Shilov boundary* of a K -affinoid space $\mathcal{X} = \mathcal{M}(\mathcal{A})$ is defined to be the smallest closed subset $\Gamma(\mathcal{X}) \subset \mathcal{X}$ such that every function $|f|$ for $f \in \mathcal{A}$ attains its maximum at a point of $\Gamma(\mathcal{X})$. Let \mathfrak{X} be a formal analytic variety over $\mathrm{Spf}(R)$ with Raynaud generic fiber \mathcal{X} . If η is a generic point of $\overline{\mathfrak{X}}$, then there is a unique preimage of η under red which we call the *Shilov point* x_η associated with η . The residue field of η is isomorphic to $\widetilde{\mathcal{H}}(x_\eta)$. If $\mathcal{X} = \mathcal{M}(\mathcal{A})$ is affinoid, then $\Gamma(\mathcal{X})$ is the set of Shilov points of $\mathfrak{X} = \mathrm{Spf}(\mathring{\mathcal{A}})$.

3.4 Analytic curves

Following [Thu05, § 2.1.3], we define a (strictly) *analytic curve* over K to be a (good) K -analytic space which is paracompact, of pure dimension 1, and without boundary. The analytification of an algebraic curve over K (by which we mean a 1-dimensional separated integral scheme of finite type over K) is always an analytic curve in this sense.

If \mathcal{X} is an analytic curve and $\mathcal{V} \subset \mathcal{X}$ is an affinoid domain, then by [Ber90, Proposition 3.1.3] and [Thu05, Proposition 2.1.12] the following three (finite) sets coincide: (i) the topological boundary $\partial_{\mathrm{top}} \mathcal{V}$ of \mathcal{V} in \mathcal{X} , (ii) the boundary $\partial \mathcal{V}$ of \mathcal{V} in the sense of [Ber90, § 2.5.7], and (iii) the Shilov boundary $\Gamma(\mathcal{V})$ of \mathcal{V} .

3.5 Types of points in an analytic curve

Let x be a point in a K -analytic curve \mathcal{X} , and let $\mathcal{H}(x)$ be its completed residue field. The extension $\mathcal{H}(x)/k$ has transcendence degree $s(x) \leq 1$, and the abelian group $|\mathcal{H}(x)^\times|/|K^\times|$ has rank $t(x) \leq 1$. Moreover, the integers $s(x)$ and $t(x)$ must satisfy the *Abhyankar inequality* [Vaq00, Theorem 9.2]

$$s(x) + t(x) \leq 1.$$

Using the terminology from [Ber90] and [Ber93] (see also [Thu05, § 2.1]), we say that x is *type 2* if $s(x) = 1$ and *type 3* if $t(x) = 1$. If $s(x) = t(x) = 0$, then x is called *type 1* if $\mathcal{H}(x) = K$ and *type 4* otherwise. Points of type 4 will not play any significant role in this paper. We define

$$\begin{aligned} \mathbf{H}_o(\mathcal{X}) &= \{\text{all points of } \mathcal{X} \text{ of types 2 and 3}\}, \\ \mathbf{H}(\mathcal{X}) &= \{\text{all points of } \mathcal{X} \text{ of types 2, 3, and 4}\}. \end{aligned}$$

We call $\mathbf{H}_o(\mathcal{X})$ the set of *skeletal points*, because it is the union of all skeleta of admissible formal models of \mathfrak{X} (see [BPR13, Corollary 5.1]), and $\mathbf{H}(\mathcal{X})$ the set of *norm points* of \mathcal{X} , because it is the set of all points corresponding to norms on the function field $K(X)$ that extend the given norm on K . If $\mathcal{X} = X^{\mathrm{an}}$ is the analytification of an algebraic curve X over K , then $X(K) \subset X^{\mathrm{an}}$ is naturally identified with the set of type 1 points of X^{an} , so $\mathbf{H}(X^{\mathrm{an}}) = X^{\mathrm{an}} \setminus X(K)$. (Recall that we are assuming throughout this discussion that K is algebraically closed.)

3.6 Some facts about admissible R -algebras

The following fact is standard and is easily proved using the results of [BL93, § 1].

PROPOSITION 3.7. (i) *If A is a finitely presented and flat R -algebra, then its ϖ -adic completion \widehat{A} is an admissible R -algebra.*

(ii) *If $f: A \twoheadrightarrow B$ is a surjective homomorphism of finitely presented and flat R -algebras with kernel \mathfrak{a} , then $\widehat{f}: \widehat{A} \twoheadrightarrow \widehat{B}$ is a surjection of admissible R -algebras with kernel $\mathfrak{a}\widehat{A}$.*

We set the following notation, which we will use until Section 3.15: A and B will denote admissible R -algebras, $\overline{A} = A \otimes_R k$ and $\overline{B} = B \otimes_R k$ their reductions, and $A_K = A \otimes_R K$ and $B_K = B \otimes_R K$ the associated K -affinoid algebras. We let $\mathfrak{X} = \mathrm{Spf}(A)$, $\mathfrak{Y} = \mathrm{Spf}(B)$, $\overline{\mathfrak{X}} = \mathrm{Spec}(\overline{A})$, and $\overline{\mathfrak{Y}} = \mathrm{Spec}(\overline{B})$. Let $f: A \rightarrow B$ be a homomorphism, let $\overline{f}: \overline{A} \rightarrow \overline{B}$ and $f_K: A_K \rightarrow B_K$ be the induced homomorphisms, and let $\phi: \mathfrak{Y} \rightarrow \mathfrak{X}$ and $\overline{\phi}: \overline{\mathfrak{Y}} \rightarrow \overline{\mathfrak{X}}$ be the induced morphisms.

PROPOSITION 3.8. (i) *The homomorphism f is flat if and only if $\overline{f}: \overline{A} \rightarrow \overline{B}$ is flat.*

(ii) *The homomorphism f is finite if and only if $f_K: A_K \rightarrow B_K$ is finite.*

Proof. The ‘only if’ directions are clear. Suppose that \overline{f} is flat. By [BL93, Lemma 1.6], it suffices to show that $f_n: A_n \rightarrow B_n$ is flat for all $n \geq 0$, where $A_n = A/\varpi^{n+1}A$ and $B_n = B/\varpi^{n+1}B$. But A_n and B_n are of finite presentation and flat over $R_n = R/\varpi^{n+1}R$, so f_n is flat by the fibral flatness criterion [GD66, Corollaire 11.3.1].

Now suppose that f_K is finite. Choose a surjection $\mathring{T}_n \twoheadrightarrow A$. The induced homomorphism $T_n \twoheadrightarrow A \otimes_R K \rightarrow B \otimes_R K$ is finite, so by [BGR84, Theorem 6.3.5/1] the composition $\mathring{T}_n \twoheadrightarrow A \rightarrow (B \otimes_R K)^\circ$ is integral. Hence $A \rightarrow (B \otimes_R K)^\circ$ is integral, so $A \rightarrow B$ is integral since $B \subset (B \otimes_R K)^\circ$. Then $f_n: A/\varpi^{n+1}A \rightarrow B/\varpi^{n+1}B$ is of finite type and integral for all $n \geq 0$, so f_n is finite, and therefore f is finite by [BL93, Lemma 1.5]. \square

COROLLARY 3.9. *Suppose that $f_K: A_K \rightarrow B_K$ is finite and dominant, that is, that $\ker(f_K)$ is nilpotent. Then $\overline{f}: \overline{A} \rightarrow \overline{B}$ is finite and $\overline{\phi}: \overline{\mathfrak{Y}} \rightarrow \overline{\mathfrak{X}}$ is surjective.*

Proof. Since $A \subset A_K$ and $B \subset B_K$, we have that $\ker(f)$ is nilpotent, and f is finite by Proposition 3.8. Hence $\mathrm{Spec}(B) \rightarrow \mathrm{Spec}(A)$ is surjective, so $\overline{\mathfrak{Y}} \rightarrow \overline{\mathfrak{X}}$ is surjective. Finiteness of f implies finiteness of \overline{f} . \square

We say that a ring is *equidimensional of dimension d* provided that every maximal ideal has height d . Let \mathcal{A} be a K -affinoid algebra, and let $\mathcal{X} = \mathcal{M}(\mathcal{A})$. Then \mathcal{A} is equidimensional of dimension d if and only if $\dim(\mathcal{O}_{\mathcal{X},x}) = d$ for every $x \in \mathrm{MaxSpec}(\mathcal{A})$ by [BGR84, Proposition 7.3.2/8]. In particular, if $\mathcal{M}(\mathcal{B})$ is an affinoid domain in $\mathcal{M}(\mathcal{A})$ and \mathcal{A} is equidimensional of dimension d , then so is \mathcal{B} .

PROPOSITION 3.10. *If A_K is equidimensional of dimension d , then \overline{A} is equidimensional of dimension d .*

Proof. Replacing \mathfrak{X} with an irreducible formal affine open subset, we may assume that \mathfrak{X} is irreducible. Let $R\langle x_1, \dots, x_n \rangle \twoheadrightarrow A$ be a presentation of A . By Noether normalization [BGR84, Theorem 6.1.2/1] we can choose the x_i such that $K\langle x_1, \dots, x_d \rangle \rightarrow A_K$ is finite and injective, where $d = \dim(A_K)$. Then $\overline{\mathfrak{X}} \rightarrow \mathbb{A}_k^d$ is finite and surjective by Corollary 3.9. \square

COROLLARY 3.11. *Suppose that $f_K: A_K \rightarrow B_K$ is finite and dominant, and that A_K and B_K are equidimensional (necessarily of the same dimension). Then $\overline{\phi}: \overline{\mathfrak{Y}} \rightarrow \overline{\mathfrak{X}}$ is finite and surjective, and the image of an irreducible component of $\overline{\mathfrak{Y}}$ is an irreducible component of $\overline{\mathfrak{X}}$.*

Proof. This follows immediately from Proposition 3.10 and Corollary 3.9. \square

The following theorem uses the fact that K is algebraically closed in an essential way. It can be found in [BL85, Proposition 1.1].

THEOREM 3.12. *Let \mathcal{A} be a K -affinoid algebra. Then $\mathring{\mathcal{A}}$ is admissible if and only if \mathcal{A} is reduced.*

Proof. Since $\mathring{\mathcal{A}}$ is always R -flat, by [BL93, Proposition 1.1(c)] the issue is whether $\mathring{\mathcal{A}}$ is topologically finitely generated. Suppose that \mathcal{A} is reduced. By [BGR84, Theorem 6.4.3/1] there is a surjection $T_n \twoheadrightarrow \mathcal{A}$ such that the residue norm on \mathcal{A} agrees with the supremum norm; then by Proposition 6.4.3/3(i) of [BGR84] the induced homomorphism $\mathring{T}_n \rightarrow \mathring{\mathcal{A}}$ is surjective. The converse follows in a similar way from Theorem 6.4.3/1 and Corollary 6.4.3/6 of [BGR84]. \square

PROPOSITION 3.13. *The ring \overline{A} is reduced if and only if $A = \mathring{A}_K$, in which case A is reduced.*

Proof. If $A = \mathring{A}_K$, then A is reduced by Theorem 3.12, so $\overline{A} = \widetilde{A}_K$ is reduced. Conversely, suppose that \overline{A} is reduced. Let $\alpha: \mathring{T}_n \twoheadrightarrow A$ be a surjection. Since the \mathring{T}_n -ideal $\mathring{T}_n + \ker(\alpha) = \mathfrak{m}\mathring{T}_n + \ker(\alpha)$ is the kernel of the composite homomorphism $\mathring{T}_n \rightarrow A \rightarrow \overline{A}$, it is a reduced ideal; hence by [BGR84, Propositions 6.4.3/4, 6.4.3/3(i)] we have $A = \alpha(\mathring{T}_n) = \mathring{A}_K$. \square

COROLLARY 3.14. *If \overline{A} is an integral domain, then A_K is an integral domain and $|\cdot|_{\text{sup}}$ is multiplicative.*

Proof. By Proposition 3.13 we have $\widetilde{A}_K = \overline{A}$, so the result follows from [BGR84, Proposition 6.2.3/5]. \square

3.15 Finite morphisms of pure degree

In general there is not a good notion of the ‘degree’ of a finite morphism $Y \rightarrow X$ between noetherian schemes when X is not irreducible, since the degree of the induced map on an irreducible component of X can vary from component to component. The notion of a morphism having ‘pure degree’ essentially means that the degree is the same on every irreducible component of X . This notion is quite well behaved in that it respects analytification of algebraic varieties and of admissible formal schemes. The definition of a morphism of pure degree is best formulated in the language of fundamental cycles. We refer to [Tho90] for a review of the theory of cycles on a noetherian scheme which is not necessarily of finite type over a field.

3.16 Let X be a noetherian scheme. A *cycle* on X is a finite formal sum $\sum_W n_W \cdot W$, where $n_W \in \mathbb{Z}$ and W ranges over the irreducible closed subsets of X . The group of cycles on X is denoted $C(X)$. The *fundamental cycle* of X is the cycle

$$[X] = \sum_{\zeta} \text{length}_{\mathcal{O}_{X,\zeta}}(\mathcal{O}_{X,\zeta}) \cdot \overline{\{\zeta\}},$$

where the sum is taken over all generic points of X . We define pushforwards and pullbacks as in intersection theory. These satisfy the usual properties; see [Tho90, Lemmas 2.4, 2.5, and 4.8].

DEFINITION 3.17. Let $f: Y \rightarrow X$ be a finite morphism of noetherian schemes. We say that f has *pure degree* δ and we write $[Y : X] = \delta$ provided that $f_*[Y] = \delta[X]$; here $\delta \in \mathbb{Q}$ need not be an integer.

Remark 3.18. Let $f: Y \rightarrow X$ be a finite morphism of noetherian schemes.

(i) If X is irreducible and every generic point of Y maps to the generic point of X , then f automatically has a pure degree, which we simply call the *degree* of f . Moreover, if X is integral with generic point ζ , then the degree of f is the dimension of $\Gamma(f^{-1}(\zeta), \mathcal{O}_{f^{-1}(\zeta)})$ as a vector space over the function field $\mathcal{O}_{X,\zeta}$. In particular, if f is a finite and dominant morphism of integral schemes, then the (pure) degree of f is the degree of the extension of function fields.

(ii) Let ζ be a generic point of X , and let $C = \overline{\{\zeta\}}$ be the corresponding irreducible component. Define the *multiplicity of C in X* to be the quantity

$$\text{mult}_X(C) = \text{length}_{\mathcal{O}_{X,\zeta}}(\mathcal{O}_{X,\zeta}),$$

so $[X] = \sum_C \text{mult}_X(C) \cdot C$. It follows that f has pure degree δ if and only if (i) every irreducible component D of Y maps to an irreducible component of X , and (ii) for every irreducible component C of X we have

$$\delta \text{mult}_X(C) = \sum_{D \rightarrow C} \text{mult}_Y(D)[D : C], \quad (3.18.1)$$

where $[D : C]$ is the usual degree of a finite morphism of integral schemes.

(iii) Let $g: X \rightarrow Z$ be another finite morphism of noetherian schemes. Suppose that f has pure degree δ and g has pure degree ϵ . Then $g \circ f$ has pure degree $\delta\epsilon$.

PROPOSITION 3.19. *Let X, Y, X' be noetherian schemes, let $f: Y \rightarrow X$ be a finite morphism, let $g: X' \rightarrow X$ be a flat morphism, let $Y' = Y \times_X X'$, and let $f': Y' \rightarrow X'$ be the projection.*

- (i) *If f has pure degree δ , then f' has pure degree δ .*
- (ii) *If g is surjective, then f has pure degree δ if and only if f' has pure degree δ .*

Proof. Let $h: Y' \rightarrow Y$ be the other projection, so h is flat. We have

$$f'_*[Y'] = f'_*h^*[Y] = g^*f_*[Y] = \delta g^*[X] = \delta[X'],$$

which proves part (i). Conversely, suppose that g is surjective (and flat) and that $f'_*[Y'] = \delta[X']$. Then

$$g^*f_*[Y] = f'_*h^*[Y] = f'_*[Y'] = \delta[X'] = g^*(\delta[X]),$$

so we are done because g^* is visibly injective in this situation. \square

3.20 Next we will define pure-degree morphisms of analytic spaces. As above, we must first review the notion of the fundamental cycle of an analytic space, as defined by Gubler [Gub98, § 2].

Let \mathcal{X} be a K -analytic space (assumed from now on to be Hausdorff and paracompact). A *Zariski-closed subspace* of \mathcal{X} is by definition an isomorphism class of closed immersions $\mathcal{V} \hookrightarrow \mathcal{X}$. A Zariski-closed subspace of \mathcal{X} is *irreducible* if it cannot be expressed as a union of two proper Zariski-closed subspaces. Gubler [Gub98, § 2] defines a *cycle* on \mathcal{X} to be a locally finite formal sum $\sum_{\mathcal{V}} n_{\mathcal{V}} \mathcal{V}$, where $n_{\mathcal{V}} \in \mathbb{Z}$ and \mathcal{V} ranges over the irreducible Zariski-closed subspaces of \mathcal{X} ; ‘locally finite’ means that there exists an admissible covering of \mathcal{X} by affinoid domains intersecting only finitely many \mathcal{V} with $n_{\mathcal{V}} \neq 0$. Let $C(\mathcal{X})$ denote the group of cycles on \mathcal{X} .

3.20.1 If $\mathcal{X} = \mathcal{M}(\mathcal{A})$ is affinoid, then the Zariski-closed subspaces of \mathcal{X} are in natural inclusion-reversing bijection with the ideals of \mathcal{A} ; therefore we have an identification $C(\mathcal{X}) = C(\text{Spec}(\mathcal{A}))$, which we will make implicitly from now on.

3.20.2 There are natural proper pushforward and flat pullback homomorphisms for cycles on analytic spaces, which satisfy the expected properties. There is a canonical *fundamental cycle* $[\mathcal{X}] \in C(\mathcal{X})$ which is uniquely determined by the property that for every affinoid domain $\iota: \mathcal{M}(\mathcal{A}) \hookrightarrow \mathcal{X}$, we have $\iota^*[\mathcal{X}] = [\mathcal{M}(\mathcal{A})] = [\text{Spec}(\mathcal{A})]$. See [Gub98, §§ 2.6, 2.7, 2.8, and Proposition 2.12].

DEFINITION 3.21. Let $f: \mathcal{Y} \rightarrow \mathcal{X}$ be a finite morphism of K -analytic spaces. We say that f has *pure degree* δ and we write $[\mathcal{Y} : \mathcal{X}] = \delta$ provided that $f_*[\mathcal{Y}] = \delta[\mathcal{X}]$. Again $\delta \in \mathbb{Q}$ need not be an integer.

Remark 3.22. Let $f: \mathcal{Y} \rightarrow \mathcal{X}$ be a finite morphism of K -analytic spaces.

(i) If $\mathcal{X} = \mathcal{M}(\mathcal{A})$ and $\mathcal{Y} = \mathcal{M}(\mathcal{B})$ are affinoid, then $f: \mathcal{M}(\mathcal{B}) \rightarrow \mathcal{M}(\mathcal{A})$ has pure degree δ if and only if the map of affine schemes $\mathrm{Spec}(\mathcal{B}) \rightarrow \mathrm{Spec}(\mathcal{A})$ has pure degree δ .

(ii) If f has pure degree δ and $g: \mathcal{X} \rightarrow \mathcal{Z}$ is a finite morphism of analytic spaces of pure degree ϵ , then $g \circ f$ has pure degree $\delta\epsilon$.

PROPOSITION 3.23. Let $f: \mathcal{Y} \rightarrow \mathcal{X}$ be a finite morphism of K -analytic spaces.

(i) If f has pure degree δ , $\mathcal{M}(\mathcal{A}) \subset \mathcal{X}$ is an affinoid domain, and $\mathcal{M}(\mathcal{B}) = f^{-1}(\mathcal{M}(\mathcal{A}))$, then $\mathcal{M}(\mathcal{B}) \rightarrow \mathcal{M}(\mathcal{A})$ has pure degree δ .

(ii) If there exists an admissible cover $\mathcal{X} = \bigcup_i \mathcal{M}(\mathcal{A}_i)$ of \mathcal{X} by affinoid domains such that $\mathcal{M}(\mathcal{B}_i) = f^{-1}(\mathcal{M}(\mathcal{A}_i)) \rightarrow \mathcal{M}(\mathcal{A}_i)$ has pure degree δ for each i , then f has pure degree δ .

Proof. Since the inclusion $\mathcal{M}(\mathcal{A}) \hookrightarrow \mathcal{X}$ is flat, the first part follows as in the proof of Proposition 3.19(i). In situation (ii), let $f_i = f|_{\mathcal{M}(\mathcal{B}_i)}: \mathcal{M}(\mathcal{B}_i) \rightarrow \mathcal{M}(\mathcal{A}_i)$, and assume $(f_i)_*[\mathcal{M}(\mathcal{B}_i)] = \delta[\mathcal{M}(\mathcal{A}_i)]$ for all i . Arguing as in the proof of Proposition 3.19(i), we see that the pullback of $f_*[\mathcal{Y}]$ to $\mathcal{M}(\mathcal{A}_i)$ is equal to $\delta[\mathcal{M}(\mathcal{A}_i)]$ for all i ; since $[\mathcal{X}]$ is the unique cycle which pulls back to $[\mathcal{M}(\mathcal{A}_i)]$ for all i , this shows $f_*[\mathcal{Y}] = \delta[\mathcal{X}]$. \square

The property of being a finite morphism of pure degree is compatible with analytification.

PROPOSITION 3.24. Let $f: Y \rightarrow X$ be a morphism of finite-type K -schemes. Then f is finite of pure degree δ if and only if $f^{\mathrm{an}}: Y^{\mathrm{an}} \rightarrow X^{\mathrm{an}}$ is finite of pure degree δ .

Proof. By [Con99, Theorem A.2.1], f is finite if and only if f^{an} is finite. Hence we may assume that $X = \mathrm{Spec}(A)$ and $Y = \mathrm{Spec}(B)$ are affine. If $\mathcal{M}(\mathcal{A}) \subset X^{\mathrm{an}}$ is an affinoid domain, then $\mathrm{Spec}(A) \rightarrow \mathrm{Spec}(A)$ is flat by Lemma A.1.2 of [Con99], and if $\{\mathcal{M}(\mathcal{A}_i)\}_{i \in I}$ is an admissible covering of X^{an} , then $\coprod_{i \in I} \mathrm{Spec}(\mathcal{A}_i) \rightarrow \mathrm{Spec}(A)$ is flat and surjective. Let $\mathcal{M}(\mathcal{B}_i) = f^{-1}(\mathcal{M}(\mathcal{A}_i))$. We claim that $\mathcal{B}_i = B \otimes_A \mathcal{A}_i$. Since $B \otimes_A \mathcal{A}_i$ is finite over \mathcal{A}_i , it is affinoid by [BGR84, Proposition 6.1.1/6], so the claim follows easily from the universal property of the analytification (see also [Con99, § A.2]). Hence by Proposition 3.19(ii), f has pure degree δ if and only if $\mathrm{Spec}(\mathcal{B}_i) \rightarrow \mathrm{Spec}(\mathcal{A}_i)$ has pure degree δ for each i ; by Remark 3.22(i), this is the case if and only if $\mathcal{M}(\mathcal{B}_i) \rightarrow \mathcal{M}(\mathcal{A}_i)$ has pure degree δ for each i , which is equivalent to f^{an} having pure degree δ by Proposition 3.23(ii). \square

The following counterpart to Proposition 3.24 allows us to compare the degrees of the generic and special fibers of a finite morphism of admissible formal schemes. It will play a key role throughout this paper.

PROPOSITION 3.25 (Projection formula). Let $f: \mathfrak{Y} \rightarrow \mathfrak{X}$ be a finite morphism of admissible formal schemes, and let $f^{\mathrm{an}}: \mathfrak{Y}^{\mathrm{an}} \rightarrow \mathfrak{X}^{\mathrm{an}}$ and $\bar{f}: \bar{\mathfrak{Y}} \rightarrow \bar{\mathfrak{X}}$ be the induced morphisms on the generic and special fibers, respectively. If f^{an} has pure degree δ , then \bar{f} has pure degree δ .

Proof. The theory of cycles on analytic spaces discussed above is part of Gubler's more general intersection theory on admissible formal schemes, and our 'projection formula' is in fact a special case of Gubler's projection formula [Gub98, Proposition 4.5]; this can be seen as follows. Choose any $\varpi \in K^\times$ with $\mathrm{val}(\varpi) \in (0, \infty)$, and let D be the Cartier divisor on \mathfrak{X} defined by ϖ .

Essentially by definition (cf. (3.8) and (3.10) of [Gub98]) the intersection product $D.[\mathfrak{X}^{\text{an}}]$ is equal to $\text{val}(\varpi)[\bar{\mathfrak{X}}]$, and likewise $(f^*D).[\mathfrak{Y}^{\text{an}}] = \text{val}(\varpi)[\bar{\mathfrak{Y}}]$. Hence if $f_*^{\text{an}}[\mathfrak{Y}^{\text{an}}] = \delta[\mathfrak{X}^{\text{an}}]$, then

$$\text{val}(\varpi)\bar{f}_*[\bar{\mathfrak{Y}}] = f_*((f^*D).[\mathfrak{Y}^{\text{an}}]) = D.f_*^{\text{an}}[\mathfrak{Y}^{\text{an}}] = D.(\delta[\mathfrak{X}^{\text{an}}]) = \text{val}(\varpi)\delta[\bar{\mathfrak{X}}],$$

where the second equality is by Gubler's projection formula. Canceling the factors of $\text{val}(\varpi)$ yields Proposition 3.25. \square

Remark 3.26. The converse to Proposition 3.25 does not hold in general. The following example is due to Gubler: let $\mathfrak{X} = \text{Spf}(R[x]/(x(x - \varpi)))$ and $\mathfrak{Y} = \mathfrak{X} \amalg \text{Spf}(R)$, and let $f: \mathfrak{Y} \rightarrow \mathfrak{X}$ be the map which is the identity on \mathfrak{X} and which maps $\text{Spf}(R)$ to \mathfrak{X} via $x \mapsto 0$. Then f^{an} does not have a pure degree, but \bar{f} does since $\bar{\mathfrak{X}}$ is a point.

3.27 Here we note some special cases of the projection formula:

(i) Suppose that $\mathfrak{X} = \text{Spf}(A)$ and $\mathfrak{Y} = \text{Spf}(B)$, and that A is an integral domain with fraction field Q . If all generic points of $\text{Spec}(B \otimes_R K)$ map to the generic point of $\text{Spec}(A \otimes_R K)$, then $\mathcal{M}(B \otimes_R K) \rightarrow \mathcal{M}(A \otimes_R K)$ is finite with pure degree equal to $\dim_Q(B \otimes_A Q)$. By (3.18.1), for every irreducible component $\bar{\mathfrak{C}}$ of $\bar{\mathfrak{X}}$ we have

$$\dim_Q(B \otimes_A Q) \cdot \text{mult}_{\bar{\mathfrak{X}}}(\bar{\mathfrak{C}}) = \sum_{\bar{\mathfrak{D}} \rightarrow \bar{\mathfrak{C}}} \text{mult}_{\bar{\mathfrak{Y}}}(\bar{\mathfrak{D}}) \cdot [\bar{\mathfrak{D}} : \bar{\mathfrak{C}}],$$

where the sum is taken over all irreducible components $\bar{\mathfrak{D}}$ of $\bar{\mathfrak{Y}}$ that surject onto $\bar{\mathfrak{C}}$.

(ii) Suppose that $f^{\text{an}}: \mathfrak{Y}^{\text{an}} \rightarrow \mathfrak{X}^{\text{an}}$ is an isomorphism. Then for every irreducible component $\bar{\mathfrak{C}}$ of $\bar{\mathfrak{X}}$ we have

$$\text{mult}_{\bar{\mathfrak{X}}}(\bar{\mathfrak{C}}) = \sum_{\bar{\mathfrak{D}} \rightarrow \bar{\mathfrak{C}}} \text{mult}_{\bar{\mathfrak{Y}}}(\bar{\mathfrak{D}}) \cdot [\bar{\mathfrak{D}} : \bar{\mathfrak{C}}],$$

where the sum is taken over all irreducible components $\bar{\mathfrak{D}}$ of $\bar{\mathfrak{Y}}$ that surject onto $\bar{\mathfrak{C}}$, because an isomorphism has pure degree 1.

4. Tropical integral models

We continue to assume that K is an algebraically closed field which is complete with respect to a nontrivial nonarchimedean valuation.

NOTATION 4.1. Let $M \cong \mathbb{Z}^n$ be a lattice, with dual lattice $N = \text{Hom}(M, \mathbb{Z})$. If H is an additive subgroup of \mathbb{R} , we write M_H for $M \otimes_{\mathbb{Z}} H$, so N_H is naturally identified with $\text{Hom}(M, H)$. We write $\langle \cdot, \cdot \rangle$ to denote the canonical pairings $M \times N \rightarrow \mathbb{Z}$ and $M_{\mathbb{R}} \times N_{\mathbb{R}} \rightarrow \mathbb{R}$.

Let $\mathbf{T} = \text{Spec } K[M]$ be the torus over K with character lattice M . For u in M , we write x^u for the corresponding character, considered as a function in $K[M]$.

4.2 Extended tropicalization

A point $\|\cdot\|$ in \mathbf{T}^{an} naturally determines a real-valued linear function on the character lattice M , taking u to $-\log \|x^u\|$. The induced tropicalization map $\text{trop}: \mathbf{T}^{\text{an}} \rightarrow N_{\mathbb{R}}$ is continuous, proper, and surjective [Pay09a]. The image of $\mathbf{T}(K)$ is exactly N_G , which is dense in $N_{\mathbb{R}}$ because G is nontrivial and divisible.

More generally, if σ is a pointed rational polyhedral cone in $N_{\mathbb{R}}$ and $Y_{\sigma} = \text{Spec } K[\sigma^{\vee} \cap M]$ is the associated affine toric variety with dense torus \mathbf{T} , then there is a natural tropicalization

map from Y_σ to the space of additive semigroup homomorphisms $\text{Hom}(\sigma^\vee \cap M, \mathbb{R} \cup \{\infty\})$ taking a point $\|\cdot\|$ to the semigroup map $u \mapsto -\log \|x^u\|$, where $-\log(0)$ is defined to be ∞ . See [Pay09a, Rab12] for further details. We write $N_{\mathbb{R}}(\sigma)$ for the image of Y_σ^{an} under this extended tropicalization map.

DEFINITION 4.3. We say that a point in $N_{\mathbb{R}}(\sigma)$ is G -rational if it is in the subspace $\text{Hom}(\sigma^\vee \cap M, G \cup \{\infty\})$. Note that the image of any K -rational point of Y_σ is G -rational.

For any toric variety Y_Δ , the tropicalization $N_{\mathbb{R}}(\Delta)$ is the union of the spaces $N_{\mathbb{R}}(\sigma)$ for σ in Δ , glued along the open inclusions $N_{\mathbb{R}}(\tau) \subset N_{\mathbb{R}}(\sigma)$ for $\tau \preceq \sigma$. The tropicalization maps on torus-invariant affine opens are compatible with this gluing, and together give a natural continuous, proper, and surjective map of topological spaces $\text{trop}: Y_\Delta^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta)$. Note that the vector space $N_{\mathbb{R}}$, which is the tropicalization of the dense torus $\mathbf{T} \subset Y_\Delta$, is open and dense in $N_{\mathbb{R}}(\Delta)$. For the purpose of constructing tropical integral models of toric varieties and their subvarieties, it will generally suffice to study polyhedral complexes in $N_{\mathbb{R}}$.

Let X be a closed subscheme of Y_Δ . The tropicalization $\text{Trop}(X)$ is the image of X^{an} under trop . Since $X(K)$ is dense in X^{an} , its image is dense in $\text{Trop}(X)$. Furthermore, every G -rational point of $\text{Trop}(X)$ is the image of a point of $X(K)$, and if X is irreducible, then the preimage of any point in $\text{Trop}(X) \cap N_G$ is Zariski dense in X . See [Pay09b, Corollary 4.2] and [Pay12, Remark 2], [Gub13, Proposition 4.14], or [OP13, Theorem 4.2.5].

4.4 Polyhedral domains

Recall that the *recession cone* σ_P of a nonempty polyhedron $P \subset N_{\mathbb{R}}$ is the set of those v in $N_{\mathbb{R}}$ such that $w + v$ is in P whenever w is in P . If P is the intersection of the halfspaces $\langle u_1, v \rangle \geq a_1, \dots, \langle u_r, v \rangle \geq a_r$, then σ_P is the dual of the cone in $M_{\mathbb{R}}$ spanned by u_1, \dots, u_r . In particular, if P is an integral G -affine polyhedron, then these halfspaces can be chosen with each u_i in M , so the recession cone σ_P is a rational polyhedral cone. The recession cone can also be characterized as the intersection with $N_{\mathbb{R}} \times \{0\}$ of the closure in $N_{\mathbb{R}} \times \mathbb{R}$ of the cone spanned by $P \times \{1\}$.

Let P be an integral G -affine polyhedron in $N_{\mathbb{R}}$ that does not contain any positive-dimensional affine linear subspace, so its recession cone $\sigma = \sigma_P$ is pointed.

DEFINITION 4.5. The *polyhedral domain* associated with P is the inverse image of the closure of P in $N_{\mathbb{R}}(\sigma)$ under $\text{trop}: Y_\sigma^{\text{an}} \rightarrow N_{\mathbb{R}}(\sigma)$ and is denoted \mathcal{U}^P .

These polyhedral domains, introduced in [Rab12], directly generalize the polytopal domains studied by Gubler in [Gub07b]. Indeed, a polytopal domain is the preimage in \mathbf{T}^{an} of an integral G -affine polytope in $N_{\mathbb{R}}$. Since the recession cone of a polytope in $N_{\mathbb{R}}$ is the zero cone, whose associated toric variety is \mathbf{T} , Gubler's polytopal domains are exactly the special case of these polyhedral domains where P is bounded.

By [Rab12, § 6] the polyhedral domain \mathcal{U}^P is an affinoid domain in Y_σ^{an} with coordinate ring

$$K\langle \mathcal{U}^P \rangle = \left\{ \sum_{u \in \sigma^\vee \cap M} a_u x^u : \lim(\text{val}(a_u) + \langle u, v \rangle) = \infty \text{ for all } v \in P \right\},$$

where the limit is taken over all complements of finite sets. Its supremum norm is given by

$$\left| \sum a_u x^u \right|_{\text{sup}} = \sup_{\substack{u \in \sigma^\vee \cap M \\ v \in P}} |a_u| \exp(-\langle u, v \rangle).$$

Since the recession cone σ is pointed, the polyhedron P contains no linear subspace and hence has vertices. The supremum above is always achieved at one of the vertices of P , so the ring of power-bounded regular functions on \mathcal{U}^P is

$$K\langle\mathcal{U}^P\rangle^\circ = \left\{ \sum_{u \in \sigma^\vee \cap M} a_u x^u \in K\langle\mathcal{U}^P\rangle : \text{val}(a_u) + \langle u, v \rangle \geq 0 \text{ for all } v \in \text{vert}(P) \right\}.$$

Since $K\langle\mathcal{U}^P\rangle$ is reduced, Theorem 3.12 implies that $\mathfrak{U}^P = \text{Spf}(K\langle\mathcal{U}^P\rangle^\circ)$ is an admissible formal scheme with analytic generic fiber \mathcal{U}^P .

Remark 4.6. If P is integral affine but not G -affine, then the inverse image \mathcal{U}^P of the closure of P under trop is a nonstrict affinoid domain. Indeed, if K' is a complete valued field extension of K whose value group G' is large enough that P is G' -affine, then $\mathcal{U}^P \widehat{\otimes}_K K'$ is strictly K' -affinoid.

4.7 Polyhedral integral models

Let P be an integral G -affine polyhedron in $N_{\mathbb{R}}$ whose recession cone $\sigma = \sigma_P$ is pointed. As usual, we let $Y_\sigma = \text{Spec } K[\sigma^\vee \cap M]$ denote the associated affine toric variety with dense torus \mathbf{T} .

DEFINITION 4.8. We define $R[Y^P] \subset K[\sigma^\vee \cap M]$ to be the subring consisting of those Laurent polynomials $\sum a_u x^u$ such that $\text{val}(a_u) + \langle u, v \rangle \geq 0$ for all $v \in P$ and all u . The scheme $Y^P := \text{Spec}(R[Y^P])$ is called a *polyhedral integral model* of Y_σ .

In other words, $R[Y^P]$ is the intersection of $K\langle\mathcal{U}^P\rangle^\circ$ with $K[M]$. It is clear that $K\langle\mathcal{U}^P\rangle^\circ$ is the ϖ -adic completion of $R[Y^P]$. Note that $R[Y^P]$ is torsion-free and hence flat over R .

LEMMA 4.9. *The tensor product $R[Y^P] \otimes_R K$ is equal to $K[Y_\sigma]$.*

Proof. By definition we have $R[Y^P] \otimes_R K \subset K[Y_\sigma]$. For the other inclusion, note that if $g = \sum b_u x^u$ is in $K[Y_\sigma]$, then the minimum over v in P of $\text{val}(b_u) + \langle u, v \rangle$ is achieved at some vertex of P . It follows that some sufficiently high power of ϖ times g is in $R[Y^P]$, and hence g is in $R[Y^P] \otimes_R K$. \square

Remark 4.10. One could equivalently define $R[Y^P]$ to be the subring of $K[M]$ satisfying the same inequalities. Since P is closed under addition of points in σ , any Laurent polynomial satisfying these inequalities for all v in P must be supported in σ^\vee .

We will use the following notation in the proof of Proposition 4.11. For each face $F \leq P$, let $\sigma(F)$ be the cone in $N_{\mathbb{R}}$ spanned by $P - v$ for any v in the relative interior of F . In other words, $\sigma(F) = \text{Star}_P(F)$. We fix a labeling v_1, \dots, v_r for the vertices of P , and write σ_i for $\sigma(v_i)$. The dual cone σ_i^\vee is

$$\sigma_i^\vee = \{u \in \sigma^\vee : \langle u, v_i \rangle \leq \langle u, v_j \rangle \text{ for all } j\}.$$

The cones $\sigma_1^\vee, \dots, \sigma_r^\vee$ are the maximal cones of the (possibly degenerate) inner normal fan of P , and their union is σ^\vee .

PROPOSITION 4.11. *Let P be a G -rational polyhedron in $N_{\mathbb{R}}$. Then $R[Y^P]$ is finitely presented over R .*

Proof. By [RG71, Corollary 3.4.7], any finitely-generated and flat algebra over an integral domain is automatically of finite presentation, so it suffices to show that $R[Y^P]$ is finitely generated.

The cones $\sigma_1^\vee, \dots, \sigma_r^\vee$ cover σ^\vee , so $R[Y^P]$ is generated by the subrings

$$A_j = R[Y^P] \cap K[\sigma_j^\vee \cap M]$$

for $1 \leq j \leq r$. Therefore, it will suffice to show that each A_j is finitely generated over R .

The semigroup $\sigma_j^\vee \cap M$ is finitely generated by Gordan's lemma [Ful93, § 1.2, Proposition 1]. Let u_1, \dots, u_s be generators, and choose a_1, \dots, a_s in K^\times such that $\text{val}(a_i) + \langle u_i, v_j \rangle = 0$. Then each monomial in A_j can be written as an element of R times a monomial in the $a_i x^{u_i}$. It follows that A_j is finitely generated over R , as required, with generating set $\{a_1 x^{u_1}, \dots, a_s x^{u_s}\}$. \square

In particular, Y^P is a flat and finitely-presented R -model of the affine toric variety Y_σ .

Remark 4.12. As in Remark 4.6, one can construct an algebraic model Y^P of Y_σ associated with an integral affine but not G -affine polyhedron P ; when P is a point this is done in [OP13]. This model is not of finite type.

4.13 Polyhedral integral and formal models of subschemes

Let P be an integral G -affine polyhedron with pointed recession cone σ . Let X be the closed subscheme of the affine toric variety Y_σ over K defined by an ideal $\mathfrak{a} \subset K[Y_\sigma]$.

DEFINITION 4.14. (i) Let $\mathcal{X}^P = X^{\text{an}} \cap \mathcal{U}^P$. This is the Zariski-closed subspace of \mathcal{U}^P defined by $\mathfrak{a}K\langle \mathcal{U}^P \rangle$.

(ii) The *polyhedral integral model* of X is the scheme-theoretic closure X^P of X in Y^P . It is defined by the ideal $\mathfrak{a}^P = \mathfrak{a} \cap R[Y^P]$.

(iii) The *polyhedral formal model* of \mathcal{X}^P is the ϖ -adic completion \mathfrak{X}^P of X^P . We will show in Proposition 4.17 that \mathfrak{X}^P is an admissible formal scheme with generic fiber \mathcal{X}^P .

(iv) The *canonical model* of \mathcal{X}^P is

$$\mathfrak{X}_{\text{can}}^P = \text{Spf} \left((K\langle \mathcal{U}^P \rangle / \mathfrak{a}K\langle \mathcal{U}^P \rangle)^\circ \right).$$

By Theorem 3.12, the canonical model is admissible if and only if \mathcal{X}^P is reduced.

NOTATION 4.15. The *P -initial degeneration* of X is defined to be

$$\text{in}_P(X) = X^P \otimes_R k = \mathfrak{X}^P \otimes_R k.$$

As usual we write $\overline{\mathfrak{X}}_{\text{can}}^P = \mathfrak{X}_{\text{can}}^P \otimes_R k$. This coincides with the canonical reduction of \mathcal{X}^P when \mathcal{X}^P is reduced. In the case where P is a single point $w \in N_G$ we write \mathcal{X}^w , \mathfrak{X}^w , $\text{in}_w(X)$, etc. In this case, $\text{in}_w(X)$ is the w -initial degeneration of X in the sense generally used in the literature (and in the introduction).

LEMMA 4.16. *The ideal \mathfrak{a}^P is finitely generated.*

Proof. Since X^P is the closure of its generic fiber, it is flat over $\text{Spec } R$, and its coordinate ring is a quotient of the finitely-generated R -algebra $R[M]$. Since any finitely-generated flat algebra over an integral domain is finitely presented [RG71, Corollary 3.4.7], it follows that \mathfrak{a}^P is finitely generated. \square

PROPOSITION 4.17. *The formal scheme \mathfrak{X}^P is the formal closed subscheme of \mathfrak{U}^P defined by $\mathfrak{a}^P K\langle \mathcal{U}^P \rangle^\circ$. It is an admissible formal scheme with generic fiber \mathcal{X}^P and special fiber $\text{in}_P(X)$.*

Proof. The admissibility of \mathfrak{X}^P is a consequence of Proposition 3.7(i). If $A = R[Y^P]/\mathfrak{a}$, then by definition $X^P = \text{Spec}(A)$ and $\mathfrak{X}^P = \text{Spf}(\widehat{A})$, where \widehat{A} is the ϖ -adic completion of A . By Proposition 3.7(ii) the sequence

$$0 \longrightarrow \mathfrak{a}^P K\langle \mathcal{U}^P \rangle^\circ \longrightarrow K\langle \mathcal{U}^P \rangle^\circ \longrightarrow \widehat{A} \longrightarrow 0$$

is exact; it follows that \mathfrak{X}^P is the closed subscheme of \mathcal{U}^P defined by $\mathfrak{a}^P K \langle \mathcal{U}^P \rangle^\circ$. We have

$$(K \langle \mathcal{U}^P \rangle^\circ / \mathfrak{a}^P K \langle \mathcal{U}^P \rangle^\circ) \otimes_R K = K \langle \mathcal{U}^P \rangle / \mathfrak{a} K \langle \mathcal{U}^P \rangle$$

since $K\mathfrak{a}^P = \mathfrak{a}$, so $\mathfrak{X}^P \otimes_R K = \mathcal{X}^P$. The special fiber of \mathfrak{X}^P agrees with the special fiber of X^P by construction. \square

The canonical inclusion

$$K \langle \mathcal{U}^P \rangle^\circ / \mathfrak{a}^P K \langle \mathcal{U}^P \rangle^\circ \hookrightarrow (K \langle \mathcal{U}^P \rangle / \mathfrak{a} K \langle \mathcal{U}^P \rangle)^\circ$$

induces a map of formal schemes

$$\mathfrak{X}_{\text{can}}^P \longrightarrow \mathfrak{X}^P.$$

As this morphism induces an isomorphism on analytic generic fibers, it is *finite* when \mathcal{X}^P is reduced by Proposition 3.8(ii). Its special fiber is a morphism $\overline{\mathfrak{X}}_{\text{can}}^P \rightarrow \text{in}_P(X)$. Many of the results of this paper are proved by using this morphism and the results of Section 3 (in particular the projection formula, Proposition 3.25) to compare these two models.

4.18 Compatibility with extension of the ground field

We continue to use the notation of Section 4.13. Let K' be an algebraically closed complete valued field extension of K , with valuation ring R' and residue field k' . Let P be an integral G -affine polyhedron in $N_{\mathbb{R}}$ with pointed recession cone σ . Let $Y'_\sigma = Y_\sigma \otimes_K K'$, so Y'_σ is the affine toric variety defined over K' with dense torus $\mathbf{T}' := \mathbf{T} \otimes_K K'$ associated with the cone σ . The triangle

$$\begin{array}{ccc} (Y'_\sigma)^{\text{an}} & \xrightarrow{\quad} & Y_\sigma^{\text{an}} \\ & \searrow \text{trop} & \swarrow \text{trop} \\ & N_{\mathbb{R}}(\sigma) & \end{array}$$

commutes, so $\mathcal{U}^P \widehat{\otimes}_K K'$ is the polyhedral domain in $(Y'_\sigma)^{\text{an}}$ associated with P . Likewise the polyhedral integral model $(Y')^P$ of Y'_σ associated with P is naturally identified with $Y^P \otimes_R R'$. Indeed, as an R -module we have

$$R[Y^P] = \bigoplus_{u \in \sigma^\vee \cap M} R_u \cdot x^u \subset K[\sigma^\vee \cap M], \quad \text{where } R_u = \{a \in R : \text{val}(a) \geq \max_{v \in \text{vert}(P)} -\langle u, v \rangle\}.$$

Since $\langle u, v \rangle \in G$ for all $u \in M$ and $v \in \text{vert}(P)$, each R_u is a free R -module of rank 1, so the image of $R_u \otimes_R R'$ in K' is exactly R'_u .

Let $X \subset Y_\sigma$ be the closed subscheme defined by an ideal $\mathfrak{a} \subset K[\sigma^\vee \cap M]$, and let $X' = X \otimes_K K' \subset Y'_\sigma$, so X' is defined by $\mathfrak{a}K'[\sigma^\vee \cap M]$. Since the above triangle is commutative, we have $\text{Trop}(X) = \text{Trop}(X') \subset N_{\mathbb{R}}(\sigma)$, and $\text{trop}: (X')^{\text{an}} \rightarrow \text{Trop}(X)$ factors through the natural map $(X')^{\text{an}} \rightarrow X^{\text{an}}$. Hence

$$(\mathcal{X}')^P = \text{trop}^{-1}(P) \cap (X')^{\text{an}} = \mathcal{X}^P \widehat{\otimes}_K K'.$$

Since schematic closure commutes with flat base change, the polyhedral integral model $(X')^P$ of X' coincides with $X^P \otimes_R R'$; hence if $\mathfrak{a}^P = \mathfrak{a} \cap R[Y^P]$ is the ideal defining X^P , then $(X')^P$ is defined by $\mathfrak{a}^P R'[(Y')^P]$. It follows from this and Proposition 4.17 that $(\mathfrak{X}')^P = \mathfrak{X}^P \widehat{\otimes}_R R'$, and in particular that $\text{in}_P(X') = \text{in}_P(X) \otimes_k k'$. As for the canonical models, suppose that X is reduced, so X' is reduced as well. Then $(\mathfrak{X}')^P_{\text{can}} = \mathfrak{X}^P_{\text{can}} \widehat{\otimes}_R R'$ because $(\mathfrak{X}^P_{\text{can}} \widehat{\otimes}_R R') \otimes_{R'} k' = (\mathfrak{X}^P_{\text{can}} \otimes_R k) \otimes_k k'$ is reduced; cf. Proposition 3.13.

Below we will make various definitions by passing to a valued field extension K' of K . In order for these definitions to be independent of the choice of K' , we will need the following fact, proven in [Duc09, § 0.3.2] or [Con08, § 4].

LEMMA 4.19. *Let K_1 and K_2 be complete valued field extensions of K . Then there is a complete valued field extension K' of K admitting isometric embeddings $K_1 \hookrightarrow K'$ and $K_2 \hookrightarrow K'$ over K .*

4.20 Relative multiplicities and tropical multiplicities

Recall Section 3.3 that if $\mathcal{X} = \mathcal{M}(\mathcal{A})$ is an affinoid space, then the reduction map induces a one-to-one correspondence between the Shilov boundary points of \mathcal{X}^{an} and the generic points of the canonical reduction $\text{Spec}(\tilde{\mathcal{A}})$. This leads to the following definition.

DEFINITION 4.21. Let $X \subset \mathbf{T}$ be a reduced and equidimensional closed subscheme, let $x \in X^{\text{an}}$, let $w = \text{trop}(x)$, and suppose $w \in N_G$. Define the *relative multiplicity* $m_{\text{rel}}(x)$ of x in $\text{trop}^{-1}(w)$ as follows. If x is not a Shilov boundary point of $\text{trop}^{-1}(w)$, then we define its multiplicity to be 0. Otherwise $\text{red}(x)$ is the generic point of an irreducible component $\bar{\mathcal{C}}$ of $\bar{\mathfrak{X}}_{\text{can}}^w$; we define the multiplicity of x to be $[\bar{\mathcal{C}} : \text{im}(\bar{\mathcal{C}})]$, where $\text{im}(\bar{\mathcal{C}})$ is the image of $\bar{\mathcal{C}}$ in $\text{in}_w(X)$ (this is an irreducible component by Corollary 3.11).

Now suppose $w \notin N_G$. Let K' be an algebraically closed complete valued field extension of K such that $w \in N_{G'}$, where G' is the value group of K' . Let $X' = X \otimes_K K'$, and let $\phi: (X')^{\text{an}} \rightarrow X^{\text{an}}$ be the natural morphism. We define

$$m_{\text{rel}}(x) = \sum_{x' \in \phi^{-1}(x)} m_{\text{rel}}(x').$$

In order for this definition to make sense, by Lemma 4.19 we only have to show that if $K \subset K' \subset K''$ are algebraically closed complete valued field extensions, then we can calculate $m_{\text{rel}}(x)$ with respect to either K' or K'' . Replacing K with K' , we are reduced to showing the following.

LEMMA 4.22. *Let K' be an algebraically closed complete valued field extension of K , and let $X' = X \otimes_K K'$. Let $x \in X^{\text{an}}$, let $w = \text{trop}(x)$, and suppose $w \in N_G$. Then the natural map $(\mathcal{X}')^w \rightarrow \mathcal{X}^w$ induces a bijection of Shilov boundary points which preserves relative multiplicities.*

Proof. Let k' be the residue field of K' . As discussed in Section 4.18, we have $(\bar{\mathfrak{X}}')^w_{\text{can}} = \bar{\mathfrak{X}}_{\text{can}}^w \otimes_k k'$ and $\text{in}_w(X') = \text{in}_w(X) \otimes_k k'$, so the first assertion follows from the fact that $\bar{\mathfrak{X}}_{\text{can}}^w \otimes_k k' \rightarrow \bar{\mathfrak{X}}_{\text{can}}^w$ induces a bijection on irreducible components. Let $\bar{\mathcal{C}}$ be an irreducible component of $\bar{\mathfrak{X}}_{\text{can}}^w$, and let $\bar{\mathcal{D}}$ be its image in $\bar{\mathfrak{X}}^w$. Then $[\bar{\mathcal{C}} : \bar{\mathcal{D}}] = [\bar{\mathcal{C}} \otimes_k k' : \bar{\mathcal{D}} \otimes_k k']$, so relative multiplicities are preserved as well. \square

Later we will relate $m_{\text{rel}}(x)$ to other geometrically-defined notions of multiplicity; see Proposition 4.31 and Theorem 5.8. For the moment we relate relative multiplicities to tropical multiplicities, defined as follows.

DEFINITION 4.23. Let $X \subset \mathbf{T}$ be a closed subscheme, and let $w \in \text{Trop}(X)$. If $w \in N_G$, then the *tropical multiplicity* of X at w is defined to be

$$m_{\text{Trop}}(w) = \sum_{\bar{C} \subset \text{in}_w(X)} \text{mult}_{\text{in}_w(X)}(\bar{C}),$$

where the sum is taken over all irreducible components \bar{C} of $\text{in}_w(X)$. If $w \notin N_G$, then let K' be an algebraically closed complete valued field extension of K such that $w \in N_{G'}$, where G' is the

value group of K' . Let $X' = X \otimes_K K'$. We define $m_{\text{Trop}}(w)$ to be the tropical multiplicity of w relative to $\text{trop}: (X')^{\text{an}} \rightarrow \text{Trop}(X)$.

The fact that $m_{\text{Trop}}(w)$ is independent of the choice of K' is proved in [OP13, Remark A.5]. It is also one of the consequences of the following proposition.

PROPOSITION 4.24. *Let $X \subset \mathbf{T}$ be a reduced and equidimensional closed subscheme, and let $w \in \text{Trop}(X)$. Then*

$$m_{\text{Trop}}(w) = \sum_{x \in \text{trop}^{-1}(w)} m_{\text{rel}}(x).$$

Proof. We immediately reduce to the case where $w \in N_G$ by extending the ground field if necessary. By definition we have

$$\sum_{x \in \text{trop}^{-1}(w)} m_{\text{rel}}(x) = \sum_{\bar{\mathcal{C}} \subset \bar{\mathfrak{X}}_{\text{can}}^w} [\bar{\mathcal{C}} : \text{im}(\bar{\mathcal{C}})],$$

where the sum is taken over all irreducible components $\bar{\mathcal{C}}$ of $\bar{\mathfrak{X}}_{\text{can}}^w$; the image $\text{im}(\bar{\mathcal{C}})$ of $\bar{\mathcal{C}}$ in $\text{in}_w(X)$ is an irreducible component by Corollary 3.11. Also by definition,

$$m_{\text{Trop}}(w) = \sum_{\bar{C} \subset \text{in}_w(X)} \text{mult}_{\text{in}_w(X)}(\bar{C}),$$

where the sum is taken over all irreducible components \bar{C} of $\text{in}_w(X)$. By Proposition 3.25, for every irreducible component \bar{C} of $\text{in}_w(X)$ we have

$$\text{mult}_{\text{in}_w(X)}(\bar{C}) = \sum_{\bar{\mathcal{C}} \rightarrow \bar{C}} [\bar{\mathcal{C}} : \bar{C}],$$

where the sum is taken over all irreducible components $\bar{\mathcal{C}}$ of $\bar{\mathfrak{X}}_{\text{can}}^w$ mapping onto \bar{C} (for any such $\bar{\mathcal{C}}$ we have $\text{mult}_{\bar{\mathfrak{X}}_{\text{can}}^w}(\bar{\mathcal{C}}) = 1$ since $\bar{\mathfrak{X}}_{\text{can}}^w$ is reduced). Therefore

$$m_{\text{Trop}}(w) = \sum_{\bar{C} \subset \text{in}_w(X)} \text{mult}_{\text{in}_w(X)}(\bar{C}) = \sum_{\bar{C} \subset \text{in}_w(X)} \sum_{\bar{\mathcal{C}} \rightarrow \bar{C}} [\bar{\mathcal{C}} : \bar{C}] = \sum_{\bar{\mathcal{C}} \subset \bar{\mathfrak{X}}_{\text{can}}^w} [\bar{\mathcal{C}} : \text{im}(\bar{\mathcal{C}})]. \quad \square$$

EXAMPLE 4.25. In Example 2.6, we have $m_{\text{rel}}(\xi) = 1$ for all $\xi \in \Gamma$ and $m_{\text{rel}}(\xi) = 0$ for all $\xi \notin \Gamma$. This follows from Proposition 4.24 and the concrete description of the tropicalization map in Example 2.6, together with the observation that $m_{\text{Trop}}(0, 0) = m_{\text{Trop}}(2, 2) = 1$ (since the initial degenerations $\text{in}_{(0,0)}(E) \cong \text{Spec } k[x, y, x^{-1}, y^{-1}]/(y^2 - x^3 - x^2)$ and $\text{in}_{(2,2)}(E) \cong \text{Spec } k[x, y, x^{-1}, y^{-1}]/(y^2 - x^2 - 1)$ are both integral schemes over k). Note that $m_{\text{rel}}(\xi) > 0$ for all $\xi \in \Gamma$ because ξ is contained in the topological boundary of $\text{trop}^{-1}(\text{trop}(\xi))$ in E^{an} , hence in the Shilov boundary; see Section 3.4.

4.26 Polyhedral structures on tropicalizations

Let W be a G -rational affine space in $N_{\mathbb{R}}$, and let W_0 be the linear space under W , so W_0 is spanned by $W_0 \cap N$. Set $N' = N/(W_0 \cap N)$ and $M' = W_0^{\perp} \cap M \subset M$, and let \mathbf{T}' be the torus $\text{Spec}(K[M'])$. We call \mathbf{T}' the *torus transverse to W* . Let $w' \in N'_G$ be the image of any point of W . Then

$$R[(\mathbf{T}')^{w'}] = \left\{ \sum_{u \in M'} a_u x^u \in K[M'] : \text{val}(a_u) + \langle u, w' \rangle \geq 0 \right\},$$

so for all $w \in N_G \cap W$ we have $R[(\mathbf{T}')^{w'}] \subset R[\mathbf{T}^w]$. Hence we have a natural morphism $\pi_w: \bar{\mathbf{T}}^w \rightarrow (\bar{\mathbf{T}}')^{w'}$ for all $w \in N_G \cap W$.

Remark 4.27. Let $N'' = \ker(N \rightarrow N') = W_0 \cap N$, and let $M'' = \text{Hom}_{\mathbb{Z}}(N'', \mathbb{Z})$, so we have exact sequences

$$0 \longrightarrow N'' \longrightarrow N \longrightarrow N' \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow M'' \longrightarrow M \longrightarrow M' \longrightarrow 0$$

inducing an exact sequence of tori

$$0 \longrightarrow \mathbf{T}'' \longrightarrow \mathbf{T} \longrightarrow \mathbf{T}' \longrightarrow 0,$$

where $\mathbf{T}'' = \text{Spec}(K[M''])$. We call \mathbf{T}'' the *torus parallel to W* . Choosing a splitting of $N \rightarrow N'$ splits all three exact sequences, and in particular furnishes an isomorphism $\mathbf{T} \cong \mathbf{T}' \times \mathbf{T}''$. Let $w \in W \cap N_G$, and let w'' be its image in $N''_{\mathbb{R}}$. Then we have an isomorphism $\overline{\mathbf{T}}^w \cong (\overline{\mathbf{T}}')^{w'} \times (\overline{\mathbf{T}}'')^{w''}$ under which π_w corresponds to the projection onto the first factor.

THEOREM 4.28. *Let $X \subset \mathbf{T}$ be an equidimensional subscheme of dimension d . The set $\text{Trop}(X)$ admits a polyhedral complex structure of pure dimension d with the following properties:*

- (i) *The tropical multiplicities are constant along the relative interior of every maximal face.*
- (ii) *Let w be contained in the relative interior of a maximal face τ of $\text{Trop}(X)$, let $W = \text{span}(\tau)$, let \mathbf{T}' be the torus transverse to W , and let $\pi_w: \overline{\mathbf{T}}^w \rightarrow (\overline{\mathbf{T}}')^{w'}$ be the natural map. Then $\text{in}_w(X) \cong \pi_w^{-1}(\overline{Y})$ for some dimension 0 subscheme \overline{Y} of $(\overline{\mathbf{T}}')^{w'}$.*

Proof. The first part is a basic result in tropical geometry; it is proved in [MS15, §3.3].⁴ Let $\mathbf{T}'' \subset \mathbf{T}$ be the torus parallel to W and write $\overline{\mathbf{T}}'' = (\overline{\mathbf{T}}'')^0$, so $(\overline{\mathbf{T}}')^{w'}$ is the quotient of $\overline{\mathbf{T}}^w$ by $\overline{\mathbf{T}}''$. By [Spe05, Proposition 2.2.4], the initial degeneration $\text{in}_w(X)$ is invariant under the action of $\overline{\mathbf{T}}''$. Therefore $\text{in}_w(X)$ is the inverse image of a closed subscheme \overline{Y} of $(\overline{\mathbf{T}}')^{w'}$ (in fact \overline{Y} is the quotient $\text{in}_w(X)/\overline{\mathbf{T}}'' \subset (\overline{\mathbf{T}}')^{w'}$); counting dimensions, we see that $\dim(\overline{Y}) = 0$. See [MFK94, Theorem 1.1.1 and Amplification 1.1.3] for basic existence results about geometric quotients of affine schemes over a field by a free action of a reductive group. \square

In the situation of Theorem 4.28(ii), let \mathbf{T}'' be the torus parallel to W , and choose a splitting $\mathbf{T} \cong \mathbf{T}' \times \mathbf{T}''$ as in Remark 4.27. Then $\text{in}_w(X) \cong \overline{Y} \times (\overline{\mathbf{T}}'')^{w''}$.

4.29 The tropical projection formula

Let $X \subset \mathbf{T}$ be a reduced and equidimensional closed subscheme of dimension d , and let P be an integral G -affine polytope contained in the relative interior of a maximal (d -dimensional) face τ of a polyhedral complex decomposition of $\text{Trop}(X)$ as in Theorem 4.28. Let W be the affine span of τ , let \mathbf{T}' be the torus transverse to W (Section 4.26), let \mathbf{T}'' be the torus parallel to W (Remark 4.27), and choose a splitting $\mathbf{T} \rightarrow \mathbf{T}''$. Note that $\dim(\mathbf{T}'') = d$. Let P'' be the image of P in $N''_{\mathbb{R}}$, so $\mathcal{U}^{P''}$ is a polytopal domain in $(\mathbf{T}'')^{\text{an}}$. The map $\mathcal{U}^P \rightarrow \mathcal{U}^{P''}$ induces a morphism $\psi^P: \mathcal{X}^P \rightarrow \mathcal{U}^{P''}$.

THEOREM 4.30. *The morphism $\psi^P: \mathcal{X}^P \rightarrow \mathcal{U}^{P''}$ is finite, and every irreducible component of \mathcal{X}^P surjects onto $\mathcal{U}^{P''}$.*

Proof. For dimension reasons it suffices to show that ψ^P is finite. Since $\mathcal{X}^P = \mathcal{M}(\mathcal{A})$ and $\mathcal{U}^P = \mathcal{M}(K\langle \mathcal{U}^{P''} \rangle)$ are both affinoid, by the rigid-analytic direct image theorem [BGR84,

⁴The proofs in [MS15, §3.3] assume that there is a section to the valuation map $\text{val}: K^\times \rightarrow G$. Such a section always exists when K is an algebraically closed nonarchimedean field; the following short proof was communicated to us by Speyer. If $G = \{0\}$, then there is nothing to prove. Otherwise, consider the short exact sequence $0 \rightarrow U \rightarrow K^\times \rightarrow G \rightarrow 0$. Since K is algebraically closed, the group U is divisible. Thus U is injective as a \mathbb{Z} -module, so $\text{Ext}^1(A, U) = 0$ and the valuation map splits.

Theorem 9.6.3/1] it suffices to show that $\mathcal{X}^P \rightarrow \mathcal{U}^{P''}$ is proper in the sense of [BGR84, §9.6.2]. In fact we will show that $\mathcal{X}^P \subseteq_{\mathcal{U}^{P''}} \mathcal{X}^P$, that is, that there exist affinoid generators f_1, \dots, f_r for \mathcal{A} over $K\langle \mathcal{U}^{P''} \rangle$ such that $|f_1|_{\text{sup}}, \dots, |f_r|_{\text{sup}} < 1$.

Choosing bases for N' and N'' , we obtain isomorphisms $N'_{\mathbb{R}} \cong \mathbb{R}^{n'}$, $N''_{\mathbb{R}} \cong \mathbb{R}^d$, and $N_{\mathbb{R}} \cong \mathbb{R}^{n'} \times \mathbb{R}^d$. Translating by an element of $\mathbf{T}(K)$, we may and do assume that $P \subset \{0\} \times N''_{\mathbb{R}}$ (so $P = P''$). For $\epsilon \in G$ with $\epsilon > 0$ we let $I_{\epsilon} \subset N'_{\mathbb{R}}$ be the cube $[-\epsilon, \epsilon]^{n'}$, so I_{ϵ} is an integral G -affine polytope in $N'_{\mathbb{R}}$, and $P_{\epsilon} := I_{\epsilon} \times P''$ is an integral G -affine polytope in $N_{\mathbb{R}} = N'_{\mathbb{R}} \times N''_{\mathbb{R}}$ containing P . Since τ is a maximal face, we have $P_{\epsilon} \cap \text{Trop}(X) = P$ for small ϵ ; we fix such an ϵ as well as an element $e \in K$ with $\text{val}(e) = \epsilon$. The polytopal subdomain $\mathcal{U}^{I_{\epsilon}} \subset (\mathbf{T}')^{\text{an}}$ is a product of annuli of inner radius $|e|$ and outer radius $|e|^{-1}$, so if $u_1, \dots, u_{n'}$ is a basis for M' , then $\{ex^{\pm u_1}, \dots, ex^{\pm u_{n'}}\}$ is a set of affinoid generators for $K\langle \mathcal{U}^{I_{\epsilon}} \rangle$. Since $\mathcal{U}^{P_{\epsilon}} = \mathcal{U}^{I_{\epsilon}} \times_K \mathcal{U}^{P''}$, it follows that $\{ex^{\pm u_1}, \dots, ex^{\pm u_{n'}}\}$ is a set of affinoid generators for $K\langle \mathcal{U}^{P_{\epsilon}} \rangle$ over $K\langle \mathcal{U}^{P''} \rangle$. Since $P_{\epsilon} \cap \text{Trop}(X) = P$, we have $X^{\text{an}} \cap \mathcal{U}^{P_{\epsilon}} = \mathcal{X}^P$, so $\{ex^{\pm u_1}, \dots, ex^{\pm u_{n'}}\}$ can be regarded as a set of affinoid generators for \mathcal{A} over $K\langle \mathcal{U}^{P''} \rangle$. But by construction $|x^{u_i}(x)| = 1$ for all $x \in \mathcal{X}^P$ and all $i = 1, \dots, n'$, so $|ex^{u_i}(x)| = |e| < 1$. This proves that ψ^P is finite. \square

It follows from Theorem 4.30 and Remarks 3.18(i) and 3.22(i) that ψ^P has a (pure) degree.

PROPOSITION 4.31. *In the situation of Section 4.29, let $\mathcal{Y} \subset \mathcal{X}^P$ be a union of connected components, and let $w \in P$. Then*

$$[\mathcal{Y} : \mathcal{U}^{P''}] = \sum_{x \in \mathcal{Y} \cap \text{trop}^{-1}(w)} m_{\text{rel}}(x).$$

Proof. Extending the ground field if necessary, we assume $w \in N_G$. Let w'' be the image of w in N''_G . Since $\mathcal{Y} \cap \mathcal{X}^w \rightarrow \mathcal{U}^{w''}$ is obtained by flat base change from $\mathcal{Y} \rightarrow \mathcal{U}^{P''}$, we may replace P by w and P'' by w'' to assume $\mathcal{Y} \subset \mathcal{X}^w$ (cf. Proposition 3.23(i)). Let \mathfrak{Y} be the canonical model of Y . The canonical reduction $\overline{\mathfrak{Y}}$ of \mathcal{Y} is a union of connected components of $\overline{\mathfrak{X}}_{\text{can}}^w$, so for $x \in \mathcal{Y}$ the relative multiplicity $m_{\text{rel}}(x)$ is nonzero if and only if $\text{red}(x)$ is the generic point of an irreducible component $\overline{\mathfrak{C}}$ of $\overline{\mathfrak{Y}}$, in which case $m_{\text{rel}}(x) = [\overline{\mathfrak{C}} : \text{im}(\overline{\mathfrak{C}})]$, where $\text{im}(\overline{\mathfrak{C}})$ is the image of $\overline{\mathfrak{C}}$ in $\text{in}_w(X)$. Noting that $\overline{\mathfrak{U}}^{w''}$ is an integral domain and $\overline{\mathfrak{Y}}$ is reduced, applying Proposition 3.25 to $\mathfrak{Y} \rightarrow \overline{\mathfrak{U}}^{w''}$ yields

$$[\mathcal{Y} : \mathcal{U}^{w''}] = \sum_{\mathfrak{C} \subset \overline{\mathfrak{Y}}} [\mathfrak{C} : \overline{\mathfrak{U}}^{w''}].$$

Since $\overline{\mathfrak{U}}^{w''} = (\overline{\mathbf{T}}'')^{w''}$ and $\text{in}_w(X) \cong \overline{D} \times (\overline{\mathbf{T}}'')^{w''}$ for some dimension 0 scheme $\overline{D} \subset (\overline{\mathbf{T}}'')^{w''}$ (cf. Remark 4.27), the reduced space underlying any irreducible component of $\text{in}_w(X)$ is isomorphic to $(\overline{\mathbf{T}}'')^{w''}$. Therefore $[\text{im}(\overline{\mathfrak{C}}) : \overline{\mathfrak{U}}^{w''}] = 1$ for any irreducible component $\overline{\mathfrak{C}} \subset \overline{\mathfrak{Y}}$, so $[\mathfrak{C} : \overline{\mathfrak{U}}^{w''}] = [\overline{\mathfrak{C}} : \text{im}(\overline{\mathfrak{C}})]$ and the proposition follows. \square

COROLLARY 4.32 (Tropical projection formula). *In the situation of Theorem 4.30, the degree of $\psi^P : \mathcal{X}^P \rightarrow \mathcal{U}^{P''}$ is equal to $m_{\text{Trop}}(w)$ for any $w \in P$.*

Proof. Assuming $P = w$ and $P'' = w''$ as in the proof of Proposition 4.31, the result follows immediately from Propositions 4.31 and 4.24. \square

Remark 4.33. The tropical projection formula is an equality of the degree of the morphism $\mathcal{X}^P \rightarrow \mathcal{U}^{P''}$ (a morphism on the generic fiber) with the degree of a morphism $\overline{\mathfrak{X}}^w \rightarrow \overline{\mathfrak{U}}^{w''}$ (a morphism on the special fiber). It is conceptually very close to the projection formula as stated in Proposition 3.25, as indeed that is the main tool used in its proof; it is for this reason that we call it the tropical projection formula.

5. The tropicalization of a nonarchimedean analytic curve

In this section we freely use the definitions and notation from [BPR13]. In particular, for $a, b \in K^\times$ we have $\mathbf{S}(a, b) = \{t \in \mathbf{G}_m^{\text{an}} : |a| \leq |t| \leq |b|\}$, a closed annulus; we also set $\mathbf{S}(a) = \mathbf{S}(a, 1)$ and $\mathbf{S}(1) = \{t \in \mathbf{G}_m^{\text{an}} : |t| = 1\}$, the closed annulus of modulus 0.

5.1 The setup

Throughout this section X denotes a smooth connected algebraic curve realized as a closed subscheme of a torus \mathbf{T} , \widehat{X} is the smooth completion of X , and $D = \widehat{X}(K) \setminus X(K)$ is the set of punctures. We will denote a choice of semistable vertex set for X by V , and we let $\Sigma = \Sigma(X, V)$ be the associated skeleton. See [BPR13, § 3].

If we choose a basis for M , then we obtain isomorphisms $N_{\mathbb{R}} \cong \mathbb{R}^n$ and $K[\mathbf{T}] \cong K[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$; if $f_i \in K[X]^\times$ is the image of x_i , then

$$\text{trop}(\|\cdot\|) = (-\log \|f_1\|, \dots, -\log \|f_n\|). \quad (5.1.1)$$

5.2 Compatible polyhedral structures

The tropicalization of X is a polyhedral complex of pure dimension 1 in $N_{\mathbb{R}}$. We can regard $\text{Trop}(X)$ as a dimension 1 abstract G -rational polyhedral complex where the metric on the edges is given by the lattice length, that is, the length in the direction of a primitive lattice vector in N .

Recall the following consequence of the slope formula [BPR13, Theorem 5.15].

LEMMA 5.3. *Let e be an edge of Σ , and let $f \in K[X]^\times$. The map $\|\cdot\| \mapsto -\log \|f\| : X^{\text{an}} \rightarrow \mathbb{R}$ restricts to a G -affine linear function from e to \mathbb{R} with integer slope.*

Since the G -rational points of an edge e of Σ are exactly the type 2 points of X^{an} contained in e (cf. [BPR13, § 3.12]), it follows from (5.1.1) and the G -rationality of $-\log \|f\|$ as above that trop maps type 2 points into N_G .

PROPOSITION 5.4. (i) *The tropicalization map $\text{trop} : X^{\text{an}} \rightarrow \text{Trop}(X)$ factors through the retraction $\tau_\Sigma : X^{\text{an}} \rightarrow \Sigma$.*

(ii) *We can choose V and a polyhedral complex structure on $\text{Trop}(X)$ as in Theorem 4.28 such that $\text{trop} : \Sigma \rightarrow \text{Trop}(X)$ is a morphism of dimension 1 abstract G -rational polyhedral complexes.*

Proof. The first part follows from (5.1.1) and the slope formula [BPR13, Theorem 5.15]. Let e be an edge of Σ . It follows from Lemma 5.3 as applied to f_1, \dots, f_n that trop restricts to an expansion by an integer multiple with respect to the intrinsic metric on e and the lattice length on its image. Hence there exist refinements of the polyhedral structures on Σ and on $\text{Trop}(X)$ such that $\text{trop} : \Sigma \rightarrow \text{Trop}(X)$ becomes a morphism of dimension 1 abstract G -rational polyhedral complexes. By [BPR13, Proposition 3.13(2)], any refinement of Σ is also a skeleton of X . \square

From now on we assume that our skeleton Σ of X and our choice of polyhedral structure on $\text{Trop}(X)$ are compatible in the sense of Proposition 5.4(ii). If $e \subset \Sigma$ is an interval contained in an edge, we let $\ell_{\text{an}}(e)$ be its length with respect to the skeletal metric, and if $e' \subset \text{Trop}(X)$ is an interval contained in an edge, we let $\ell_{\text{Trop}}(e')$ be its lattice length. As a consequence of Proposition 5.4(ii), if $\text{trop}(e) = e'$, then $\ell_{\text{Trop}}(e')$ is an integer multiple of $\ell_{\text{an}}(e)$.

DEFINITION 5.5. Let $e \subset \Sigma$ be an edge, and let $e' \subset \text{Trop}(X)$ be its image. We define the *expansion factor of e* to be the unique integer $m_{\text{rel}}(e) \in \mathbb{Z}_{\geq 0}$ such that

$$\ell_{\text{Trop}}(\text{trop}(\tilde{e})) = m_{\text{rel}}(e) \cdot \ell_{\text{an}}(\tilde{e})$$

for any finite-length segment \tilde{e} contained in e .

Remark 5.6. For $u \in M$ let $f_u \in K[X]^\times$ be the image of the character $x^u \in K[M]$. Let e be an edge of Σ , and let $s_u \in \mathbb{Z}_{\geq 0}$ be the absolute value of the slope of $-\log \|f_u\|$ on e . It follows easily from the definitions that $m_{\text{rel}}(e) = \gcd\{s_u : u \in M\}$. More concretely, let u_1, \dots, u_n be a basis for M , and let $f_i \in K[X]^\times$ be the image of x^{u_i} , so $\text{trop}(\|\cdot\|) = (-\log \|f_1\|, \dots, -\log \|f_n\|)$. Let $s_i \in \mathbb{Z}_{\geq 0}$ be the absolute value of the slope of $-\log \|f_i\|$ on e . Then

$$m_{\text{rel}}(e) = \gcd(s_1, \dots, s_n). \quad (5.6.1)$$

5.7 We now come to one of the key results of this section. Let $e \subset \Sigma$ be a bounded edge,⁵ and assume that $e' = \text{trop}(e)$ is an edge of $\text{Trop}(X)$ (as opposed to a vertex). The inverse image of the interior of e' under trop is a disjoint union of open annuli, one of whose skeleta is the interior of e . Hence if $x \in e$ is the unique point mapping to some $w \in \text{relint}(e') \cap N_G$, then $\mathcal{Y}_x := \tau_\Sigma^{-1}(x) \cong \mathbf{S}(1)$ is a connected component of $\mathcal{X}^w = \mathcal{U}^w \cap X^{\text{an}}$ (Definition 4.14). Let W be the affine span of e' , let \mathbf{T}' be the torus transverse to W (Section 4.26), let \mathbf{T}'' be the torus parallel to W (Remark 4.27), and choose a splitting $\mathbf{T} \rightarrow \mathbf{T}''$. We have a finite surjective morphism $\mathcal{X}^w \rightarrow \mathcal{U}^{w''}$ by Theorem 4.30, where w'' is the image of w in N_G'' .

For $y \in e$ the relative multiplicity $m_{\text{rel}}(y)$ was defined in Definition 4.21.

THEOREM 5.8 (Compatibility of multiplicities). *With the above notation,*

$$m_{\text{rel}}(e) = [\mathcal{Y}_x : \mathcal{U}^{w''}] = m_{\text{rel}}(x).$$

Moreover, $m_{\text{rel}}(e) = m_{\text{rel}}(y)$ for any y in the interior of e (even if $\text{trop}(y) \notin N_G$).

Proof. Let P be a G -rational closed interval contained in the interior of e' and containing w . Then

$$\mathcal{Y} := \tau_\Sigma^{-1}(\text{trop}^{-1}(P) \cap e) \cong \mathbf{S}(a, b)$$

is a connected component of \mathcal{X}^P ; it is a closed annulus of nonzero modulus with skeleton $\text{trop}^{-1}(P) \cap e$. Let P'' be the image of P in N_G'' , so $\ell_{\text{Trop}}(P'') = \ell_{\text{Trop}}(P)$. By Theorem 4.30 the morphism $\mathcal{Y} \rightarrow \mathcal{U}^{P''}$ is finite and surjective, and $[\mathcal{Y} : \mathcal{U}^{P''}] = [\mathcal{Y}_x : \mathcal{U}^{w''}]$ by Proposition 3.23(i). The torus \mathbf{T}'' is 1-dimensional, so $\mathcal{U}^{P''} = \text{trop}^{-1}(P'')$ is an annulus and P'' is by definition the skeleton of $\mathcal{U}^{P''}$. We have

$$m_{\text{rel}}(e) \cdot \ell_{\text{an}}(\text{trop}^{-1}(P) \cap e) = \ell_{\text{Trop}}(P) = \ell_{\text{Trop}}(P'') = [\mathcal{Y} : \mathcal{U}^{P''}] \cdot \ell_{\text{an}}(\text{trop}^{-1}(P) \cap e),$$

where the final equality is by [BPR13, Corollary 2.6]. Since x is the unique Shilov boundary point of $\mathcal{Y}_x \cong \mathbf{S}(1)$, the equality $[\mathcal{Y}_x : \mathcal{U}^{w''}] = m_{\text{rel}}(x)$ is a consequence of Proposition 4.31.

By a standard argument involving extension of the ground field (cf. Lemma 4.22 and Section 4.18), the second statement follows from the first. \square

⁵If e is an infinite ray, we can compute $m_{\text{rel}}(e)$ by refining the polyhedral structure on Σ so that $m_{\text{rel}}(e)$ coincides with $m_{\text{rel}}(\tilde{e})$ for some bounded edge \tilde{e} in the refinement. Thus we can assume without loss of generality in Theorem 5.8 that e is bounded.

COROLLARY 5.9. *Fix an edge e' of $\text{Trop}(X)$, and let e_1, \dots, e_r be the edges of Σ mapping homeomorphically onto e' . Then*

$$m_{\text{Trop}}(e') = \sum_{i=1}^r m_{\text{rel}}(e_i).$$

Proof. This follows immediately from Theorem 5.8 and Proposition 4.24. \square

Remark 5.10. With the notation in Corollary 5.9, let w be a G -rational point contained in the relative interior of e' . The affinoid space $\text{trop}^{-1}(w)$ is isomorphic to $\coprod_{i=1}^r \mathbf{S}(1)$ as in Section 5.7. Since the canonical reduction of $\mathbf{S}(1)$ is isomorphic to \mathbf{G}_m , the integer r is equal to the number of irreducible components in the canonical reduction of \mathcal{X}^w .

COROLLARY 5.11. *If e' is an edge of $\text{Trop}(X)$ and $m_{\text{Trop}}(e') = 1$, then there is a unique edge e in Σ such that trop maps e homeomorphically and isometrically onto e' . The edge e is in fact the unique geodesic segment (or ray) in X^{an} which is mapped homeomorphically by trop onto e' .*

COROLLARY 5.12. *Let $x \in \mathbf{H}(X^{\text{an}})$. Then $m_{\text{rel}}(x) > 0$ if and only if x belongs to an edge of Σ mapping homeomorphically onto its image via trop .*

Proof. Suppose that x is contained in an edge $e \subset \Sigma$ mapping homeomorphically onto its image $e' = \text{trop}(e)$. If x is in the interior of e , then $m_{\text{rel}}(x) = m_{\text{rel}}(e) > 0$ by Theorem 5.8. Otherwise $w = \text{trop}(x) \in N_G$, and x is contained in the limit boundary $\partial_{\text{lim}} \mathcal{X}^w$ of $\mathcal{X}^w = \text{trop}^{-1}(w)$ because it is a limit of points of e which are not contained in \mathcal{X}^w . Since $\partial_{\text{lim}} \mathcal{X}^w$ is the Shilov boundary of \mathcal{X}^w , by definition we have $m_{\text{rel}}(x) > 0$.

Now suppose that x is not contained in an edge of Σ mapping homeomorphically onto its image. If x has type 4, then $m_{\text{rel}}(x) = 0$ by definition. Otherwise by [BPR13, Corollary 5.1] we can enlarge Σ if necessary to assume $x \in \Sigma$. Recall that every edge of Σ maps homeomorphically onto its image or is crushed to a vertex of $\text{Trop}(X)$. By hypothesis all edges containing x are crushed to a vertex of $\text{Trop}(X)$, so $w = \text{trop}(x) \in N_G$. Hence there is an open neighborhood U of x in Σ contained in \mathcal{X}^w . Then $\tau_{\Sigma}^{-1}(U)$ is a neighborhood of x in X^{an} contained in \mathcal{X}^w , so $x \notin \partial \mathcal{X}^w$, and hence $m_{\text{rel}}(x) = 0$. \square

5.13 Slopes as orders of vanishing

The slope formula [BPR13, Theorem 5.15] provides a useful interpretation of the quantities s_i appearing in Remark 5.6 in terms of orders of vanishing. Assume that V is a strongly semistable vertex set [BPR13, § 3] of \widehat{X} (in addition to being a semistable vertex set of X). Let \mathfrak{X} be the strongly semistable formal model of \widehat{X} associated with V , let $x \in V$, and let $\overline{\mathfrak{C}}$ be the irreducible component of $\overline{\mathfrak{X}}$ whose generic point is $\text{red}(x)$. Let e be an edge of Σ adjacent to x , and let $\xi \in \overline{\mathfrak{C}}(k)$ be the reduction of the interior of e . The slope formula [BPR13, Theorem 5.15] says that if f is a nonzero rational function on \widehat{X} , then the slope s of $-\log |f|$ along e (in the direction away from x) is equal to $\text{ord}_{\xi}(\tilde{f}_x)$. One can use this fact to give a simple proof of the well-known balancing formula for tropical curves.

THEOREM 5.14 (Balancing formula for tropical curves). *Let w be a vertex of $\text{Trop}(X)$, and let $\vec{v}_1, \dots, \vec{v}_t$ be the primitive integer tangent directions at w corresponding to the various edges e'_1, \dots, e'_t incident to w . Then $\sum_{j=1}^t m_{\text{Trop}}(e'_j) \vec{v}_j = 0$.*

Proof. We use the setup of Section 5.13. Let $f_1, \dots, f_n \in K[X]^{\times}$ be the coordinate functions as in Remark 5.6, and let $F_i = -\log |f_i|$. Let $x \in \Sigma \cap \text{trop}^{-1}(w)$ be a vertex. By the slope

formula [BPR13, Theorem 5.15] we have $0 = \sum_{v \in T_x} d_v F_i(x)$ for each $i = 1, \dots, n$. Since we are assuming that V is a strongly semistable vertex set, each tangent direction $v \in T_x$ along which some F_i has nonzero slope is represented by a unique edge $e_v = [x, y_v]$ of Σ adjoining x , and $d_v F_i(x)$ is just the slope of $-\log |f_i|$ along e_v . If $\text{trop}(e_v) = \{w\}$, then $d_v F_i(x) = 0$ for all i , and otherwise

$$d_v F_i(x) = \frac{\log |f_i(x)| - \log |f_i(y_v)|}{\ell_{\text{an}}(e_v)} = m_{\text{rel}}(e_v) \frac{\log |f_i(x)| - \log |f_i(y_v)|}{\ell_{\text{Trop}}(\text{trop}(e_v))}.$$

By Corollary 5.9, for each i we have

$$\sum_{j=1}^t m_{\text{Trop}}(e'_j)(\vec{v}_j)_i = \sum_{j=1}^t \left(\sum_{e=[x,y] \xrightarrow{\sim} e'_j} m_{\text{rel}}(e) \right) \frac{\log |f_i(x)| - \log |f_i(y)|}{\ell_{\text{Trop}}(e'_j)} = \sum_{x \rightarrow w} \sum_{v \in T_x} d_v F_i(x) = 0,$$

which implies the result. \square

5.15 Faithful representations

If Y_Δ is a proper toric variety with dense torus \mathbf{T} , then $X \hookrightarrow \mathbf{T}$ extends in a unique way to a morphism $\iota: \widehat{X} \rightarrow Y_\Delta$, which is a closed immersion for suitable Y_Δ . The intersection $\widehat{X}^{\text{an}} \cap (Y_\Delta^{\text{an}} \setminus \mathbf{T}^{\text{an}})$ is the finite set of type 1 points $D = \widehat{X}^{\text{an}} \setminus X^{\text{an}}$. We write $\text{trop}(\iota): \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta)$ for the induced tropicalization map, and we set $\text{Trop}(\widehat{X}, \iota) = \text{trop}(\iota)(\widehat{X}^{\text{an}}) \subset N_{\mathbb{R}}(\Delta)$.

5.15.1 Let Y_Δ and $Y_{\Delta'}$ be toric varieties with dense tori \mathbf{T} and \mathbf{T}' , respectively, and let $\iota: \widehat{X} \hookrightarrow Y_\Delta$ and $\iota': \widehat{X} \hookrightarrow Y_{\Delta'}$ be closed immersions whose images meet the dense torus. We say that ι' *dominates* ι and we write $\iota' \geq \iota$ provided that there exists a morphism $\psi: Y_{\Delta'} \rightarrow Y_\Delta$ of toric varieties such that $\psi \circ \iota' = \iota$. In this case we have an induced morphism $\text{Trop}(\psi): \text{Trop}(\widehat{X}, \iota') \rightarrow \text{Trop}(\widehat{X}, \iota)$ making the triangle

$$\begin{array}{ccc} & \widehat{X}^{\text{an}} & \\ \text{trop}(\iota') \swarrow & & \searrow \text{trop}(\iota) \\ \text{Trop}(\widehat{X}, \iota') & \xrightarrow{\text{Trop}(\psi)} & \text{Trop}(\widehat{X}, \iota) \end{array}$$

commute. Since $\text{trop}(\iota)$ and $\text{trop}(\iota')$ are surjective, the map $\text{Trop}(\psi)$ is independent of the choice of ψ , so the set of ‘tropicalizations of toric embeddings’ is a filtered inverse system.

5.15.2 By a *finite subgraph* of \widehat{X}^{an} we mean a *connected* compact subgraph of a skeleton of \widehat{X} . Any finite union of geodesic segments in $\mathbf{H}_o(\widehat{X}^{\text{an}})$ is contained in a skeleton by [BPR13, Corollary 5.10], so we can equivalently define a finite subgraph of \widehat{X}^{an} to be an isometric embedding of a finite connected metric graph Γ into $\mathbf{H}_o(\widehat{X}^{\text{an}})$. Let $\widehat{X} \hookrightarrow Y_\Delta$ be a closed immersion into a toric variety with dense torus \mathbf{T} such that $\widehat{X} \cap \mathbf{T} \neq \emptyset$. We say that a finite subgraph Γ of \widehat{X}^{an} is *faithfully represented* by $\text{trop}: \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta)$ if trop maps Γ homeomorphically and isometrically onto its image Γ' (which is contained in $N_{\mathbb{R}}(\Delta)$). We say that trop is *faithful* if it faithfully represents a skeleton Σ of \widehat{X} .

Remark 5.16. When considering a closed connected subset Γ of $\mathbf{H}(X^{\text{an}})$ or Γ' of $\text{Trop}(X)$, we will always implicitly endow it with the shortest-path metric. In general this is *not* the same as the metric on Γ (respectively, Γ') induced by the (shortest-path) metric on $\mathbf{H}(X^{\text{an}})$ (respectively, $\text{Trop}(X)$). With this convention, Γ (respectively, Γ') is a *length space* in the sense of [Pap05,

Definition 2.1.2], so any homeomorphism $\Gamma \rightarrow \Gamma'$ which is a local isometry is automatically an isometry by Corollary 3.4.6 of [Pap05]. This will be used several times in what follows.

The following result shows that if Γ is faithfully represented by a given tropicalization, then it is also faithfully represented by all ‘larger’ tropicalizations.

LEMMA 5.17. *Let $\iota: \widehat{X} \hookrightarrow Y_\Delta$ and $\iota': \widehat{X} \hookrightarrow Y_{\Delta'}$ be closed immersions of \widehat{X} into toric varieties whose images meet the dense torus and which satisfy $\iota' \geq \iota$. If a finite subgraph Γ of \widehat{X}^{an} is faithfully represented by $\text{trop}(\iota): \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta)$, then Γ is faithfully represented by $\text{trop}(\iota'): \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta')$.*

Proof. Without loss of generality, we may replace \widehat{X} by $X = \widehat{X} \setminus D$ and assume that $Y_{\Delta'}$ and Y_Δ are tori with $\iota = (f_1, \dots, f_n)$ and $\iota' = (f_1, \dots, f_m)$ for some $m \geq n$. The result is now clear from (5.6.1). \square

We will show in Theorem 5.20 that any finite subgraph of \widehat{X}^{an} is faithfully represented by some tropicalization. First we need two lemmas.

LEMMA 5.18. *Let e be an edge of a skeleton Σ of \widehat{X} with distinct endpoints x and y . There exists a nonzero meromorphic function f on \widehat{X} such that $F = -\log|f|$ has the following properties:*

- (i) *We have $F \geq 0$ on e , and $F(x) = F(y) = 0$.*
- (ii) *There exist (not necessarily distinct) type 2 points x' and y' in the interior e° of e such that $\rho(x, x') = \rho(y, y')$, such that F has slope ± 1 on $[x, x']$ and $[y, y']$, and such that F is constant on $[x', y']$, as shown in Figure 4.*




FIGURE 4. The graph of the function $F = -\log|f|: e \rightarrow \mathbb{R}_{\geq 0}$ constructed in Lemma 5.18.

Proof. Without loss of generality, we may assume $\Sigma = \Sigma(\widehat{X}, V(\mathfrak{X}))$ for a strongly semistable formal model \mathfrak{X} of \widehat{X} [BPR13, § 4]. For each irreducible component $\overline{\mathfrak{C}}_\nu$ of $\overline{\mathfrak{X}}$, a simple argument using the Riemann–Roch theorem allows us to choose a rational function \tilde{f}_ν on $\overline{\mathfrak{C}}_\nu$ which vanishes to order 1 at every singular point of $\overline{\mathfrak{X}}$ lying on $\overline{\mathfrak{C}}_\nu$. By [BL85, Corollary 3.8] there exists a nonzero rational function \tilde{f} on \widehat{X} whose poles all reduce to smooth points of $\overline{\mathfrak{X}}$ and which induces the rational function \tilde{f}_ν on each irreducible component $\overline{\mathfrak{C}}_\nu$ of $\overline{\mathfrak{X}}$ (the gluing condition from loc. cit. is trivial in this situation). The function f constructed in the proof of loc. cit. is defined on an affinoid domain U of \widehat{X}^{an} containing x and y , and \tilde{f}_x and \tilde{f}_y are the restrictions of the residue of f in the canonical reduction of U . Therefore we have $|f(x)| = |f(y)| = 1$. Since $\{x, y\}$ is the Shilov boundary of $\tau_\Sigma^{-1}(e)$, this proves part (i).

By [BPR13, Theorem 5.15(3)] the outgoing slope of F at x or y in the direction of e is 1. Let ξ be the singular point of $\overline{\mathfrak{X}}$ whose formal fiber is $\tau_\Sigma^{-1}(e^\circ)$. Since f has no poles on the formal fibers above singular points, f restricts to an analytic function on the open annulus $\tau_\Sigma^{-1}(e^\circ)$. Part (ii) now follows from part (i) and [BPR13, Proposition 2.10(1)]. \square

LEMMA 5.19. *Let $A \subset \widehat{X}^{\text{an}}$ be an affinoid domain isomorphic to a closed annulus $\mathbf{S}(a)$ with nonzero modulus. There exists a nonzero meromorphic function f on \widehat{X} such that $F = -\log |f|$ is linear with slope ± 1 on $\Sigma(A)$.*

Proof. Choose an identification of A with $\mathbf{S}(a) = \mathcal{M}(K\langle at^{-1}, t \rangle)$. By [FM86, Théorème 4, § 2.4] the ring of meromorphic functions on \widehat{X} which are regular on A is dense in $K\langle at^{-1}, t \rangle$. Hence there exists a meromorphic function f on \widehat{X} such that $f \in K\langle at^{-1}, t \rangle$ and $|f - t|_{\text{sup}} < 1$. It follows from [BPR13, Proposition 2.2] that f is also a parameter for the annulus A , so $-\log |f|$ is linear with slope ± 1 on $\Sigma(A)$. \square

THEOREM 5.20. *If Γ is any finite subgraph of \widehat{X}^{an} , then there is a closed immersion $\widehat{X} \hookrightarrow Y_{\Delta}$ of \widehat{X} into a quasiprojective toric variety Y_{Δ} such that $\text{trop}: \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta)$ faithfully represents Γ . In particular, there exists a faithful tropicalization.*

Proof. Since Γ is by definition contained in a skeleton Σ , we may assume $\Gamma = \Sigma$. Taking a refinement of Σ if necessary, we assume without loss of generality that Σ does not have any loop edges. We claim that after possibly refining Σ further, for each edge $e \subset \Sigma$ there exists a nonzero meromorphic function f on \widehat{X} such that $\log |f|$ has slope ± 1 on e .

Let $e = [x, y]$ be an edge of Σ , and let f and $x', y' \in e^{\circ}$ be as in Lemma 5.18. Then $[x, x']$ and $[y', y]$ are edges in a refinement of Σ , and $\log |f|$ has slope ± 1 on $[x, x']$ and $[y', y]$. If $x' = y'$, then we are done with e ; otherwise we let $e' = [x', y']$. By construction $e' \subset e^{\circ}$, so $\tau_{\Sigma}^{-1}(e')$ is a closed annulus of nonzero modulus, and we may apply Lemma 5.19 to find f' such that $\log |f'|$ has slope ± 1 on e' . This proves the claim.

By (5.6.1), if $\Phi = \{f_1, \dots, f_r\}$ is any collection of meromorphic functions on X such that (i) for each edge e of Σ there is an i such that $\log |f_i|$ has slope ± 1 on e , and (ii) $\phi = (f_1, \dots, f_r)$ induces a closed immersion of a dense open subscheme X of \widehat{X} into a torus $\mathbf{T} \cong \mathbf{G}_m^r$, then $\text{trop} \circ \phi$ maps each edge of Σ isometrically onto its image. Since ϕ extends to a closed immersion $\widehat{X} \hookrightarrow Y_{\Delta}$ into a suitable compactification Y_{Δ} of \mathbf{T} , it only remains to show that we can enlarge Φ so that $\phi|_{\Sigma}$ is injective; then $\text{trop} \circ \phi$ maps Σ isometrically onto its image by Remark 5.16.

Let e be an edge of Σ . Since Σ has at least two edges, the proof of [BPR13, Theorem 4.11] (specifically case 1 in (4.15.1) of [BPR13]) shows that $\tau_{\Sigma}^{-1}(e)$ is an affinoid domain in \widehat{X}^{an} . By [FM86, Théorème 1, § 1.4], there is a meromorphic function f on \widehat{X} such that $\tau_{\Sigma}^{-1}(e) = \{x \in \widehat{X}^{\text{an}} : |f(x)| \leq 1\}$. Adding such an f to Φ for every edge e , we may assume that $\text{trop} \circ \phi$ is injective on $\Sigma \setminus V$, that is, that if $\text{trop}(\phi(x)) = \text{trop}(\phi(y))$ for $x, y \in \Sigma$, then x and y are vertices. By the definition of \widehat{X}^{an} , if $x, y \in \widehat{X}^{\text{an}}$ are distinct points, then there exists a meromorphic function f on \widehat{X}^{an} such that $|f(x)| \neq |f(y)|$. Adding such f to Φ for every pair of vertices yields a faithful tropicalization. \square

We obtain the following theorem as a consequence.

THEOREM 5.21. *Let Γ be a finite subgraph of \widehat{X} . Then there exists a quasiprojective toric embedding $\iota: \widehat{X} \hookrightarrow Y_{\Delta}$ such that for every quasiprojective toric embedding $\iota': \widehat{X} \hookrightarrow Y_{\Delta'}$ with $\iota' \geq \iota$, the tropicalization map $\text{trop}(\iota'): \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta')$ maps Γ homeomorphically and isometrically onto its image.*

Proof. By Theorem 5.20, there exists a closed embedding ι such that $\text{trop}(\iota)$ maps Γ homeomorphically and isometrically onto its image. By Lemma 5.17, the same property holds for any closed embedding $\iota' \geq \iota$. \square

As mentioned in the introduction, Theorem 5.21 can be interpreted colloquially as saying that the homeomorphism in Theorem 1.2 is an isometry.

With a little more work, we obtain the following strengthening of Theorem 5.21 in which the finite metric graph Γ is replaced by an arbitrary skeleton of X (which is no longer required to be compact or of finite length).

THEOREM 5.22. *Let Σ be any skeleton of X . Then there exists a quasiprojective toric embedding $\iota: X \hookrightarrow Y_\Delta$ such that for every quasiprojective toric embedding $\iota': X \hookrightarrow Y_{\Delta'}$ with $\iota' \geq \iota$, the tropicalization map $\text{trop}(\iota'): \widehat{X}^{\text{an}} \rightarrow N_{\mathbb{R}}(\Delta')$ maps Σ homeomorphically and isometrically onto its image.*

Proof. Using Lemma 5.17, it suffices to prove that there exists a closed embedding ι such that $\text{trop}(\iota)$ maps Σ homeomorphically and isometrically onto its image.

For each point $p \in D = \widehat{X} \setminus X$, choose a pair of relatively prime integers $m_1(p), m_2(p)$ greater than $2g$, where g is the genus of \widehat{X} . By the Riemann–Roch theorem, there are rational functions $f_1^{(p)}$ and $f_2^{(p)}$ on \widehat{X} such that $f_i^{(p)}$ has a pole of exact order $m_i(p)$ at p and no other poles for $i = 1, 2$. Let U_p be an (analytic) open neighborhood of p on which $f_1^{(p)}$ and $f_2^{(p)}$ have no zeros, and let U be the union of U_p for all $p \in D$.

Let $\Gamma = \Sigma \setminus (\Sigma \cap U)$. Then Γ is a finite subgraph of X , hence by Theorem 5.20 there exists a closed embedding ι_0 such that $\text{trop}(\iota_0)$ maps Γ homeomorphically and isometrically onto its image. We can choose the U_p such that the complement $\Sigma \setminus \Gamma$ consists of finitely many open infinite rays r_p , one for each point $p \in D$. By the slope formula, the absolute value of the slope of $\log |f_i^{(p)}|$ along r_p is $m_i(p)$. Since $\gcd(m_1(p), m_2(p)) = 1$ for all $p \in D$, if we enhance the embedding ι_0 to a larger embedding ι by adding the coordinate functions $f_1^{(p)}$ and $f_2^{(p)}$ for all $p \in D$, $\text{trop}(\iota)$ has multiplicity 1 along each ray r_p by Remark 5.6. By Lemma 5.17, $\text{trop}(\iota)$ also has multiplicity 1 at every edge of Γ . It follows easily (as in the proof of Theorem 5.20) that $\text{trop}(\iota)$ maps Σ homeomorphically and isometrically onto its image, as desired. \square

5.23 Certifying faithfulness

It is useful to be able to certify that a given tropicalization map is faithful using only ‘tropical’ computations.

THEOREM 5.24. *Let Γ' be a compact connected subset of $\text{Trop}(X)$, and suppose that $m_{\text{Trop}}(w) = 1$ for all $w \in \Gamma' \cap N_G$. Then there is a unique closed subset $\Gamma \subset \mathbf{H}_o(X^{\text{an}})$ mapping homeomorphically onto Γ' , and this homeomorphism is an isometry.*

Proof. Since m_{Trop} is constant along the interior of each edge of $\text{Trop}(X)$ by Theorem 4.28(i) (for points not contained in N_G this is proved by a standard ground-field extension argument) and Γ' is a finite union of closed intervals, we have $m_{\text{Trop}}(w) = 1$ for all $w \in \Gamma'$. By Proposition 4.24, for each $w \in \Gamma'$ there is a unique point $x = x_w \in \mathbf{H}_o(X^{\text{an}})$ such that $\text{trop}(x) = w$ and $m_{\text{rel}}(x) > 0$. Let $\Gamma = \{x_w : w \in \Gamma'\}$. The natural continuous map $\text{trop}: \Gamma \rightarrow \Gamma'$ is bijective. It follows from Corollaries 5.11 and 5.12 that Γ is also a finite union of closed intervals, hence compact. Thus $\text{trop}: \Gamma \rightarrow \Gamma'$, being a continuous bijection between compact Hausdorff spaces, is a homeomorphism. By Corollary 5.11 this homeomorphism is an isometry.

As for the uniqueness of Γ , let $\widetilde{\Gamma}$ be any closed subset of $\mathbf{H}_o(X^{\text{an}})$ mapping homeomorphically onto Γ' . Fix $w \in \Gamma'$, and let x be the point in $\widetilde{\Gamma}$ with $\text{trop}(x) = w$. Since x belongs to a closed

segment of X^{an} mapping homeomorphically onto its image via trop (namely the inverse image in $\widetilde{\Gamma}$ of an edge in $\text{Trop}(X)$ containing w), it follows from Corollary 5.12 that $m_{\text{rel}}(x) > 0$. Hence $x = x_w$, so $\widetilde{\Gamma} = \Gamma$. \square

In order to apply Theorem 5.24, it is useful to know that one can sometimes determine the multiplicity at a point $w \in \text{Trop}(X) \cap N_G$ just from the local structure of $\text{Trop}(X)$ at w , that is, from the combinatorics of $\text{Star}(w)$. Recall that if $\vec{v}_1, \dots, \vec{v}_r$ are the primitive generators of the edge directions in $\text{Trop}(X)$ at w , and a_i is the tropical multiplicity of the edge corresponding to \vec{v}_i , then the balancing condition says that $a_1\vec{v}_1 + \dots + a_r\vec{v}_r = 0$. Now, if Z_1, \dots, Z_s are the irreducible components of $\text{in}_w(X)$, then the tropicalization of each Z_j (as a subscheme of the torus torsor $\overline{\mathbf{T}}^w$ over the trivially valued field k) is a union of rays spanned by a subset of $\{\vec{v}_1, \dots, \vec{v}_r\}$. If b_{ij} is the multiplicity of the ray spanned by \vec{v}_i in $\text{Trop}(Z_j)$ and m_i is the multiplicity of Z_i in $\text{in}_w(X)$, then the balancing condition implies $b_{1j}\vec{v}_1 + \dots + b_{rj}\vec{v}_r = 0$, and we also have $m_1b_{i1} + \dots + m_sb_{is} = a_i$, since $\text{Trop}(\text{in}_w(X)) = \text{Star}_w(\text{Trop}(X))$ by [Spe14, Proposition 10.1].

THEOREM 5.25. *Let $w \in \text{Trop}(X) \cap N_G$. If $\text{Trop}(X)$ is trivalent at w and one of the edges adjacent to w has multiplicity 1, then w has multiplicity 1.*

Proof. Let $\vec{v}_1, \vec{v}_2, \vec{v}_3$ be the primitive generators of the edge directions in $\text{Trop}(X)$ at w , and let a_i be the multiplicity of the edge in direction \vec{v}_i . The linear span $\langle \vec{v}_1, \vec{v}_2, \vec{v}_3 \rangle$ is 2-dimensional, since the \vec{v}_i are distinct and satisfy the balancing condition, so any relation among them is a scalar multiple of the relation $a_1\vec{v}_1 + a_2\vec{v}_2 + a_3\vec{v}_3 = 0$.

Let Z be an irreducible component of $\text{in}_w(X)$, and let b_i be the multiplicity of the ray spanned by \vec{v}_i in $\text{Trop}(Z)$. Then b_i is a nonnegative integer bounded above by a_i and $b_1\vec{v}_1 + b_2\vec{v}_2 + b_3\vec{v}_3 = 0$. This relation must be a scalar multiple of the relations given by the a_i , so there is a positive rational number $\lambda \leq 1$ such that $b_i = \lambda a_i$ for all i . If some a_i is 1, then λ must also be 1 and $b_i = a_i$ for all i . Since a_i is the sum of the multiplicities of the ray spanned by \vec{v}_i in the tropicalizations of the components of $\text{in}_w(X)$, it follows that $\text{in}_w(X)$ has no other components, and w has multiplicity 1. \square

Remark 5.26. Initial degenerations at interior points of an edge of multiplicity 1 are always smooth, since they are isomorphic to \mathbf{G}_m by Theorem 4.28.

Remark 5.27. There are other natural combinatorial conditions which can guarantee multiplicity 1 at a point $w \in N_G$ of $\text{Trop}(X)$ or smoothness of the corresponding initial degeneration. In the case of curves, for example, the argument above works more generally if $\text{Trop}(X)$ is r -valent, the linear span of the edge directions at w has dimension $r - 1$, and the multiplicities of the edges at w have no nontrivial common factor.

Combining the previous two results and the discussion of tropical hypersurfaces in Section 2.1, we obtain the following. Recall that a *leaf* in a graph is a vertex of valence 1.

COROLLARY 5.28. *Suppose $g(\widehat{X}) \geq 1$, and let Σ be the minimal skeleton of \widehat{X}^{an} .*

(i) *If all vertices of $\text{Trop}(X)$ are trivalent, all edges of $\text{Trop}(X)$ have multiplicity 1, the minimal skeleton Σ has no leaves,⁶ and $\dim H_1(\Sigma, \mathbb{R}) = \dim H_1(\text{Trop}(X), \mathbb{R})$, then $\text{trop}: \Sigma \rightarrow \text{Trop}(X)$ is an isometry onto its image.*

⁶In earlier versions of this paper, the skeleton Σ was assumed to be bridgeless; however, the proof only used that Σ has no leaves.

(ii) If $X \subset \mathbf{G}_m^2$ is defined by a polynomial $f \in K[x, y]$ whose Newton complex (see Remark 2.2) is a unimodular triangulation, then \widehat{X} has totally degenerate reduction and $\text{trop}: \Sigma \rightarrow \text{Trop}(X)$ induces an isometry from Σ onto its image.

Proof. Let $g' = \dim H_1(\Sigma, \mathbb{R}) = \dim H_1(\text{Trop}(X), \mathbb{R})$. Choose a compact, connected subgraph $\Sigma' \subset \text{Trop}(X)$ with $g' = \dim H_1(\Sigma', \mathbb{R})$. By Theorem 5.25, all $w \in \text{Trop}(X) \cap N_G$ have multiplicity 1, so according to Theorem 5.24, there is a (unique) finite subgraph $\widetilde{\Sigma}$ of X^{an} mapping homeomorphically and isometrically onto Σ' . Since Σ is the minimal skeleton of X^{an} and Σ has no leaves, any subgraph of X^{an} whose first homology has dimension at least g' must contain Σ . In particular, $\Sigma \subset \widetilde{\Sigma}$, so Σ maps isometrically onto its image, which proves part (i).

We now prove part (ii). By Section 2.1, $\text{Trop}(X)$ is trivalent with all edges of multiplicity 1. By Baker's theorem [Bak93] (see also [Bee09]), the genus $g(\widehat{X})$ is bounded by the number of internal vertices in the Newton polytope of f . Since the Newton complex is dual to the tropicalization, each internal vertex corresponds to a region in $\mathbb{R}^2 \setminus \text{Trop}(X)$; from this and the above, one sees that $g(\widehat{X}) \leq \dim H_1(\text{Trop}(X), \mathbb{R})$. With Σ' as above, by Theorem 5.24, there is a (unique) finite subgraph $\widetilde{\Sigma}$ of X^{an} mapping homeomorphically onto Σ' via trop . Since $\dim H_1(\Sigma, \mathbb{R}) \geq \dim H_1(\widetilde{\Sigma}, \mathbb{R}) \geq g(\widehat{X})$ in any case, it follows from the genus formula [BPR13, Remark 4.18] that $\dim H_1(\Sigma, \mathbb{R}) = g(\widehat{X})$ and that \widehat{X} has totally degenerate reduction. In particular, the minimal skeleton Σ has no leaves. By part (i), $\text{trop}: \Sigma \rightarrow \text{Trop}(X)$ is an isometry onto its image. \square

Corollary 5.28 will be important for applications to Tate curves in Section 6.

EXAMPLE 5.29. As an example where the hypotheses of Corollary 5.28 are satisfied, consider the genus 3 curve $X = V(f)$ with

$$f = t^4(x^4 + y^4 + z^4) + t^2(x^3y + x^3z + xy^3 + xz^3 + y^3z + yz^3) + t(x^2y^2 + x^2z^2 + y^2z^2) + x^2yz + xy^2z + xyz^2.$$

See Figure 5.




FIGURE 5. The Newton complex and tropicalization of the curve X defined by the polynomial f from Example 5.29. The tropicalization faithfully represents the minimal skeleton of X^{an} .

EXAMPLE 5.30. The following example shows that it is possible, even under the hypotheses of Theorem 5.24, for the tropicalization map to fail to be faithful. Let K be the completion

of $C\{\{t\}\}$, let $\widehat{X} \subset \mathbb{P}^2$ be defined by the equation

$$(y - 1)^2 = (x - 1)^2(y + 1) + t \cdot xy$$

over $\mathbb{C}\{\{t\}\}$, and let $X = \widehat{X} \cap \mathbf{G}_m^2$. The above equation degenerates to a nodal rational curve when $t = 0$, the node being $[1 : 1 : 0]$, so it defines a (not strongly) semistable algebraic integral model \mathcal{X} of \widehat{X} —in fact, \mathcal{X} is a minimal stable model for \widehat{X} , and the associated semistable vertex set only contains one point, so it is minimal as well (see [BPR13, Corollary 4.23]). Therefore \widehat{X} is an elliptic curve with bad reduction and $\text{in}_0(X) \cong \overline{\mathcal{X}} \cap \mathbf{G}_{m,k}^2$ is reduced and irreducible. The points in the set of punctures

$$D := \widehat{X}(K) \setminus X(K) = \{[0 : 0 : 1], [0 : 1 : 0], [1 : 0 : 0], [0 : 3 : 1], [2 : 0 : 1]\}$$

reduce to distinct smooth points of $\overline{\mathcal{X}}$, so if Σ is the minimal skeleton of \widehat{X}^{an} and $\tau_\Sigma: \widehat{X}^{\text{an}} \rightarrow \Sigma$ is the retraction, then $\tau_\Sigma(x)$ reduces to the generic point of $\overline{\mathcal{X}}$ for all $x \in D$. Therefore the minimal skeleton Γ of X and the tropicalization of X are as shown in Figure 6. We see that $\text{Trop}(X)$ is contractible and has everywhere multiplicity 1 but the image of the section does not contain the loop in X^{an} (the loop is contracted to the origin). In particular, trop is not faithful despite the fact that all points in $\text{Trop}(X)$ have multiplicity 1.




FIGURE 6. The skeleton, tropicalization, and Newton complex of the curve X from Example 5.30. One sees from the Newton complex that the initial degenerations are all of multiplicity 1 away from 0, and $\text{in}_0(X)$ is a rational nodal curve. However, the tropicalization crushes the loop in X^{an} to the origin.

6. Elliptic curves

Let \widehat{E}/K be an elliptic curve. If \widehat{E} has good reduction, then the minimal skeleton Σ of \widehat{E}^{an} is a point, while if \widehat{E} has multiplicative reduction, then the minimal skeleton Σ of \widehat{E}^{an} is homeomorphic to a circle of length $-\text{val}(j_{\widehat{E}}) = \text{val}(q_{\widehat{E}})$, where $\widehat{E}^{\text{an}} \cong \mathbf{G}_m^{\text{an}}/q_{\widehat{E}}^{\mathbb{Z}}$ is the Tate uniformization of \widehat{E} (see [BPR13, Remark 4.24]). In this section, we use our results on nonarchimedean analytic curves and their tropicalizations to prove some new results (and reinterpret some old results) about tropicalizations of elliptic curves.

6.1 Faithful tropicalization of elliptic curves

As noted in [KMM08, KMM09], a curve in \mathbf{G}_m^2 given by a Weierstrass equation $y^2 = x^3 + ax^2 + bx + c$ cannot have a cycle in its tropicalization, because the Newton complex of a Weierstrass equation does not have an interior vertex. Thus Weierstrass equations are always ‘bad’ from the point of view of tropical geometry. On the other hand, the following result shows that there do always exist ‘good’ plane embeddings of elliptic curves with multiplicative reduction.




FIGURE 7. The skeleton Γ of E , the tropicalization of E , and the Newton complex of the equation $ax^2y + bxy^2 + cxyz = dz^3$ defining E , where E is as in the proof of Theorem 6.2. The minimal skeleton Σ of \widehat{E} is the circle contained in Γ . The tropicalization faithfully represents Σ , so $\ell_{\text{Trop}}(\text{Trop}(\Sigma)) = \ell_{\text{an}}(\Sigma)$.

THEOREM 6.2. *Let \widehat{E}/K be an elliptic curve with multiplicative reduction. Then there is a closed embedding of \widehat{E} in \mathbb{P}^2 , given by a projective plane equation of the form $ax^2y + bxy^2 + cxyz = dz^3$, such that (letting E be the open affine subset of \widehat{E} mapping into the torus \mathbf{G}_m^2) $\text{Trop}(E)$ is a trivalent graph, every point of $\text{Trop}(E)$ has a smooth and irreducible initial degeneration (hence tropical multiplicity 1), and the minimal skeleton of \widehat{E} is faithfully represented by the tropicalization map. In particular, $\text{Trop}(E)$ contains a cycle of length $-\text{val}(j_{\widehat{E}})$.*

Proof. Let $q = q_{\widehat{E}}$ be the Tate parameter, so that $\widehat{E}^{\text{an}} \cong \mathbf{G}_m^{\text{an}}/q^{\mathbb{Z}}$. Choose a cube root $q^{1/3} \in K^\times$ of q , and let $\alpha, \beta \in \widehat{E}(K)$ correspond under the Tate isomorphism to the classes of $q^{1/3}$ and $(q^{1/3})^2$, respectively. Recall that a divisor $D = \sum a_P(P)$ on an elliptic curve is principal if and only if $\sum a_P = 0$ and $\sum a_PP = 0$ in the group law on the curve. In particular, there exist rational functions f and g on \widehat{E} (unique up to multiplication by a nonzero constant) such that $\text{div}(f) = 2(\alpha) - (\beta) - (0)$ and $\text{div}(g) = 2(\beta) - (\alpha) - (0)$. Let $\psi: \widehat{E} \rightarrow \mathbb{P}^2$ be the morphism associated with the rational map $[f : g : 1]$. Since $1, f, g$ form a basis for $L(D)$ with $D = (\alpha) + (\beta) + (0)$ and D is very ample [Har77, Corollary IV.3.2(b)], we see that ψ is a closed immersion.

Let Γ be the minimal skeleton of E , that is, the smallest closed connected subset of \widehat{E}^{an} containing the skeleton Σ of \widehat{E}^{an} and the three points $\alpha, \beta, 0$, with those three points removed. Recall that Σ is isometric to a circle of circumference $\ell_{\text{an}}(\Sigma) = -\text{val}(j_{\widehat{E}}) = \text{val}(q)$. The natural map

$$K^\times \rightarrow \widehat{E}(K) \hookrightarrow \widehat{E}^{\text{an}} \rightarrow \Sigma \xrightarrow{\sim} \mathbb{R}/\ell\mathbb{Z}$$

is given by $z \mapsto [\text{val}(z)]$; in particular, if $\tau_\Sigma: \widehat{E}^{\text{an}} \rightarrow \Sigma$ denotes the canonical retraction, we have $\tau_\Sigma(0) = [0]$, $\tau_\Sigma(\alpha) = [\frac{1}{3}\text{val}(q)]$, and $\tau_\Sigma(\beta) = [\frac{2}{3}\text{val}(q)]$. Thus Γ is a circle with an infinite ray emanating from each of three equally spaced points $O = \tau_\Sigma(0)$, $A = \tau_\Sigma(\alpha)$, $B = \tau_\Sigma(\beta)$ along the circle (see Figure 7). The tropicalization map $\text{trop}: E = \widehat{E}^{\text{an}} \setminus \{0, \alpha, \beta\} \rightarrow \mathbb{R}^2$ corresponding to the embedding $E \hookrightarrow \mathbf{G}_m^2$ given by (f, g) factors through the retraction onto Γ .

The map $\text{trop}: \Gamma \rightarrow \mathbb{R}^2$ can be determined (up to an additive translation) using the slope formula [BPR13, Theorem 5.15] by solving an elementary graph potential problem. The result is as follows. The function $\text{val}(f) = -\log|f|$ has slope -1 along the ray from O to 0 , slope 2 along the ray from A to α , and slope -1 along the ray from B to β . On Σ , it has slope 1 along the segment from O to A , slope -1 along the segment from A to B , and slope 0 along the segment

from B to O . Similarly, the function $\text{val}(g) = -\log |g|$ has slope -1 along the ray from O to 0 , slope -1 along the ray from A to α , and slope 2 along the ray from B to β . On Σ , it has slope 0 along the segment from O to A , slope 1 along the segment from A to B , and slope -1 along the segment from B to O . Thus (up to a translation on \mathbb{R}^2) $\text{Trop}(E)$ is a trivalent graph consisting of a triangle with an infinite ray emanating from each of the vertices, as in Figure 7.

Since the expansion factor along every edge of Γ is equal to 1 by (5.6.1), it follows from Corollary 5.9 that the tropical multiplicity of every edge of $\text{Trop}(E)$ is 1. By Theorem 5.25, the multiplicity at every vertex of $\text{Trop}(E)$ is 1 as well, and in fact the initial degenerations are smooth and irreducible since the Newton complex is unimodular (see Figure 7). Since the expansion factor is 1 along every edge of Σ and $\text{trop}|_{\Sigma}$ is a homeomorphism, it follows that Σ is faithfully represented. The bounded edges of $\text{Trop}(E)$ form a triangle each of whose sides has lattice length $\text{val}(q)/3$.

The only thing which remains to be proved is that $\psi(\widehat{E}) \subset \mathbb{P}^2$ is cut out by an equation of the form indicated in the statement of the theorem. This follows from the Riemann–Roch theorem: the functions $1, fg, f^2g, fg^2$ all belong to the 3-dimensional vector space $L(3(0))$ and hence there is a nonzero linear relation between them. (This argument is similar to [Har77, Proposition IV.4.6].) \square

We can also use our theorems to give more conceptual proofs of many of the results from [KMM08, KMM09]. For example, we have the following theorem which was proved in [KMM09] by a brute-force computation.

THEOREM 6.3. *Let $E \subset \mathbf{G}_m^2$ be the intersection of an elliptic curve $\widehat{E} \subset \mathbb{P}^2$ with \mathbf{G}_m^2 . Assume that (i) $\text{Trop}(E)$ contains a cycle C , (ii) all edges of $\text{Trop}(E)$ have multiplicity 1, and (iii) $\text{Trop}(E)$ is trivalent. Then $\ell_{\text{Trop}}(C) = -\text{val}(j_{\widehat{E}})$.*

Proof. This follows immediately from Corollary 5.28(i). \square

Remark 6.4. Conditions (i)–(iii) from Theorem 6.3 are automatically satisfied if the Newton complex of the defining polynomial for E is a unimodular triangulation with a vertex lying in the interior of the Newton polygon, as in Figure 7. Varying the valuation of the coefficient corresponding to the interior vertex while keeping all other coefficients fixed gives a natural map from an annulus in \mathbf{G}_m to the j -line, which is finite and flat onto an annulus in the j -line, by [BPR13, Proposition 2.2(2)]. In particular, given a tropical plane curve dual to such a Newton complex and an elliptic curve E with j -invariant equal to minus the length of the loop, there is an embedding of E into a toric variety such that the tropicalization of the intersection with \mathbf{G}_m^2 is faithful and equal to the given tropical curve. See [CS13] for explicit constructions of such embeddings for tropical curves of ‘honeycomb normal form’, including an algorithm for finding the honeycomb form of an elliptic curve, parameterization by theta functions, the tropicalization of the inflection points, and relations to the group law.

Remark 6.5. A different (but related) conceptual explanation for Theorem 6.3 is given in [HK12, Proposition 7.7].

Let us say that a closed embedding of an elliptic curve \widehat{E}/K in some toric variety is *certifiably of genus 1* if $\text{Trop}(E)$ satisfies conditions (i)–(iii) from Theorem 6.3. Note that the cycle C in any such embedding satisfies $\ell_{\text{Trop}}(C) = -\text{val}(j_{\widehat{E}})$, by Theorem 6.3. Combining Theorems 6.2 and 6.3, we obtain the following result.

COROLLARY 6.6. *An elliptic curve \widehat{E}/K has multiplicative reduction if and only if it has a closed embedding in \mathbb{P}^2 which is certifiably of genus 1.*

6.7 Speyer's well-spacedness condition

In this section we explain how Speyer's well-spacedness condition [Spe14] follows from a more general result (possibly of independent interest) about the analytification of an elliptic curve \widehat{E}/K .

Let Σ be the minimal skeleton of \widehat{E} . For $P, Q \in \widehat{E}(K)$, define $i(P, Q) \in \mathbb{R}_{\geq 0} \cup \{\infty\}$ as follows:

$$i(P, Q) = \begin{cases} 0 & \text{if } \tau_{\Sigma}(P) \neq \tau_{\Sigma}(Q), \\ \text{dist}(P \vee Q, \Sigma) & \text{if } \tau_{\Sigma}(P) = \tau_{\Sigma}(Q), \end{cases}$$

where $\tau_{\Sigma}: \widehat{E}^{\text{an}} \rightarrow \Sigma$ is the retraction map, $P \vee Q$ is the first point where the geodesic paths from P to Σ and Q to Σ meet, and $\text{dist}(x, \Sigma)$ is the distance (in the natural metric on $\mathbf{H}(\widehat{E}^{\text{an}})$) from $x \in \widehat{E}^{\text{an}}$ to its retraction $\tau_{\Sigma}(x) \in \Sigma$. By convention we set $i(P, P) = +\infty$. Since translation by a point $P \in \widehat{E}(K)$ is an automorphism of \widehat{E} , it induces an isometry on $\mathbf{H}(\widehat{E}^{\text{an}})$; therefore $i(P, Q)$ only depends on the difference $P - Q$ in $\widehat{E}(K)$; that is, $i(P, Q) = \iota(P - Q)$ with $\iota(R) = i(R, 0)$.

The following lemma shows that $\|P, Q\| := \exp(-i(P, Q))$ is an *ultrametric* on $\widehat{E}(K)$.

LEMMA 6.8. (i) *For any points $P, Q, R \in \widehat{E}(K)$ we have $i(P, Q) \geq \min\{i(P, R), i(Q, R)\}$, with equality if $i(P, R) \neq i(Q, R)$.*

(ii) *If $m \in \mathbb{Z}$ is an integer such that $|m| = 1$ in K , then $i(mP, mQ) = i(P, Q)$ for any $P, Q \in \widehat{E}(K)$ such that $i(P, Q) > 0$.*

Proof. We begin by proving part (i). If either $i(P, R) = 0$ or $i(Q, R) = 0$, then the inequality is trivial. Moreover, by translation invariance of i we may assume $R = 0$. So we are reduced to showing that if $\iota(P) > 0$ and $\iota(Q) > 0$, then $\iota(P - Q) \geq \min\{\iota(P), \iota(Q)\}$.

Let \mathfrak{E} be the semistable formal model of \widehat{E} corresponding to the semistable vertex set $\{\tau_{\Sigma}(0)\}$ (see [BPR13, Remark 4.24]); note that $\Sigma = \Sigma(\widehat{E}, \{\tau_{\Sigma}(0)\})$. Then $\overline{\mathfrak{E}}$ is a nodal rational curve, and the smooth locus $\overline{\mathfrak{E}}^{\text{sm}}$ is a group scheme isomorphic to $\mathbf{G}_{m,k}$. The subset $\widehat{E}^1(K) := \{P \in \widehat{E}(K) : \iota(P) > 0\}$ is the formal fiber over the identity element of $\overline{\mathfrak{E}}^{\text{sm}}$, hence is a subgroup. In fact, $\widehat{E}^1(K)$ is isomorphic to the group $\mathfrak{m} = \{z \in K : |z| < 1\}$ with the law of composition given by a 1-parameter formal group law F over R [Sil09, Proposition VII.2.1], and the restriction of ι to $\widehat{E}^1(K)$ corresponds to the valuation on \mathfrak{m} under this identification. The desired inequality follows since a group law on \mathfrak{m} given by a power series with coefficients in R is obviously ultrametric.

In the situation of part (ii), as above we are reduced to showing that $\iota(mP) = \iota(P)$ when $\iota(P) > 0$. This is true because m is the coefficient of the linear term of the power series for multiplication by m under F [Sil09, Proposition IV.2.3], and all other terms have greater valuation. \square

Since $\iota(P) = \iota(-P)$, an equivalent formulation of Lemma 6.8(i) is that for any $P, Q \in \widehat{E}(K)$ we have $\iota(P + Q) \geq \min\{\iota(P), \iota(Q)\}$, with strict inequality if $\iota(P) \neq \iota(Q)$.

If f is a nonconstant rational function on \widehat{E} , define N_f to be the set of all $x \in \mathbf{H}(\widehat{E}^{\text{an}})$ such that $\log|f|$ is nonconstant in every open neighborhood of x . Equivalently, for $x \in \mathbf{H}(\widehat{E}^{\text{an}})$ let $T_x(f)$ be the (finite) set of tangent directions at x along which the derivative of $\log|f|$ is nonzero. Then N_f is the set of all $x \in \widehat{E}^{\text{an}}$ such that $T_x(f) \neq \emptyset$. By [BPR13, Theorem 5.15(1,2)], N_f is a union of finitely many edges of the minimal skeleton Γ_f of the curve obtained from \widehat{E}^{an} by removing all zeros and poles of f .

THEOREM 6.9. *Suppose that K has residue characteristic 0. Let f be a nonconstant rational function on \widehat{E} , and assume that there exists an $x \in N_f$ such that $\text{dist}(x, \Sigma) < \text{dist}(y, \Sigma)$ for all $y \in N_f$ with $y \neq x$. Assume also $\Sigma \cap N_f = \emptyset$. Then $|T_x(f)| \geq 3$.*

In other words, either the minimum distance from N_f to the skeleton is achieved at two distinct points, or else the minimum is achieved at a unique point at which $\log |f|$ has nonzero slope in at least three different tangent directions.

Proof. By the slope formula [BPR13, Theorem 5.15], the sum of the outgoing slopes of $\log |f|$ at x is 0, so $|T_x(f)| \geq 2$. For the sake of contradiction, assume $|T_x(f)| = 2$ and write $T_x(f) = \{v, v'\}$. Our hypotheses imply $x \notin \Sigma$ and that Σ lies in a single connected component of $\widehat{E}^{\text{an}} \setminus \{x\}$.

Let $B(x, v)$ (respectively, $B(x, v')$) be the open set consisting of all $z \in \widehat{E}^{\text{an}}$ lying in the tangent direction v (respectively, v'), so that $B(x, v)$ and $B(x, v')$ are connected components of $\widehat{E}^{\text{an}} \setminus \{x\}$ which are disjoint from Σ . Let D_v be the restriction of $\text{div}(f)$ to $B(x, v)$, and let $D_{v'}$ be the restriction of $\text{div}(f)$ to $B(x, v')$. By the slope formula, we have $m = \deg(D_v) = -\deg(D_{v'})$ for some nonzero integer m . Without loss of generality, we may assume $m > 0$. Let $\delta = \text{dist}(x, \Sigma) > 0$.

We claim that $\text{div}(f)$ can be written as

$$\text{div}(f) = m((P) - (Q)) + \sum_j ((A_j) - (B_j))$$

with $i(P, Q) = \delta$ and $i(A_j, B_j) > \delta$ for all j . Because we have assumed that K has residue characteristic 0, we have $|m| = 1$, and therefore $i(m(P), m(Q)) = i(P, Q) = \delta$ by Lemma 6.8(ii). Since $mP - mQ = \sum (B_j - A_j)$ in the group law on $\widehat{E}(K)$, we obtain a contradiction to Lemma 6.8(i).

To prove the claim, we use a trick due to Speyer. Suppose $D_v = (P_1) + \cdots + (P_r) - (Q_1) - \cdots - (Q_s)$ and $D_{v'} = (P'_1) + \cdots + (P'_{r'}) - (Q'_1) - \cdots - (Q'_{s'})$ with $r - s = m$ and $s' - r' = m$. Then

$$\begin{aligned} D_v + D_{v'} &= m((P_1) - (Q'_1)) + \sum_{j=1}^s ((P_j) - (Q_j)) + \sum_{j=s+1}^r ((P_j) - (P_1)) \\ &\quad + \sum_{j=1}^{r'} ((P'_j) - (Q'_j)) + \sum_{j=r'+1}^{s'} ((Q'_1) - (Q'_j)). \end{aligned}$$

Note that $i(P_1, Q'_1) = \delta$ but that $i(P_j, Q_j) > \delta$ for all $j = 1, \dots, s$; $i(P_j, P_1) > \delta$ for all $j = s + 1, \dots, r$; $i(P'_j, Q'_j) > \delta$ for all $j = 1, \dots, r'$; and $i(Q'_1, Q'_j) > \delta$ for all $j = r' + 1, \dots, s'$.

Let C_1, \dots, C_t be the connected components of N_f , labeled so that C_1 is the component containing x . By the slope formula [BPR13, Theorem 5.15], for each j the restriction D_j of $\text{div}(f)$ to C_j is a nonzero divisor of degree 0 and $\text{div}(f) = \sum_j D_j$. Moreover, if $A, B \in \widehat{E}(K) \cap C_j$, then the unique geodesic paths from A to Σ and B to Σ must pass through the unique point x_j of C_j closest to Σ , so for $j \geq 2$ we have $i(A, B) > \delta$. The claim now follows since $D_1 = D_v + D_{v'}$ can be written as above, and by what we have just said we can write each D_j for $j \geq 2$ as a sum of divisors of the form $(A) - (B)$ with $i(A, B) > \delta$. \square

In particular, we obtain the necessity of Speyer's well-spacedness condition for a genus 1 tropical curve to lift.

COROLLARY 6.10 (Speyer). *Suppose that K has residue characteristic 0. Let E be a dense open subset of an elliptic curve \widehat{E} over K with multiplicative reduction and let $\psi: E \hookrightarrow \mathbf{T}$ be a closed embedding of E in a torus \mathbf{T} . Assume that (i) every vertex of $\text{Trop}(E)$ is trivalent, (ii) every edge of $\text{Trop}(E)$ has multiplicity 1, and (iii) $\text{Trop}(E)$ contains a cycle Σ' which is contained in a hyperplane H . If W_H denotes the closure in $N_{\mathbb{R}}$ of the set of points of $\text{Trop}(E)$ not lying in H , then there is no single point of W_H which is closest to Σ' .*

In other words, ‘the minimum distance from points of $\text{Trop}(E)$ not lying in H to the cycle must be achieved twice’.

Proof. We may assume that $\psi: E \rightarrow \mathbf{T} \cong \mathbf{G}_m^n$ is given by (f_1, \dots, f_n) with $\log |f_n|$ equal to a constant c on Σ and that H is the hyperplane $x_n = c$. Let Γ be the minimal skeleton of E . By Corollary 5.28(i), we see that $\text{trop}: \Gamma \rightarrow \text{Trop}(E)$ is an isometry. Since $N_{f_n} \subset \Gamma$, the result now follows from Theorem 6.9. \square

Remark 6.11. Katz [Kat12] and Nishinou [Nis09] have recently obtained other kinds of generalizations of Speyer’s well-spacedness condition. Their generalized conditions apply to curves of higher genus.

7. A generalization of the Sturmfels–Tevelev multiplicity formula

As an illustration of the tools developed in this paper, we conclude with a generalization of the Sturmfels–Tevelev multiplicity formula [ST08, Theorem 1.1] to the nonconstant coefficient case (and also to nonsmooth points). A different proof is given in an appendix to [OP13].

Let $X \subset \mathbf{T}$ be a closed subvariety, that is, a reduced and irreducible closed subscheme. Let $\alpha: \mathbf{T} \rightarrow \mathbf{T}'$ be a homomorphism of tori that induces a generically finite map of degree δ from X to X' , where X' is the closure of $\alpha(X)$. Then, set theoretically, $\text{Trop}(X')$ is the image of $\text{Trop}(X)$ under the induced linear map $A: N_{\mathbb{R}} \rightarrow N'_{\mathbb{R}}$ [Tev07, Proposition 3]. The fundamental problem of tropical elimination theory is to determine the multiplicities on the maximal faces of $\text{Trop}(X')$ from those on the maximal faces of $\text{Trop}(X)$.

THEOREM 7.1. *Let $\alpha: \mathbf{T} \rightarrow \mathbf{T}'$ be a homomorphism of algebraic tori over K , and let X be a closed subvariety of \mathbf{T} . Let X' be the schematic image of X in \mathbf{T}' , let $f: X \rightarrow X'$ be the restriction of α to X , and let $F = \text{trop}(f): \text{Trop}(X) \rightarrow \text{Trop}(X')$ be the restriction of the linear map $A: N_{\mathbb{R}} \rightarrow N'_{\mathbb{R}}$ induced by α . Suppose that f is generically finite of degree δ . Then for any point $w' \in \text{Trop}(X') \cap N'_G$ such that $|F^{-1}(w')| < \infty$, we have*

$$m_{\text{Trop}}(w') = \frac{1}{\delta} \sum_{w \in F^{-1}(w')} \sum_{\bar{\mathcal{C}} \subset \bar{\mathfrak{X}}^w} \text{mult}_{\bar{\mathfrak{X}}^w}(\bar{\mathcal{C}})[\bar{\mathcal{C}} : \text{im}(\bar{\mathcal{C}})], \quad (7.1.1)$$

where the second sum runs over all irreducible components $\bar{\mathcal{C}}$ of $\bar{\mathfrak{X}}^w$ and where $\text{im}(\bar{\mathcal{C}})$ is the image of $\bar{\mathcal{C}}$ in $(\bar{\mathfrak{X}}')^{w'}$.

In order to use the projection formula (Proposition 3.25), we will need the following lemma.

LEMMA 7.2. *Let X and X' be integral finite-type K -schemes, and let $f: X \rightarrow X'$ be a generically finite dominant morphism of degree δ . Let $\mathcal{U}' \subset (X')^{\text{an}}$ be an analytic domain, and let $\mathcal{U} = (f^{\text{an}})^{-1}(\mathcal{U}')$. If $f^{\text{an}}|_{\mathcal{U}}: \mathcal{U} \rightarrow \mathcal{U}'$ is finite, then it has pure degree δ (Section 3.15).*

Proof. By [Har77, Exercise II.3.7], there is a dense open subscheme $U' \subset X'$ such that $U := f^{-1}(U') \rightarrow U'$ is finite. Shrinking U' if necessary, we assume that U' is smooth. By Proposition 3.24 the morphism $U^{\text{an}} \rightarrow (U')^{\text{an}}$ is pure of degree δ . Let \mathcal{U} and \mathcal{U}' be as in the statement of the lemma. By Proposition 3.23(ii), we may assume that $\mathcal{U} = \mathcal{M}(\mathcal{A})$ and $\mathcal{U}' = \mathcal{M}(\mathcal{A}')$ are affinoid. By [Con99, Lemma A.1.2(2)], \mathcal{U} and \mathcal{U}' are equidimensional of the same dimension as X and X' , respectively. Therefore $\mathcal{U}' \cap (X' \setminus U')^{\text{an}}$ is nowhere dense in \mathcal{U}' . If $\mathcal{V}' = \mathcal{M}(\mathcal{B}')$ is any connected affinoid subdomain of $\mathcal{U}' \cap (U')^{\text{an}}$, then \mathcal{B}' is a domain because \mathcal{V}' is smooth, and $\mathcal{V} := (f^{\text{an}})^{-1}(\mathcal{V}') \rightarrow \mathcal{V}'$ has (pure) degree δ because $\mathcal{V}' \subset (U')^{\text{an}}$. Since $\mathcal{U}' \cap (X' \setminus U')^{\text{an}}$ is

nowhere dense in \mathcal{U}' , we can choose \mathcal{V}' such that $\text{Spec}(\mathcal{B}') \rightarrow \text{Spec}(\mathcal{A}')$ takes the generic point of $\text{Spec}(\mathcal{B}')$ to any given generic point of $\text{Spec}(\mathcal{A}')$. Hence $\mathcal{U} \rightarrow \mathcal{U}'$ has pure degree δ . \square

Proof of Theorem 7.1. Let $w' \in \text{Trop}(X') \cap N'_G$ be a point with finite preimage under F . Let

$$\mathcal{X}^{w'} = \text{trop}^{-1}(F^{-1}(w')) \cap X^{\text{an}} = \coprod_{w \in F^{-1}(w')} \mathcal{X}^w.$$

This is an affinoid domain in X^{an} because it is a closed subspace of the affinoid $\text{trop}^{-1}(F^{-1}(w')) = \coprod_{w \in F^{-1}(w')} \mathcal{U}^w$. We claim that $\mathcal{X}^{w'} \rightarrow (\mathcal{X}')^{w'}$ is a finite morphism. It suffices to show that the composite $\mathcal{X}^{w'} \rightarrow \mathcal{U}^{w'}$ is a finite morphism, where $\mathcal{U}^{w'} = \text{trop}^{-1}(w') \subset (\mathbf{T}')^{\text{an}}$. This follows exactly as in the proof of Theorem 4.30: Since $F^{-1}(w')$ is bounded, there is an affinoid domain of \mathbf{T}^{an} contained in $(\alpha^{\text{an}})^{-1}(\mathcal{U}^{w'})$ and containing $\mathcal{X}^{w'}$ in its relative interior. This means that the morphism $\mathcal{X}^{w'} \rightarrow (\mathcal{X}')^{w'}$ is proper, thus finite because both spaces are affinoid. Hence by Lemma 7.2 the morphism $\mathcal{X}^{w'} \rightarrow (\mathcal{X}')^{w'}$ has pure degree δ . Let $\mathfrak{X}^{w'} := \coprod_{w \in F^{-1}(w')} \mathfrak{X}^w$. The generic fiber of $\mathfrak{X}^{w'}$ is $\mathcal{X}^{w'}$, and the natural morphism $\mathfrak{X}^{w'} \rightarrow (\mathfrak{X}')^{w'}$ is finite by Proposition 3.8(ii) and takes generic points to generic points by Proposition 3.10. By Proposition 3.25 the induced morphism $\overline{\mathfrak{X}}^{w'} \rightarrow (\overline{\mathfrak{X}}')^{w'}$ has pure degree δ , so summing (3.18.1) over all irreducible components $\overline{\mathfrak{C}}'$ of $(\overline{\mathfrak{X}}')^{w'}$ yields

$$\delta \cdot m_{\text{Trop}}(w') = \delta \sum_{\overline{\mathfrak{C}}' \subset (\overline{\mathfrak{X}}')^{w'}} \text{mult}_{(\overline{\mathfrak{X}}')^{w'}}(\overline{\mathfrak{C}}') = \sum_{\overline{\mathfrak{C}} \subset \overline{\mathfrak{X}}^{w'}} \text{mult}_{\overline{\mathfrak{X}}^{w'}}(\overline{\mathfrak{C}}) [\overline{\mathfrak{C}} : \text{im}(\overline{\mathfrak{C}})].$$

Since $\mathfrak{X}^{w'} = \coprod_{w \in F^{-1}(w')} \mathfrak{X}^w$, this is the desired multiplicity formula. \square

As a consequence of Theorem 7.1, we obtain the following result.

COROLLARY 7.3. *Let $\alpha: \mathbf{T} \rightarrow \mathbf{T}'$ be a homomorphism of algebraic tori over K , and let $A = \text{trop}(\alpha): N_{\mathbb{R}} \rightarrow N'_{\mathbb{R}}$ be the natural linear map. Let X be a closed subvariety of \mathbf{T} , and suppose that α induces a generically finite morphism of degree δ from X onto its schematic image X' in \mathbf{T}' . After subdividing, we may assume that A maps each face of $\text{Trop}(X)$ onto a face of $\text{Trop}(X')$. Let σ' be a maximal face of $\text{Trop}(X')$. Then*

$$m(\sigma') = \frac{1}{\delta} \sum_{A(\sigma) = \sigma'} m(\sigma) \cdot [N'_{\sigma'} : A(N_{\sigma})].$$

(Here N_{σ} and $N'_{\sigma'}$ are the sublattices of N and N' parallel to σ and σ' , respectively.)

Proof. If w' is a smooth point of $\text{Trop}(X')$ and w is a smooth point of $\text{Trop}(X)$ with $A(w) = w'$, then $(\overline{\mathfrak{X}}')^{w'} \cong Y' \times T'(w')$ and $\overline{\mathfrak{X}}^w \cong Y \times T(w)$, where Y and Y' are 0-dimensional schemes of length $m_{\text{Trop}}(w)$ and $m_{\text{Trop}}(w')$, respectively, and $T(w)$ and $T'(w')$ are algebraic tori of dimension $\dim(X) = \dim(X')$ (cf. Remark 4.27). Moreover, α induces a finite homomorphism $T(w) \rightarrow T'(w')$ of degree $[N'_{\sigma'} : A(N_{\sigma})]$. In this situation, the quantity $[\overline{\mathfrak{C}} : \text{im}(\overline{\mathfrak{C}})]$ appearing in (7.1.1) is equal to $[T(w) : T'(w')] = [N'_{\sigma'} : A(N_{\sigma})]$, and $m_{\text{Trop}}(w) = \sum_{\overline{\mathfrak{C}} \subset \overline{\mathfrak{X}}^w} \text{mult}_{\overline{\mathfrak{X}}^w}(\overline{\mathfrak{C}})$, so we are reduced to Theorem 7.1. \square

Remark 7.4. The original Sturmfels–Tevelev multiplicity formula is the special case of Corollary 7.3 in which $K = k\{\{T\}\}$ and X is defined over k .

ACKNOWLEDGEMENTS

The authors would like to express their thanks to María Angélica Cueto, Eric Katz, Brian Osserman, David Speyer, and Josephine Yu for helpful discussions, and to Bernd Sturmfels for his interest and encouragement. Special thanks are due to Walter Gubler for some illuminating conversations which helped shape the direction of this work. The authors also thank Melanie Dunn for computing several nice examples of tropicalizations, and the anonymous referees for many helpful comments.

REFERENCES

- ABBR15a O. Amini, M. Baker, E. Brugallé, and J. Rabinoff, *Lifting harmonic morphisms I: Metrized complexes and Berkovich skeleta*, Res. Math. Sci. **2** (2015), no. 7; <http://dx.doi.org/10.1186/s40687-014-0019-0>.
- ABBR15b ———, *Lifting harmonic morphisms II: Tropical curves and metrized complexes*, Algebra Number Theory **9** (2015), no. 2, 267–315; <http://dx.doi.org/10.2140/ant.2015.9.267>.
- Bak93 H. F. Baker, *Examples of applications of Newton's polygon to the theory of singular points of algebraic functions*, Trans. Cambridge Philos. Soc. **15** (1893), 403–450.
- Bak08 M. Baker, *An introduction to Berkovich analytic spaces and non-Archimedean potential theory on curves*, *p-Adic Geometry*, Univ. Lecture Ser., vol. 45 (Amer. Math. Soc., Providence, RI, 2008), 123–174; <http://dx.doi.org/10.1090/ulect/045/04>.
- Bee09 P. Beelen, *A generalization of Baker's theorem*, Finite Fields Appl. **15** (2009), no. 5, 558–568; <http://dx.doi.org/10.1016/j.ffa.2009.04.003>.
- Ber90 V. G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Math. Surveys and Monogr., vol. 33 (Amer. Math. Soc., Providence, RI, 1990).
- Ber93 ———, *Étale cohomology for non-Archimedean analytic spaces*, Publ. Math. Inst. Hautes Études Sci. (1993), no. 78, 5–161; <http://dx.doi.org/10.1007/BF02712916>.
- Ber99 ———, *Smooth p-adic analytic spaces are locally contractible*, Invent. Math. **137** (1999), no. 1, 1–84; <http://dx.doi.org/10.1007/s002220050323>.
- Ber04 ———, *Smooth p-adic analytic spaces are locally contractible II*, in *Geometric Aspects of Dwork Theory* (Walter de Gruyter GmbH & Co. KG, Berlin, 2004), 293–370.
- BG84 R. Bieri and J. R. J. Groves, *The geometry of the set of characters induced by valuations*, J. reine angew. Math. **347** (1984), 168–195; <http://dx.doi.org/10.1515/crll.1984.347.168>.
- BGR84 S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis. A systematic approach to rigid analytic geometry*, Grundlehren Math. Wiss. vol. 261 (Springer-Verlag, Berlin, 1984); <http://dx.doi.org/10.1007/978-3-642-52229-1>.
- BL85 S. Bosch and W. Lütkebohmert, *Stable reduction and uniformization of abelian varieties I*, Math. Ann. **270** (1985), no. 3, 349–379; <http://dx.doi.org/10.1007/BF01473432>.
- BL93 ———, *Formal and rigid geometry I: Rigid spaces*, Math. Ann. **295** (1993), no. 2, 291–317; <http://dx.doi.org/10.1007/BF01444889>.
- Bos14 S. Bosch, *Lectures on formal and rigid geometry*, Lecture Notes in Math., vol. 2105 (Springer, Cham, 2014); <http://dx.doi.org/10.1007/978-3-319-04417-0>.
- BPR13 M. Baker, S. Payne, and J. Rabinoff, *On the structure of non-Archimedean analytic curves*, in *Tropical and non-Archimedean geometry*, Contemp. Math. vol. 605, Amer. Math. Soc. (Providence, RI, 2013), 93–121; <http://dx.doi.org/10.1090/conm/605/12113>.
- BR10 M. Baker and R. Rumely, *Potential theory and dynamics on the Berkovich projective line*, Math. Surveys and Monogr. vol. 159 (Amer. Math. Soc., Providence, RI, 2010); <http://dx.doi.org/10.1090/surv/159>.

- CHW14 M. A. Cueto, M. Hübich, and A. Werner, *Faithful tropicalization of the Grassmannian of planes*, Math. Ann. **360** (2014), no. 1-2, 391–437; <http://dx.doi.org/10.1007/s00208-014-1037-3>.
- Con99 B. Conrad, *Irreducible components of rigid spaces*, Ann. Inst. Fourier (Grenoble) **49** (1999), no. 2, 473–541; <http://dx.doi.org/10.5802/aif.1681>.
- Con08 ———, *Several approaches to non-Archimedean geometry*, in *p-Adic Geometry*, Univ. Lecture Ser., vol. 45, Amer. Math. Soc. (Providence, RI, 2008), 9–63; <http://dx.doi.org/10.1090/ulect/045/02>.
- CS13 M. Chan and B. Sturmfels, *Elliptic curves in honeycomb form*, in *Algebraic and Combinatorial Aspects of Tropical Geometry*, Contemp. Math., vol. 589 (Amer. Math. Soc., Providence, RI, 2013), 87–107; <http://dx.doi.org/10.1090/comm/589/11743>.
- DP14 J. Draisma and E. Postingshel, *Faithful tropicalisation and torus actions*, [arXiv:1404.4715](https://arxiv.org/abs/1404.4715).
- Duc09 A. Ducros, *Les espaces de Berkovich sont excellents*, Ann. Inst. Fourier (Grenoble) **59** (2009), no. 4, 1443–1552; <http://dx.doi.org/10.5802/aif.2470>.
- EKL06 M. Einsiedler, M. Kapranov, and D. Lind, *Non-Archimedean amoebas and tropical varieties*, J. Reine Angew. Math. **601** (2006), 139–157; <http://dx.doi.org/10.1515/CRELLE.2006.097>.
- FM86 J. Fresnel and M. Matignon, *Sur les espaces analytiques quasi-compacts de dimension 1 sur un corps valué complet ultramétrique*, Ann. Mat. Pura Appl. (4) **145** (1986), 159–210; <http://dx.doi.org/10.1007/BF01790541>.
- Ful93 W. Fulton, *Introduction to toric varieties*, Ann. of Math. Stud., vol. 131 (Princeton University Press, Princeton, NJ, 1993).
- GD66 A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III*, Publ. Math. Inst. Hautes Études Sci. (1966), no. 28, 5–248; <http://dx.doi.org/10.1007/BF02684343>, http://www.numdam.org/item?id=PMIHES_1966__28__5_0.
- GRW14 W. Gubler, J. Rabinoff, and A. Werner, *Skeletons and tropicalizations*, [arXiv:1404.7044](https://arxiv.org/abs/1404.7044).
- Gub98 W. Gubler, *Local heights of subvarieties over non-Archimedean fields*, J. reine angew. Math. **498** (1998), 61–113; <http://dx.doi.org/10.1515/crll.1998.054>.
- Gub07a ———, *The Bogomolov conjecture for totally degenerate abelian varieties*, Invent. Math. **169** (2007), no. 2, 377–400; <http://dx.doi.org/10.1007/s00222-007-0049-y>.
- Gub07b ———, *Tropical varieties for non-Archimedean analytic spaces*, Invent. Math. **169** (2007), no. 2, 321–376; <http://dx.doi.org/10.1007/s00222-007-0048-z>.
- Gub13 ———, *A guide to tropicalizations*, in *Algebraic and Combinatorial Aspects of Tropical Geometry*, Contemp. Math., vol. 589 (Amer. Math. Soc., Providence, RI, 2013), 125–189; <http://dx.doi.org/10.1090/comm/589/11745>.
- Har77 R. Hartshorne, *Algebraic geometry*, Grad. Texts in Math., vol. 52 (Springer-Verlag, New York – Heidelberg, 1977); <http://dx.doi.org/10.1007/978-1-4757-3849-0>.
- HK12 D. Helm and E. Katz, *Monodromy filtrations and the topology of tropical varieties*, Canad. J. Math. **64** (2012), no. 4, 845–868; <http://dx.doi.org/10.4153/CJM-2011-067-9>.
- Kat12 E. Katz, *Lifting tropical curves in space and linear systems on graphs*, Adv. Math. **230** (2012), no. 3, 853–875; <http://dx.doi.org/10.1016/j.aim.2012.03.017>.
- KMM08 E. Katz, H. Markwig, and T. Markwig, *The j -invariant of a plane tropical cubic*, J. Algebra **320** (2008), no. 10, 3832–3848; <http://dx.doi.org/10.1016/j.jalgebra.2008.08.018>.
- KMM09 ———, *The tropical j -invariant*, LMS J. Comput. Math. **12** (2009), 275–294; <http://dx.doi.org/10.1112/S1461157000001522>.
- MFK94 D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, third ed., Ergeb. Math. Grenzgeb. (2), vol. 34 (Springer-Verlag, Berlin, 1994); <http://dx.doi.org/10.1007/978-3-642-57916-5>.

- MS15 D. Maclagan and B. Sturmfels, *Introduction to tropical geometry*, Grad. Stud. Math., vol. 161 (Amer. Math. Soc., Providence, RI, 2015).
- Nis09 T. Nishinou, *Correspondence theorems for tropical curves*, [arXiv:0912.5090](https://arxiv.org/abs/0912.5090).
- OP13 B. Osserman and S. Payne, *Lifting tropical intersections*, Doc. Math. **18** (2013), 121–175.
- Pap05 A. Papadopoulos, *Metric spaces, convexity and nonpositive curvature*, IRMA Lect. Math. Theor. Phys., vol. 6, (European Mathematical Society, Zürich, 2005); <http://dx.doi.org/10.4171/132>.
- Pay09a S. Payne, *Analytification is the limit of all tropicalizations*, Math. Res. Lett. **16** (2009), no. 3, 543–556; <http://dx.doi.org/10.4310/MRL.2009.v16.n3.a13>.
- Pay09b ———, *Fibers of tropicalization*, Math. Z. **262** (2009), no. 2, 301–311; <http://dx.doi.org/10.1007/s00209-008-0374-x>.
- Pay12 ———, *Erratum to “Fibers of tropicalization”*, Math. Z. **272** (2012), no. 3-4, 1403–1406; <http://dx.doi.org/10.1007/s00209-012-1080-2>.
- Rab12 J. Rabinoff, *Tropical analytic geometry, Newton polygons, and tropical intersections*, Adv. Math. **229** (2012), no. 6, 3192–3255; <http://dx.doi.org/10.1016/j.aim.2012.02.003>.
- RG71 M. Raynaud and L. Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), 1–89; <http://dx.doi.org/10.1007/BF01390094>.
- Rob00 A. M. Robert, *A course in p-adic analysis*, Grad. Texts in Math., vol. 198 (Springer-Verlag, New York, 2000); <http://dx.doi.org/10.1007/978-1-4757-3254-2>.
- Sil09 J. H. Silverman, *The arithmetic of elliptic curves*, second ed., Grad. Texts in Math., vol. 106 (Springer, Dordrecht, 2009), <http://dx.doi.org/10.1007/978-0-387-09494-6>.
- Spe05 D. E. Speyer, *Tropical geometry*, Ph.D. thesis, University of California, Berkeley, 2005, available at <http://www-personal.umich.edu/~speyer/thesis.pdf>.
- Spe14 ———, *Parameterizing tropical curves I: Curves of genus zero and one*, Algebra Number Theory **8** (2014), no. 4, 963–998; <http://dx.doi.org/10.2140/ant.2014.8.963>.
- ST08 B. Sturmfels and J. Tevelev, *Elimination theory for tropical varieties*, Math. Res. Lett. **15** (2008), no. 3, 543–562; <http://dx.doi.org/10.4310/MRL.2008.v15.n3.a14>.
- Tev07 J. Tevelev, *Compactifications of subvarieties of tori*, Amer. J. Math. **129** (2007), no. 4, 1087–1104; <http://dx.doi.org/10.1353/ajm.2007.0029>.
- Tho90 A. Thorup, *Rational equivalence theory on arbitrary Noetherian schemes*, Enumerative Geometry (Sitges, 1987), Lecture Notes in Math., vol. 1436 (Springer, Berlin, 1990), 256–297; <http://dx.doi.org/10.1007/BFb0084049>.
- Thu05 A. Thuillier, *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne. Applications à la théorie d’Arakelov*, Ph.D. thesis, University of Rennes, 2005, available at <http://tel.ccsd.cnrs.fr/documents/archives0/00/01/09/90/index.html>.
- Vaq00 M. Vaquié, *Valuations, Resolution of Singularities* (Obergrugl, 1997), Progr. Math., vol. 181 (Birkhäuser, Basel, 2000), 539–590; http://dx.doi.org/10.1007/978-3-0348-8399-3_21.

Matthew Baker mbaker@math.gatech.edu

School of Mathematics, Georgia Institute of Technology, Atlanta GA 30332-0160, USA

Sam Payne sam.payne@yale.edu

Mathematics Department, Yale University, New Haven, CT 06511, USA

Joseph Rabinoff rabinoff@math.gatech.edu

Department of Mathematics, Harvard University, Cambridge, MA 02138, USA

Current address: School of Mathematics, Georgia Institute of Technology, Atlanta GA 30332-0160, USA