

ANALYTIFICATION AND TROPICALIZATION OVER NON-ARCHIMEDEAN FIELDS

ANNETTE WERNER

Abstract:

We provide a survey of recent progress on the interplay between tropical geometry and non-archimedean analytic geometry in the sense of Berkovich. After briefly discussing results by Baker, Payne and Rabinoff [BPR11] in the case of curves, we explain the comparison between the tropical Grassmannian of planes and the analytic Grassmannian obtained in joint work with Maria Angelica Cueto and Mathias Häbich [GRW14]. We also give an overview of the general higher-dimensional approach developed in joint work with Walter Gubler and Joseph Rabinoff [GRW14]. We explain the construction of generalized skeleta in [GRW14] which are polyhedral substructures of Berkovich spaces lending themselves to comparison with tropicalizations. We discuss the slope formula for the valuation of rational functions and explain two results on the comparison between polyhedral substructures of Berkovich spaces and tropicalizations.

2010 MSC: 14G22, 14T05

1. INTRODUCTION

Tropical varieties are polyhedral images of varieties over non-archimedean fields. They are obtained by applying the valuation map to a set of toric coordinates. From the very beginning, analytic geometry was present in the systematic study of tropical varieties, e.g. in [EKL06] where rigid analytic varieties are used. The theory of Berkovich spaces which leads to spaces with nicer topological properties than rigid varieties is even better suited to study tropicalizations.

A result of Payne [Pay09] states that the Berkovich space associated to an algebraic variety is homeomorphic to the inverse limit of all tropicalizations in toric varieties. However, individual tropicalizations may fail to capture topological features of an analytic space. The present paper is an overview of recent results on the relationship between analytic spaces and tropicalizations. In particular, we address the question of whether a given tropicalization is contained in a Berkovich space as a combinatorial substructure.

A novel feature of Berkovich spaces compared to rigid analytic varieties is that they contain interesting piecewise linear combinatorial structures. In fact, Berkovich curves are, very roughly speaking, generalized graphs, where infinite ramifications along a dense set of points is allowed, see [Ber90], chapter 4, and [BaRu10].

In higher dimensions, the structure of Berkovich analytic spaces is more involved, but still they often contain piecewise linear substructures as deformation retracts. This was a crucial tool in Berkovich’s proof of local contractibility for smooth analytic spaces, see [Ber99] and [Ber04]. Berkovich constructs these piecewise linear substructures as so-called skeleta of suitable models (or fibration of models). These skeleta basically capture the incidence structure of the irreducible components in the special fiber.

In dimension one, Baker, Payne and Rabinoff studied the relationship between tropicalizations and subgraphs of Berkovich curves in [BPR11]. Their results show that every finite subgraph of a Berkovich curve admits a faithful, i.e. homeomorphic and isometric tropicalization. They also prove that every tropicalization with tropical multiplicity one everywhere is isometric to a finite subgraph of the Berkovich curve.

As a first higher dimensional example, the Grassmannian of planes was studied in [CHW14]. The tropical Grassmannian of planes has an interesting combinatorial structure and is a moduli space of phylogenetic trees. It is shown in [CHW14] that it is homeomorphic to a closed subset of the Berkovich analytic Grassmannian.

In [GRW14], the higher dimensional situation is analyzed from a general point of view. This approach is based on a generalized notion of a Berkovich skeleton which is associated to the datum of a semistable model plus a horizontal divisor. This naturally leads to unbounded skeleta and generalizes a well-known construction on curves, see [Ty012] and [BPR11].

For every rational function f with support in the fixed horizontal divisor it is shown in [GRW14] that the “tropicalization” $\log |f|$ factors through a piecewise linear function on the generalized skeleton. This function satisfies a slope formula which is a kind of balancing condition around any 1-codimensional polyhedral face. Moreover, for every generalized skeleton there exists a faithful tropicalization – where faithful in higher dimension refers to the preservation of the integral affine structures. In dimension one, this can be expressed via metrics. It is also proven that tropicalizations with tropical multiplicity one everywhere admit sections of the tropicalization map, which generalizes the above-mentioned result in [BPR11] on curves.

The paper is organized as follows. In section 2 we collect basic facts on Berkovich spaces and tropicalizations, giving references to the literature for proofs and more details. In section 3 we briefly recall some of the results by Baker, Payne and Rabinoff [BPR11] on curves. Section 4 starts with the definition and basic properties of the tropical Grassmannian. Theorem 4.1 claims the existence of a continuous section to the tropicalization map on the projective tropical Grassmannian. We give a sketch of the proof for the dense torus orbit, where some constructions are easier to explain than in the general case. In section 5 we explain the construction of generalized skeleta from [GRW14], and in section 6 we investigate $\log |f|$ for rational functions f . In particular, Theorem 6.5 states the slope formula. Section 7 explains the faithful tropicalization results in higher dimension.

Acknowledgements: The author would like to thank Walter Gubler for his helpful comments. Collaboration on the paper [GRW14] started during an open problem session at the Simons symposium on *Non-archimedean and Tropical Geometry* in 2013. The author is very grateful to the Simons foundation and to the organizers Matt Baker and Sam Payne for bringing us together at this exciting meeting.

2. BERKOVICH SPACES AND TROPICALIZATIONS

2.1. Notation and conventions. A non-archimedean field is a field with a non-archimedean absolute value. Our ground field is a non-archimedean field which is complete with respect to its absolute value. Examples are the field \mathbb{Q}_p , which is the completion of \mathbb{Q} after the p -adic absolute value, finite extensions of \mathbb{Q}_p and the p -adic cousin \mathbb{C}_p of the complex numbers which is defined as the completion of the algebraic closure of \mathbb{Q}_p . The field of formal Laurent series $k((t))$ over an arbitrary base field k is another example. Besides, we can endow any field k with the trivial absolute value (which is one on all non-zero elements). A nice feature of Berkovich's general approach is that it also gives an interesting theory in the case of a trivially valued field. The reason behind this is that Berkovich geometry encompasses also points with values in transcendental field extensions – and those may well carry interesting non-trivial valuations.

Let K be a complete non-archimedean field. We write $K^\circ = \{x \in K : |x| \leq 1\}$ for the ring of integers in K , and $K^{\circ\circ} = \{x \in K : |x| < 1\}$ for the valuation ideal. The quotient $\tilde{K} = K^\circ / K^{\circ\circ}$ is the residue field of K . The valuation on K^* associated to the absolute value is given by $v(x) = -\log |x|$. By $\Gamma = v(K^*) \subset \mathbb{R}$ we denote the value group.

A variety over K is an irreducible, reduced and separated scheme of finite type over K .

2.2. Berkovich spaces. Let us briefly recall some basic results about Berkovich spaces. This theory was developed in the ground-breaking treatise [Ber90]. The survey papers [Co08], [Du07] and [Tem] provide additional information. For background information on non-archimedean fields and Banach algebras and an account of rigid analytic geometry see [BGR84] and [Bo14].

For $n \in \mathbb{N}$ and every n -Tupel $r = (r_1, \dots, r_n)$ of positive real numbers we write $x = (x_1, \dots, x_n)$ and define the associated *Tate algebra* as

$$K\{r^{-1}x\} = \left\{ \sum_{I=(i_1, \dots, i_n) \in \mathbb{N}_0^n} a_I x^I : |a_I| r^I \rightarrow 0 \text{ as } |I| \rightarrow \infty \right\}.$$

Here we put $x = (x_1, \dots, x_n)$ and write $x^I = x^{i_1} \dots x^{i_n}$. Moreover, for $I = (i_1, \dots, i_n)$ we write $|I| = i_1 + \dots + i_n$.

A Banach algebra A is called *K -affinoid*, if there exists a surjective K -algebra homomorphism $\alpha : K\{r^{-1}x\} \rightarrow A$ for some n and r such that the Banach norm on A is equivalent the quotient seminorm induced by α . If there exists such an α for $r = (1, \dots, 1)$, then A

is called *strictly K -affinoid*. In rigid analytic geometry only strictly K -affinoid algebras are considered (and called affinoid algebras).

The *Berkovich spectrum* $\mathcal{M}(A)$ of an affinoid algebra is defined as the set of all non-zero, bounded (by the Banach norm), multiplicative seminorms on A . It is endowed with the coarsest topology such that all evaluation maps on functions in A are continuous.

If x is a seminorm on an algebra A , and $f \in A$, we follow the usual notational convention and write $|f(x)|$ for $x(f)$, i.e. for the real number which we get by evaluating the seminorm on f .

The *Shilov boundary* of a Berkovich spectrum $\mathcal{M}(A)$ of a K -affinoid algebra is the smallest subset Γ (with respect to inclusion) such that for every $f \in A$ the evaluation map $\mathcal{M}(A) \rightarrow \mathbb{R}_{\geq 0}$ given by $x \mapsto |f(x)|$ attains its maximum on Γ . Hence for every $f \in A$ there exists a point z in the Shilov boundary such that $|f(z)| \geq |f(x)|$ for all $x \in \mathcal{M}(A)$. The Shilov boundary exists and is a finite set by [Ber90], Corollary 2.4.5.

The Berkovich spectrum of a K -affinoid algebra carries a sheaf of analytic functions which we will not define here. Some care is needed since, very roughly speaking, not all coverings are suitable for glueing analytic functions. Analytic spaces over K are ringed spaces which are locally modelled on Berkovich spectra of affinoid algebras.

There is a *GAGA-functor*, associating to every K -scheme X locally of finite type an analytic space X^{an} over K . It has the property that X is connected, separated over K or proper over K , respectively, if and only if the topological space X^{an} is arcwise connected, Hausdorff, or compact, respectively. If $X = \text{Spec}R$ is affine, the topological space X^{an} can be identified with the set of all multiplicative seminorms on the coordinate ring R extending the absolute value on K . This space is endowed with the coarsest topology such that for all $f \in R$ the evaluation map on f is continuous.

With the help of the GAGA-functor we associate analytic spaces to algebraic varieties. Another way of obtaining analytic spaces is via *admissible formal schemes*. An admissible formal scheme over K° is, roughly speaking, a formal scheme over K° such that the formal affine building blocks are given by K° -flat algebras of the form $K^\circ\{x_1, \dots, x_n\}/\mathfrak{a}$ for a finitely generated ideal \mathfrak{a} in the ring $K^\circ\{x_1, \dots, x_n\} = \{\sum_{I \in \mathbb{N}_0^n} a_I x^I : a_I \in K^\circ \text{ and } |a_I| \rightarrow 0 \text{ as } |I| \rightarrow \infty\}$. For a precise definition see [Bo14] or [Co08].

An admissible formal scheme \mathcal{X} has a natural *analytic generic fiber* \mathcal{X}_η . On the formal affine building block given by $K^\circ\{x_1, \dots, x_n\}/\mathfrak{a}$ the analytic generic fiber is the Berkovich spectrum of $K\{x_1, \dots, x_n\}/\mathfrak{a}K\{x_1, \dots, x_n\}$.

If we start with a K° -scheme \mathcal{X} , which is separated, flat and of finite presentation, its completion after any element $\pi \in K^{\circ\circ} \setminus \{0\}$ is an admissible formal scheme \mathcal{X} . This has an analytic generic fiber \mathcal{X}_η . On the other hand, we can apply the GAGA-functor to the algebraic generic fiber $X = \mathcal{X} \otimes_{K^\circ} K$ and get another analytic space X^{an} . They are connected by a morphism $\mathcal{X}_\eta \hookrightarrow X^{\text{an}}$, which is an isomorphism if \mathcal{X} is proper over K° .

In the basic example $\mathcal{X} = \text{Spec}R[x]$ we get the natural inclusion $\mathcal{M}(K^\circ\{x\}) \hookrightarrow (\mathbb{A}_K^1)^{\text{an}}$, whose image is the set of all multiplicative seminorms on the polynomial ring $K[x]$ which extend the absolute value on K and whose value on x is bounded by 1.

2.3. Tropicalization. We start by considering a split torus $T = \text{Spec}K[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$. Evaluating seminorms on characters we get a map $\text{trop} : T^{\text{an}} \rightarrow \mathbb{R}^n$ on the associated Berkovich space, which is given by

$$\text{trop}(p) = (-\log |x_1(p)|, \dots, -\log |x_n(p)|).$$

Here we are using the valuation map to define tropicalizations. In some papers, tropicalizations are defined via the negative valuation map, i.e. the logarithmic absolute value of the coordinates.

If X is a variety over K together with a closed embedding $\varphi : X \hookrightarrow T$, we consider the composition

$$(2.1) \quad \text{trop}_\varphi : X^{\text{an}} \xrightarrow{\varphi^{\text{an}}} T^{\text{an}} \xrightarrow{\text{trop}} \mathbb{R}^n$$

of the tropicalization map with the embedding φ . Note that the map trop_φ is continuous. Its image $\text{Trop}_\varphi(X) = \text{trop}_\varphi(X^{\text{an}})$ is the support of a polyhedral complex Σ in \mathbb{R}^n . This complex is integral Γ -affine in the sense of section 5.1. It has pure dimension $d = \dim(X)$. Moreover, the polyhedral complex $\text{Trop}_\varphi(X)$ can be enriched with weights on the maximal faces such that it satisfies a balancing condition. For a nice introductory text on tropical geometry see the textbook [Ma-St]. The survey paper [Gu13] is also very useful.

Note that for all $\omega \in \text{Trop}_\varphi(X)$ the preimage $\text{trop}_\varphi^{-1}(\{\omega\})$ is homeomorphic to the Berkovich spectrum of an affinoid algebra, see [Gu07], Proposition 4.1.

For every point $\omega \in \text{Trop}_\varphi(X)$ there is an associated *initial degeneration*, which is the special fiber of a model of X over the valuation ring in a suitable non-archimedean extension field of K . The geometric number of irreducible components of this special fiber is the *tropical multiplicity* of ω . There is a balancing formula involving tropical multiplicities on all maximal-dimensional faces of $\text{Trop}_\varphi(X)$ around a fixed codimension one face, see [Ma-St], chapter 3.4.

Let \overline{K} be the algebraic closure of K . It can be endowed with an absolute value extending the one on K . By means of the coordinates x_1, \dots, x_n we can identify $T(\overline{K}) = \overline{K}^n$. Then we can define a natural tropicalization map $\text{trop} : T(\overline{K}) \rightarrow \mathbb{R}^n$ by $\text{trop}(t_1, \dots, t_n) = (-\log |t_1|, \dots, -\log |t_n|)$. Note that every point $t = (t_1, \dots, t_n)$ in $T(\overline{K})$ gives rise to the multiplicative seminorm $f \mapsto |f(t)|$ on the coordinate ring $K[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$, which is an element in T^{an} . Hence the tropicalization map on $T(\overline{K})$ is induced by the map trop on T^{an} .

We could define the tropicalization $\text{Trop}_\varphi(X)$ without recourse to Berkovich spaces. If the absolute value on K is non-trivial, the tropicalization $\text{Trop}_\varphi(X)$ is equal to the closure

of the image of the map

$$X(\overline{K}) \xrightarrow{\varphi} T(\overline{K}) \xrightarrow{\text{trop}} \mathbb{R}^n.$$

Considering a tropical variety as the image of an analytic space makes some topological considerations easier. For example, in this way it is evident that the tropicalization of a connected variety is connected since it is a continuous image of a connected space.

Let Y be a toric variety associated to the fan Δ in $N_{\mathbb{R}}$, where N is the cocharacter group of the dense torus T . Then Δ defines a natural compactification $N_{\mathbb{R}}^{\Delta}$ of the space $N_{\mathbb{R}}$. Roughly speaking, we compactify cones by dual spaces. For details see [Pay09], section 3. There is also a natural tropicalization map

$$\text{trop} : Y^{\text{an}} \rightarrow N_{\mathbb{R}}^{\Delta},$$

which extends the tropicalization map on the dense torus. As an example, we consider $Y = \mathbb{P}_K^n$ with its dense torus $\mathbb{G}_{m,K}^n$ and $N = \mathbb{Z}^n$. Put $\overline{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$ and endow it with the natural topology such that half-open intervals $]a, \infty]$ form a basis of open neighbourhoods of ∞ . Then the associated compactification of $N_{\mathbb{R}} = \mathbb{R}^n$ is the tropical projective space

$$\mathbb{TP}^n = (\overline{\mathbb{R}}^{n+1} \setminus \{(\infty, \dots, \infty)\}) / \mathbb{R}(1, \dots, 1)$$

which is endowed with the quotient topology. The Berkovich projective space $(\mathbb{P}_K^n)^{\text{an}}$ can be described as the set of equivalence classes of elements in $(\mathbb{A}_K^{n+1})^{\text{an}} \setminus \{0\}$ with respect to the following equivalence relation: Let x and y be points in $(\mathbb{A}_K^{n+1})^{\text{an}} \setminus \{0\}$, i.e. multiplicative seminorms on $K[x_0, \dots, x_n]$ extending the absolute value on K . They are equivalent if and only if there exists a constant $c > 0$ such that for every homogeneous polynomial f of degree d we have $|f(y)| = c^d |f(x)|$.

We can describe the tropicalization map on the toric variety \mathbb{P}_K^n as follows:

$$(2.2) \quad \text{trop} : (\mathbb{P}_K^n)^{\text{an}} \rightarrow \mathbb{TP}^n$$

maps the class of the seminorm p on $K[x_0, \dots, x_n]$ to the class $(-\log |p(x_0)|, \dots, -\log |p(x_n)|) + \mathbb{R}(1, \dots, 1)$ in tropical projective space.

An important result about the relation between tropical and analytic geometry due to Payne says that for every quasi-projective variety X over K the associated analytic space X^{an} is homeomorphic to the inverse limit over all $\text{Trop}_{\varphi}(X)$, where the limit runs over all all closed embeddings $\varphi : X \hookrightarrow Y$ in a quasi-projective toric variety Y (see [Pay09], Theorem 4.2). For a generalization which omits the quasi-projectivity hypothesis see [FGP14].

Hence tropicalizations are combinatorial images of analytic spaces, which recover the full analytic space in the projective limit. Individuell tropicalizations however may not faithfully depict all topological features of the analytic space. We will discuss a basic example at the beginning of the next section.

3. THE CASE OF CURVES

Let K be a field which is algebraically closed and complete with respect to a non-archimedean, non-trivial absolute value, and let X be a smooth curve over K .

As an example, let us consider an elliptic curve E in Weierstrass form $y^2 = x^3 + ax + b$ over K . The affine curve E has a natural embedding in \mathbb{A}_K^2 . The intersection E_0 of E with $\mathbb{G}_{m,K}^2$ has a natural tropicalization in \mathbb{R}^2 , which is the tree depicted in the next figure.




FIGURE 1. Tropicalization of an elliptic curve in Weierstrass form. The left hand side shows the case $3v(a) \geq 2v(b)$, the right hand side the case $3v(a) < 2v(b)$. The numbers on the edges indicate the tropical multiplicities. If no number is given, the multiplicity is one.

If however E is a Tate curve, then the analytic space E^{an} contains a circle, i.e. it has topological genus 1. Hence the Weierstrass tropicalization does not faithfully depict the topology of the analytic space.

As stated before, if X is a smooth curve over K , the Berkovich space X^{an} is some kind of generalized graph, allowing infinite ramification along a dense set of points. In particular, it makes sense to talk about its leaves. The space of non-leaves $H_0(X^{\text{an}})$ in the Berkovich analytification X^{an} admits a natural metric which is defined via semistable models, see [BPR11], section 5. Any tropicalization of X with respect to a closed embedding in a torus carries a natural metric which is locally given by lattice length on each edge (and globally by shortest paths). Another problem in the case of the Weierstrass tropicalizations from figure 1 is the fact that tropical multiplicities $\neq 1$ are present. This may indicate that the tropicalization map will not be isometric on a subgraph, see [BPR11], Corollary 6.9.

It is explained in [BPR11], Theorem 7.2, how one can construct better tropicalizations of the Tate curve.

Based on a detailed study of the structure of analytic curves there are the following two important comparison theorems in [BPR11].

Theorem 3.1 ([BPR11], Theorem 6.20). *Let Γ be any finite subgraph of X^{an} for a smooth, complete curve X over K . Then there exists a closed immersion $\varphi : X \hookrightarrow Y$ into a toric variety Y with dense torus T such that the restriction $\varphi_0 : X_0 = X \cap \varphi^{-1}(T) \hookrightarrow T$ induces a tropicalization map $\text{trop}_{\varphi_0} : X_0^{\text{an}} \rightarrow \mathbb{R}^n$ which maps Γ homeomorphically and isometrically onto its image.*

Theorem 3.2 ([BPR11], Theorem 6.23). *Consider a smooth curve X_0 over K and a closed immersion $\varphi : X_0 \hookrightarrow T$ into a split torus with associated tropical variety $\text{Trop}_{\varphi_0}(X_0)$. If Γ' is a compact connected subset of $\text{Trop}_{\varphi_0}(X_0)$ which has tropical multiplicity one at every point with coordinates in $\log |K^*|$, then there exists a unique closed subset Γ in $H_0(X^{\text{an}})$ mapping homeomorphically onto Γ' , and this homeomorphism is in fact an isometry.*

We will discuss higher-dimensional generalizations of these theorems in section 7.

4. TROPICAL GRASSMANNIANS

4.1. The setting. In view of the results [BPR11] for curves, it is a natural question whether they can be generalized to varieties of higher dimensions. As a first example, the tropical Grassmannian of planes was studied in [CHW14]. In this section we discuss the main theorem of this paper.

For natural numbers $d \leq n$ we denote by $\text{Gr}(d, n)$ the Grassmannian of d -dimensional subspaces of n -space. The tropical Grassmannian is defined as the tropicalization of $\text{Gr}(d, n)$ with respect to the Plücker embedding $\varphi : \text{Gr}(d, n) \rightarrow \mathbb{P}^{\binom{n}{d}-1}$. Roughly speaking, the Plücker embedding maps the point corresponding to the d -dimensional subspace W to the line in $\binom{n}{d}$ -space given by the d -th exterior power of W .

During this section we will deviate from the exposition in section 2.3 and consider tropicalizations with respect to the negative valuation map. Hence our tropical projective space is

$$\mathbb{TP}^{\binom{n}{d}-1} = ((\mathbb{R} \cup \{-\infty\})^{\binom{n}{d}} \setminus \{(-\infty, \dots, -\infty)\}) / \mathbb{R}(1, \dots, 1),$$

and the tropical Grassmannian $\mathcal{T}\text{Gr}(d, n) = \text{Trop}_{\varphi}(\text{Gr}(d, n))$ is defined as the image of $\text{trop} \circ \varphi^{\text{an}} : \text{Gr}(d, n)^{\text{an}} \rightarrow \mathbb{TP}^{\binom{n}{d}-1}$, where trop is given by the map

$$p \mapsto (\log |p(x_0)|, \dots, \log |p(x_n)|) + \mathbb{R}(1, \dots, 1)$$

on the analytic projective space, see (2.2).

Tropical Grassmannians were first studied by Speyer and Sturmfels [SS04] who focused on the toric part $\text{Gr}_0(d, n) = \text{Gr}(d, n) \cap \varphi^{-1}(\mathbb{G}_{m, K}^{\binom{n}{d}-1})$ which is embedded in the torus $\mathbb{G}_{m, K}^{\binom{n}{d}-1}$ via φ . The tropicalization $\mathcal{T}\text{Gr}_0(d, n) = \text{Trop}_{\varphi}(\text{Gr}_0(d, n))$ (which is called $\mathcal{G}'_{d, n}$ in [SS04]) is a fan of dimension $d(n-d)$ containing a lineality space of dimension $n-1$, see [SS04], section 3.

Moreover, if $d = 2$, Speyer and Sturmfels proved that $\mathcal{T}\mathrm{Gr}_0(2, n)$ can be identified with the space of phylogenetic trees, see [SS04], section 4. For our purposes, a phylogenetic tree on n leaves is a pair (T, ω) , where T is a combinatorial tree with no degree-two vertices together with a labeling of its leaves in bijection with $\{1, \dots, n\}$, and ω is a real-valued function on the set of edges of T . The tree T is called the combinatorial type of the phylogenetic tree (T, ω) . For every phylogenetic tree (T, ω) and all $i \neq j$ in $\{1, \dots, n\}$ we denote by x_{ij} the sum of the weights along the uniquely determined path from leaf i to leaf j . This tree-distance function satisfies the four-point-condition, which states that for all pairwise distinct indices i, j, k, l in $\{1, \dots, n\}$ the maximum among

$$x_{ij} + x_{kl}, \quad x_{ik} + x_{jl}, \quad x_{il} + x_{jk}$$

is attained at least twice.

4.2. A section of the tropicalization map. For the rest of this section we will always consider the case $d = 2$. In [CHW14], we investigate the full projective Grassmannian $\mathcal{T}\mathrm{Gr}(2, n) = \mathrm{Trop}_\varphi \mathrm{Gr}(2, n)$ in tropical projective space $\mathbb{TP}^{(n)}{}^{-1}$. The main result of this paper is the following theorem:

Theorem 4.1. *There exists a continuous section $\sigma: \mathcal{T}\mathrm{Gr}(2, n) \rightarrow \mathrm{Gr}(2, n)^{\mathrm{an}}$ of the tropicalization map $\mathrm{trop} \circ \varphi^{\mathrm{an}}: \mathrm{Gr}(2, n)^{\mathrm{an}} \rightarrow \mathcal{T}\mathrm{Gr}(2, n)$. Hence, the tropical Grassmannian $\mathcal{T}\mathrm{Gr}(2, n)$ is homeomorphic to a closed subset of the Berkovich analytic space $\mathrm{Gr}(2, n)^{\mathrm{an}}$.*

Note that we are not considering semistable models or their Berkovich skeleta in this approach.

The first idea one might have is to look at the big open cells of the Grassmannians. Since $d = 2$, the coordinates of the ambient projective space are indexed by the two element subsets $\{i, j\}$ of $\{1, \dots, n\}$. If $i < j$, we write p_{ij} for this coordinate, and we put $p_{ji} = -p_{ij}$. We intersect the Grassmannian with the standard open affine covering of projective space and get open affine subvarieties

$$U_{ij} = \mathrm{Gr}(2, n) \cap \varphi^{-1}\{p_{ij} \neq 0\}.$$

The affine Plücker coordinates are $u_{kl} = p_{kl}/p_{ij}$. Since the Plücker ideal is generated by the relations $p_{ij}p_{kl} - p_{ik}p_{jl} + p_{il}p_{jk} = 0$ for $\{i, j, k, l\}$ running over the four-element subsets of $\{1, \dots, n\}$, the subvariety U_{ij} is isomorphic to $\mathbb{A}^{n(n-2)}$ via the coordinates $u_{ik} = p_{ik}/p_{ij}$ and $u_{jk} = p_{jk}/p_{ij}$ for all k different from i and j .

Recall that $(\mathbb{A}^{n(n-2)})^{\mathrm{an}}$ is the set of all multiplicative seminorms on $K[u_{ik}, u_{jk} : k \notin \{i, j\}]$ which extend the absolute value on K . As explained in section 2, the toric variety $\mathbb{A}^{n(n-2)}$ has a natural tropicalization. With the sign conventions in this section, it is given by

$$\begin{aligned} \mathrm{trop} : (\mathbb{A}^{n(n-2)})^{\mathrm{an}} &\rightarrow (\mathbb{R} \cup \{-\infty\})^{2(n-2)} \\ p &\mapsto (\log |p(u_{ik})|, \log |p(u_{jk})|)_{k \notin \{i, j\}} \end{aligned}$$

If we denote by $\iota : U_{ij} \rightarrow \mathbb{A}^{n(n-2)}$ the natural isomorphism, we get the tropicalization map


$$\text{trop} \circ \iota^{\text{an}} : U_{ij}^{\text{an}} \longrightarrow (\mathbb{R} \cup \{-\infty\})^{2(n-2)}.$$

Now the tropicalization map on an affine space $(\mathbb{A}_K^N)^{\text{an}}$ has a natural section, which is given by the map $\delta : (\mathbb{R} \cup \{-\infty\})^N \rightarrow (\mathbb{A}_K^N)^{\text{an}}$, mapping a point $r = (r_1, \dots, r_N) \in (\mathbb{R} \cup \{-\infty\})^N$ to the seminorm

$$(4.1) \quad \delta(r) : K[x_1, \dots, x_N] \longrightarrow \mathbb{R}_{\geq 0} \quad \left| \sum_{\alpha \in \mathbb{N}_0^N} c_\alpha x^\alpha(\delta(r)) \right| = \max_\alpha \{ |c_\alpha| \prod_{i=1}^N \exp(r_i \alpha_i) \}.$$

Here we put $\exp(-\infty) = 0$. Note that the restriction of δ to \mathbb{R}^N is a map from \mathbb{R}^N to the torus $(\mathbb{G}_m^N)^{\text{an}}$. We call the image of δ on $(\mathbb{R} \cup \{-\infty\})^N$ the standard skeleton of $(\mathbb{A}_K^N)^{\text{an}}$, and the image of $\delta|_{\mathbb{R}^N}$ the standard skeleton of the torus $(\mathbb{G}_m^N)^{\text{an}}$.

However, the map δ for $N = n(n-2)$ does not in general provide a section of the tropicalization map on the whole of U_{ij} . Consider $n \geq 4$, and let $x = (x_{kl})_{kl}$ be a point in the tropical Grassmannian $\mathcal{TGr}(2, n)$. We may assume that $x_{ij} = 0$. Let ω be the projection to the affine coordinates $\omega = (x_{ik}, x_{jk})_{k \notin \{i, j\}} \in \mathbb{R}^{2(n-2)}$. Then $\delta(\omega)$ is a point in the Berkovich space U_{ij}^{an} with $\log |u_{ik}(\delta(\omega))| = x_{ik}$ and $\log |u_{jk}(\delta(\omega))| = x_{jk}$. Since $u_{kl} = u_{ik}u_{jl} - u_{il}u_{jk}$ by the Plücker relations, the definition of δ gives us $\log |u_{kl}(\delta(\omega))| = \max\{x_{ik} + x_{jl}, x_{il} + x_{jk}\}$. Hence δ can only provide a section of the tropicalization map if this maximum is equal to x_{kl} . This may fail if the labelled tree T has the wrong shape, e.g. if it looks like this:



The strategy for the proof of Theorem 4.1 is to compare the tropical Grassmannian with a standard skeleton of an affine space on a smaller pieces of the tropicalization, namely on the part consisting of phylogenetic trees such that the underlying combinatorial tree has the right shape. If we want to make this precise, the definition of these smaller pieces is quite involved. A considerable part of the difficulties is due to the fact that we take into account the boundary strata of the Grassmannian in projective space. In order to explain the general strategy we will therefore restrict our attention to the torus part of the tropical Grassmannian. The general case can be found in [CHW14].

4.3. Sketch of proof in the dense torus orbit. In this section we explain the proof of Theorem 4.1 for the subset $\mathcal{TGr}_0(2, n)$ of the tropical Grassmannian $\mathcal{TGr}(2, n)$. Recall that $\mathcal{TGr}_0(2, n)$ is the tropicalization of the dense open subset $\text{Gr}_0(2, n)$ of the Grassmannian which is mapped to the torus via the Plücker map.

We fix a pair ij as above and work in the big open cell U_{ij} . By means of the isomorphism ι , the coordinate ring R_{ij} of U_{ij} is a polynomial ring $K[u_{ik}, u_{jk} : k \notin \{i, j\}]$ in $2(n-2)$ variables. The other Plücker coordinates are expressed as $u_{kl} = u_{ik}u_{jl} - u_{il}u_{jk}$ in R_{ij} . Note that the affine variety $\text{Gr}_0(2, n)$ is contained in U_{ij} . The coordinate ring of $\text{Gr}_0(2, n)$ is equal to the localization of R_{ij} after the multiplicative subset generated by all u_{kl} for $\{k, l\} \neq \{i, j\}$.

We also fix a labelled tree T with n leaves $1, \dots, n$, and arrange T as in the following figure




FIGURE 2. Arrangement of T

with subtrees T_1, \dots, T_r .

Definition 4.2. Let \preceq be a partial order on the set $\{1, \dots, n\} \setminus \{i, j\}$. We write $k \prec l$ if $k \preceq l$ and $k \neq l$. Then \preceq has the cherry property on T with respect to i and j if the following conditions hold:

- (i) Two leaves of different subtrees T_a and T_b for $a, b \in \{1, \dots, r\}$ as in figure 2 cannot be compared by \preceq .
- (ii) The partial order \preceq restricts to a total order on the leaf set of each T_a , $a = 1, \dots, r$.
- (iii) If $k \prec l \prec m$, then either $\{k, l\}$ or $\{l, m\}$ is a cherry of the quartet $\{i, k, l, m\}$, i.e. the subtree given by this quartet of leaves contains a node which is adjacent to both elements of the cherry:




FIGURE 3. Cherry property

An induction argument shows the following lemma:

Lemma 4.3 ([CHW14], Lemma 4.7). Fix a pair of indices i, j , and let T be a tree on n labelled leaves. Then, there exists a partial order \preceq on the set $\{i, \dots, n\} \setminus \{i, j\}$ that has the cherry property on T with respect to i and j .

For an example of the inductive construction of such a partial order see [CHW14], Figure 3.

The leaves in each subtree T_a are totally ordered, say as $s_1 \prec s_2 \prec \dots \prec s_p$, if T_a contains $p = p(a)$ leaves. We consider the variable set $I_a = \{u_{is_1}, \dots, u_{is_p}\} \cup \{u_{js_1}, u_{s_1s_2}, \dots, u_{s_{p-1}s_p}\}$. Then $I = I_1 \cup \dots \cup I_r$ is a set of $2(n-2)$ affine Plücker coordinates of the form u_{kl} .

We can successively reconstruct the variables u_{jl} not contained in I as follows: In the tree T_a with leaves $s_1 \prec s_2 \prec \dots \prec s_p$, we have $u_{s_1 s_2} = u_{i s_1} u_{j s_2} - u_{i s_2} u_{j s_1}$, hence

$$u_{j s_2} = u_{i s_1}^{-1} (u_{s_1 s_2} + u_{i s_2} u_{j s_1}).$$

The right hand side is an expression in the variables contained in the affine coordinate set I with $u_{i s_1}$ inverted. Now we use the relation $u_{s_2 s_3} = u_{i s_2} u_{j s_3} - u_{i s_3} u_{j s_2}$ to express $u_{j s_3} = u_{i s_2}^{-1} (u_{s_2 s_3} + u_{i s_3} u_{j s_2})$. Plugging in the expression for $u_{j s_2}$ we can write $u_{j s_3}$ as a polynomial in the variables in I plus all $u_{i k}^{-1}$.

Proceeding by induction, we find for all $m \neq i, j$ that

$$u_{jm} \in K[u_{kl} : u_{kl} \in I][u_{ik}^{-1} : k \neq i, j]$$

and hence

$$(4.2) \quad K[u_{kl} : u_{kl} \in I] \subset R_{ij} \subset K[u_{kl} : u_{kl} \in I][u_{ik}^{-1} : k \neq i, j].$$

This shows that the variable set I generates the function field $\text{Quot}(R_{ij})$ of U_{ij} . Recall that the coordinate ring of $\text{Gr}_0(2, n)$ is $K[\text{Gr}_0(2, n)] = S^{-1}R_{ij}$, where S is the multiplicative subset of R_{ij} generated by all u_{kl} for $\{kl\} \neq \{ij\}$. By the previous result, $K[\text{Gr}_0(2, n)]$ is equal to the localization of the Laurent polynomial ring $K[u_{kl}^{\pm} : u_{kl} \in I]$ after the multiplicative subset generated by *all* u_{kl} expressed as Laurent polynomials in the coordinates contained in I .

Definition 4.4. Let \mathcal{C}_T be the cone in $\mathcal{T}\text{Gr}_0(2, n)$ whose interior corresponds to the phylogenetic trees with underlying tree T . Let $x = (x_{kl})_{kl} + \mathbb{R}(1, \dots, 1)$ be a point in $\mathcal{C}_T \subset \mathcal{T}\text{Gr}_0(2, n)$. We associate to it a point $\sigma_T^{ij}(x)$ in $\text{Gr}_0(2, n)^{\text{an}}$, i.e. a multiplicative seminorm on the coordinate ring of $\text{Gr}_0(2, n)$, as follows. For every Laurent polynomial $f = \sum_{\alpha} c_{\alpha} u^{\alpha} \in K[u_{kl}^{\pm} : u_{kl} \in I]$ (where α runs over the maps from I to \mathbb{Z}) we put

$$|f(\sigma_T^{ij}(x))| = \max_{\alpha} \{ |c_{\alpha}| \prod_{u_{kl} \in I} \exp(\alpha_{kl}(x_{kl} - x_{ij})) \}, \quad \text{where } \alpha = (\alpha_{kl})_{u_{kl} \in I}.$$

This defines a multiplicative norm on $K[u_{kl}^{\pm} : \{kl\} \in I]$ which has a unique extension to a multiplicative norm on the localization $K[\text{Gr}_0(2, n)]$. Let $\sigma_T^{ij}(x)$ be the resulting point in the Berkovich space $\text{Gr}_0(2, n)^{\text{an}}$.

Since for every $f \in K[\text{Gr}_0(2, n)]$ the evaluation map on \mathcal{C}_T given by

$$x \mapsto |f(\sigma_T^{ij}(x))|$$

is continuous, we have constructed a continuous map

$$\sigma_T^{ij} : \mathcal{C}_T \rightarrow \text{Gr}_0(2, n)^{\text{an}}.$$

We want to show that it is a section of the tropicalization map $\text{trop} \circ \varphi^{\text{an}} : \text{Gr}_0(2, n)^{\text{an}} \rightarrow \mathcal{T}\text{Gr}_0(2, n)$ on $\text{trop}^{-1}(\mathcal{C}_T)$, which amounts to checking that

$$\log |u_{kl}(\sigma_T^{ij}(x))| = x_{kl} - x_{ij} \text{ for all } \{kl\} \neq \{ij\} \text{ and for all } x \in \mathcal{C}_T.$$

For variables u_{kl} in I this is clear from the definition of σ_T^{ij} . This holds in particular for all indices of the form $\{ik\}$ for $k \notin \{i, j\}$. In order to check this fact for the other indices, recall the definition of I after Lemma 4.3. Since $u_{js_2} = u_{is_1}^{-1}(u_{s_1s_2} + u_{is_2}u_{js_1})$, we find

$$\log |u_{js_2}(\sigma_T^{ij}(x))| = \max\{-x_{is_1} + x_{s_1s_2}, -x_{is_1} + x_{is_2} + x_{js_1} - x_{ij}\}.$$

Since s_1 and s_2 are in the same subtree (see figure 2), we find that $x_{is_1} + x_{js_2} = x_{is_2} + x_{js_1} \geq x_{ij} + x_{s_1s_2}$, which implies $\log |u_{js_2}(\sigma_T^{ij}(x))| = x_{js_2} - x_{ij}$. Inductively, we can show in this way our claim for all indices of the form $\{jk\}$. If we consider an index of the form $\{kl\}$ where $k, l \notin \{i, j\}$, we have $u_{kl} = u_{ik}u_{jl} - u_{il}u_{jk}$. If k is a leaf in the subtree T_a and l is a leaf in the subtree T_b for $a < b$, we find $x_{kl} + x_{ij} = x_{il} + x_{jk} > x_{ik} + x_{jl}$. Hence the non-archimedean triangle inequality gives $\log |u_{kl}(\sigma_T^{ij}(x))| = \log |(u_{il}u_{jk})(\sigma_T^{ij}(x))| = x_{kl} - x_{ij}$. If k and l are leaves in the same subtree T_a , we may assume that $k \prec l$. If k is the predecessor of l in the total ordering \preceq restricted to T_a , we are done, since then u_{kl} is contained in I . If not, we let m be the predecessor of l , so that $k \prec m \prec l$. The Plücker relations give $u_{im}u_{kl} = u_{ik}u_{ml} + u_{il}u_{km}$, hence

$$u_{kl} = u_{im}^{-1}(u_{ik}u_{ml} + u_{il}u_{km}).$$

Note that all variables on the right hand side except possibly u_{km} are contained in I . Hence we can calculate $\log |u_{kl}(\sigma_T^{ij}(x))|$ by developing u_{km} in a Laurent series in the variables in I . The resulting Laurent series does not allow any cancellation between the variables. Now we can apply the cherry property for the ordering \preceq , which says that $\{k, m\}$ or $\{m, l\}$ is a cherry for the quartet $\{i, k, m, l\}$. Hence we have

$$x_{il} + x_{km} \leq x_{im} + x_{kl} = x_{ik} + x_{ml}$$

or

$$x_{ik} + x_{ml} \leq x_{im} + x_{kl} = x_{il} + x_{km}.$$

In both cases, one can check directly that our claim holds.

In fact, the section σ_T^{ij} has a more conceptual description. Let $\psi : \text{Gr}_0(2, n) \rightarrow \text{Spec}K[u_{kl}^{\pm} : u_{kl} \in I] = \mathbb{G}_m^{2(n-2)}$ be the open embedding induced by (4.2), and let Σ be the skeleton of $(\mathbb{G}_m^{2(n-2)})^{\text{an}}$ as in (4.1). It is contained in the open analytic subvariety $\text{Gr}_0(2, n)^{\text{an}}$, since it consists entirely of norms and does therefore not meet a closed subvariety of strictly lower dimension. Then the map σ_T^{ij} is the composition of the projection from \mathcal{C}_T to the coordinates $(x_{kl} - x_{ij})_{u_{kl} \in I} \in \mathbb{R}^{2(n-2)} = \Sigma$ followed by the inclusion of Σ in $\text{Gr}_0(2, n)^{\text{an}}$.

Recall from section 2.3 that for every $x \in \mathcal{T}\text{Gr}_0(2, n)$ the preimage $\text{trop}_{\varphi}^{-1}(x)$ under the tropicalization map is the Berkovich spectrum $\mathcal{M}(A_x)$ of an affinoid algebra A_x . It follows from our construction that $\sigma_T^{ij}(x)$ is the unique Shilov boundary point of A_x . We can formulate this fact explicitly as follows.

Lemma 4.5. *[[CHW14], lemma 4.17] For every $x \in \mathcal{C}_T$ the seminorm $\sigma_T^{ij}(x)$ constructed above has the following maximality property: For every f in the coordinate ring*

$K[\mathrm{Gr}_0(2, n)]$ and every seminorm $p \in \mathrm{trop}_\varphi^{-1}(x) \subset \mathrm{Gr}_0(2, n)^{\mathrm{an}}$ we have

$$|f(p)| \leq |f(\sigma_T^{ij}(x))|.$$

As an immediate consequence we see that $\sigma_T^{ij}(x)$ does not depend on the choice of the partial ordering \preceq or on the pair ij , and that $\sigma_T^{ij}(x) = \sigma_{T'}^{ij}(x)$ if x is contained in the intersection $\mathcal{C}_T \cap \mathcal{C}_{T'}$. Hence we can patch these maps together and get a well-defined section $\sigma : \mathcal{T}\mathrm{Gr}_0(2, n) \rightarrow \mathrm{Gr}_0(2, n)^{\mathrm{an}}$ of the tropicalization map. It follows from the construction that it is continuous.

This proves theorem 4.1 on the dense torus orbit $\mathrm{Gr}_0(2, n)$. In order to define a section on the tropicalization map also on the boundary strata of $\mathrm{Gr}(2, n)$, we follow the same strategy of constructing an index set I such that the associated Plücker variables generate the function field of the Grassmannian. This construction is more involved, see [CHW14], section 4 for details. The most important problem here is to find these local index sets in such a way that the section is also continuous when passing from one stratum of the Grassmannian for another. This continuity statement is shown in [CHW14], Theorem 4.19.

Using the index set I , we may also calculate the initial degenerations of all points in the tropical Grassmannian and deduce that their tropical multiplicity is one everywhere. We will explain a general argument showing the existence of a section in this case in paragraph 7.2.

In [DP14], Draisma and Postingshel give a different proof of Theorem 4.1. They work with the affine cone over the Grassmannian $\mathrm{Gr}(2, n)$, define the section on a suitable subset and use torus actions and tropicalized torus actions to move it around. Also in this approach a maximality statement such as Lemma 4.5 is used.

5. SKELETA OF SEMISTABLE PAIRS

In the next three sections, we give an overview of the results in [GRW14]. In order to compare polyhedral substructures of Berkovich spaces with tropicalizations, we start by generalizing Berkovich's notion of skeleta. Such skeleta are induced by the incidence complex of the special fiber of suitable models. We extend this notion by adding a horizontal divisor on the model. For the rest of this paper, we fix an algebraically closed ground field K which is complete with respect to a non-archimedean, non-trivial absolute value.

5.1. Integral affine structures. For curves, metrics play an important role in the comparison results between tropical and analytic varieties, as we have seen in section 3. The right way to generalize this to higher dimensions is to consider integral affine structures. Let M be a lattice in the finite-dimensional real vector space $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$, and let

$N = \text{Hom}(N, \mathbb{Z})$ its dual, which is a lattice in the dual space $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$ of $M_{\mathbb{R}}$. We denote the associated pairing by $\langle \cdot, \cdot \rangle : M_{\mathbb{R}} \times N_{\mathbb{R}} \rightarrow \mathbb{R}$. Recall that $\Gamma = \log |K^*|$.

An *integral Γ -affine polyhedron* in $N_{\mathbb{R}}$ is a subset of $N_{\mathbb{R}}$ of the form

$$\Delta = \{v \in N_{\mathbb{R}} \mid \langle u_i, v \rangle + \gamma_i \geq 0 \text{ for all } i = 1, \dots, r\}$$

for some $u_1, \dots, u_r \in M$ and $\gamma_1, \dots, \gamma_r \in \Gamma$. Any face of an integral Γ -affine polyhedron Δ is again integral Γ -affine. An *integral Γ -affine polyhedral complex* in $N_{\mathbb{R}}$ is a polyhedral complex whose faces are integral Γ -affine.

An *integral Γ -affine function* on $N_{\mathbb{R}}$ is a function from $N_{\mathbb{R}}$ to \mathbb{R} which is of the form

$$v \mapsto \langle u, v \rangle + \gamma$$

for some $u \in M$ and $\gamma \in \Gamma$. More generally, let M' be a second finitely generated free abelian group and let $N' = \text{Hom}(M', \mathbb{Z})$. An *integral Γ -affine map* from $N_{\mathbb{R}}$ to $N'_{\mathbb{R}}$ is a function of the form $F = \phi^* + v$, where $\phi : M' \rightarrow M$ is a homomorphism, $\phi^* : N_{\mathbb{R}} \rightarrow N'_{\mathbb{R}}$ is the dual homomorphism extended to $N_{\mathbb{R}}$, and $v \in N' \otimes_{\mathbb{Z}} \Gamma$. If $N' = M' = \mathbb{Z}^m$ and $F = (F_1, \dots, F_m) : N_{\mathbb{R}} \rightarrow \mathbb{R}^m$ is a function, then F is integral Γ -affine if and only if each coordinate $F_i : N_{\mathbb{R}} \rightarrow \mathbb{R}$ is integral Γ -affine.

An *integral Γ -affine map* from an integral Γ -affine polyhedron $\Delta \subset N_{\mathbb{R}}$ to $N'_{\mathbb{R}}$ is defined as the restriction to Δ of an integral Γ -affine map $N_{\mathbb{R}} \rightarrow N'_{\mathbb{R}}$. If $\Delta' \subset N'_{\mathbb{R}}$ is an integral Γ -affine polyhedron then a function $F : \Delta \rightarrow \Delta'$ is *integral Γ -affine* if the composition $\Delta \rightarrow \Delta' \hookrightarrow N'_{\mathbb{R}}$ is integral Γ -affine. We say that an integral Γ -affine map $F : \Delta \rightarrow N'_{\mathbb{R}}$ is *unimodular* if F is injective and if the inverse map $F(\Delta) \rightarrow \Delta$ is integral Γ -affine. Note that $F(\Delta)$ is an integral Γ -affine polyhedron in $N'_{\mathbb{R}}$.

5.2. Semistable pairs. Note that since our ground field K is algebraically closed, the ring of integers K° is a valuation ring which is not discrete and not noetherian. We begin by describing the building blocks of the polyhedral substructures of Berkovich spaces which lend themselves to comparison with tropicalizations.

Let $0 \leq r \leq d$ be natural numbers and consider the K° -scheme

$$\mathcal{S} = \text{Spec}(K^\circ[x_0, \dots, x_d]/(x_0 \dots x_r - \pi))$$

for some $\pi \in K^\circ$ satisfying $|\pi| < 1$. Then \mathcal{S} is a flat scheme over K° with smooth generic fiber. Its special fibre contains $r + 1$ irreducible components whose incidence complex is an r -dimensional simplex. Now fix a natural number $s \geq 0$ such that $r + s \leq d$, and consider the principal Cartier divisor $H(s) = \text{div}(x_{r+1}) + \dots + \text{div}(x_{r+s})$ on \mathcal{S} .

As explained in section 2.2, the K° -scheme \mathcal{S} gives rise to two analytic spaces in the following way. On the one hand, we have an associated admissible formal scheme

$$(5.1) \quad \mathcal{S} = \text{Spf}(K^\circ\{x_0, \dots, x_d\}/(x_0 \dots x_r - \pi))$$

which we get by completing \mathcal{S} with respect to a non-zero element in K of absolute value < 1 . Its analytic generic fiber is $\mathcal{M}(A_{\mathcal{S}})$ for the affinoid algebra

$$A_{\mathcal{S}} = K\{x_0, \dots, x_d\}/(x_0 \dots x_r - \pi).$$

It is a subset of the analytification of the generic fibre $\mathcal{S}_K = \text{Spec}(K[x_0, \dots, x_d]/(x_0 \dots x_r - \pi))$ of \mathcal{S} .

Now we look at the tropicalization map on $\mathcal{M}(A_{\mathcal{S}})$ which only takes into account the first $r + s + 1$ coordinates, i.e. the map

$$\begin{aligned} \text{Val} : \mathcal{M}(A_{\mathcal{S}}) &\rightarrow \mathbb{R}_{\geq 0}^{r+s} \\ p &\mapsto (-\log |x_0(p)|, \dots, -\log |x_{r+s}(p)|). \end{aligned}$$

Its image is $\Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s$, where $\Delta(r, \pi) = \{(v_0, \dots, v_r) \in \mathbb{R}_{\geq 0}^{r+1} : v_0 + \dots + v_r = -\log(|\pi|)\}$ is a simplex in \mathbb{R}^{r+1} .

We define a continuous section $\psi : \Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s \rightarrow \mathcal{M}(A_{\mathcal{S}})$ as follows. Note that the projection to (x_1, \dots, x_d) induces an isomorphism from $\mathcal{M}(A_{\mathcal{S}})$ to the affinoid subdomain $B = \{p : -\log |x_1(p)| - \dots - \log |x_r(p)| \leq -\log |\pi|\}$ of $\mathcal{M}(K\{x_1, \dots, x_d\})$.

Similarly, projection to (v_1, \dots, v_r) induces a homeomorphism

$$\Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s \rightarrow \Sigma = \{(v_1, \dots, v_{r+s}) \in \mathbb{R}_{\geq 0}^{r+s} : v_1 + \dots + v_r \leq -\log |\pi|\}$$

For every $v = (v_1, \dots, v_{r+s}) \in \Sigma$ there is a bounded multiplicative norm $\|\cdot\|_v$ on $K\{x_1, \dots, x_d\}$ which is defined as follows:

$$\| \sum_{I=(i_1, \dots, i_d)} a_I x^I \|_v = \max_I \{ |a_I| \exp(-i_1 v_1 - \dots - i_{r+s} v_{r+s}) \}.$$

It satisfies $-\log \|x_1\|_v - \dots - \log \|x_r\|_v = v_1 + \dots + v_d \leq -\log |\pi|$. Therefore the point $\|\cdot\|_v$ is contained in the affinoid domain B . Hence there is a uniquely determined continuous map

$$\sigma : \Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s \rightarrow \mathcal{M}(A_{\mathcal{S}})$$

making the diagram

$$\begin{array}{ccc} \Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s & \xrightarrow{\sigma} & \mathcal{M}(A_{\mathcal{S}}) \\ \downarrow & & \downarrow \\ \Sigma & \xrightarrow{v \mapsto \|\cdot\|_v} & B \end{array}$$

commutative. The map σ is by construction a section of the map Val . We define $S(\mathcal{S}, H(s)) \subset \mathcal{M}(A_{\mathcal{S}}) \subset (\mathcal{S}_K)^{\text{an}}$ as the image of σ and call it the skeleton of the pair $(\mathcal{S}, H(s))$.

Now we consider schemes which étale locally look like some \mathcal{S} .

Definition 5.1. A strictly semistable pair (\mathcal{X}, H) consists of an irreducible proper flat scheme \mathcal{X} over the valuation ring K° and a sum $H = H_1 + \cdots + H_S$ of effective Cartier divisors H_i on \mathcal{X} such that \mathcal{X} is covered by open subsets \mathcal{U} which admit an étale morphism

$$(5.2) \quad \psi : \mathcal{U} \longrightarrow \mathcal{S} = \text{Spec}(K^\circ[x_0, \dots, x_d]/(x_0 \cdots x_r - \pi))$$

for some $r \leq d$ and $\pi \in K^\circ$ with $|\pi| < 1$. We assume that each H_i has irreducible support and that the restriction of H_i to \mathcal{U} is either trivial or defined by $\psi^*(x_j)$ for some $j \in \{r+1, \dots, d\}$.

This is a generalization of de Jong's notion of a strictly semistable pair over a discrete valuation ring [dJ96], if we add the divisor of the special fiber to the horizontal divisor H . It is sometimes convenient to include only the horizontal divisor H as part of the data, since it is a Cartier divisor, whereas the special fiber of \mathcal{X} may not be one. For a more detailed discussion of this issue see [GRW14], Proposition 4.17. If $d = 1$, i.e. in the case of curves, semistable models and therefore nice skeletons are always available after an extension of the ground field. In higher dimensions, we have to allow alterations in order to find semistable models, see [dJ96], Theorem 6.5 and (for a generalization to the non-noetherian case) [GRW14], Proposition 5.6.

5.3. Skeleta. Let (\mathcal{X}, H) be a semistable pair in the sense of definition 5.1. The generic fiber X of \mathcal{X} is smooth of dimension d . Hence d is constant in every chart \mathcal{U} , whereas the numbers r and s may vary with \mathcal{U} . We denote the special fiber of \mathcal{X} by $\mathcal{X}_s = \mathcal{X} \otimes_{K^\circ} \tilde{K}$.

Denote by V_1, \dots, V_R the irreducible components of the special fiber \mathcal{X}_s of \mathcal{X} . The Cartier divisors H_i which are part of our data give rise to horizontal closed subschemes \mathcal{H}_i of \mathcal{X} , which are locally cut out by a defining equation of H_i . Putting $D_i = V_i$ for $i = 1, \dots, R$ and $D_{i+R} = \mathcal{H}_i$, we get a Weil divisor $D = \sum_{i=1}^{R+S} D_i$ on \mathcal{X} . This gives rise to a stratification of \mathcal{X} , where a stratum is defined as an irreducible component of a set of the form $\bigcap_{i \in I} D_i \setminus \bigcup_{i \notin I} D_i$ for some $I \subset \{1, \dots, R+S\}$. We call any stratum contained in the special fiber \mathcal{X}_s a vertical stratum and denote the set of all vertical strata by $\text{str}(\mathcal{X}_s, H)$.

Now we want to glue skeleta of local charts together in such a way that the faces of the resulting polyhedral complex are in bijective correspondence with the vertical strata in $\text{str}(\mathcal{X}_s, H)$. In order to achieve this, we may have to pass to a smaller covering in the category of admissible formal schemes.

Let (\mathcal{X}, H) be a strictly semistable pair with a covering as in Definition 5.1. We consider the induced formal open covering of the associated admissible formal scheme \mathcal{X} which is defined by completion. Hence \mathcal{X} is covered by formal open subsets \mathcal{U} which admit an étale morphism $\psi : \mathcal{U} \rightarrow \mathcal{S}$ to a formal scheme \mathcal{S} as in (5.1).

It is shown in [GRW14], Proposition 4.1 that, after passing to a refinement, the formal étale covering

$$\psi : \mathcal{U} \rightarrow \mathcal{S}$$

has the property that $\psi^{-1}\{x_0 = \dots = x_{r+s} = 0\}$ is a vertical stratum S in the special fiber \mathcal{X}_s such that for every vertical stratum T the following condition holds: The closure \bar{T} of T in \mathcal{X}_s meets \mathcal{U}_s if and only if $S \subset \bar{T}$.

We define the skeleton of $(\mathcal{U}, H|_{\mathcal{U}})$ as the preimage of the skeleton of the standard pair: $S(\mathcal{U}, H|_{\mathcal{U}}) = \psi^{-1}(S(\mathcal{S}, H(s)))$. This is a subset of the analytic generic fiber \mathcal{U}_η of \mathcal{U} which does not meet the horizontal divisor H .

It follows from results of Berkovich [Ber99] that the étale map ψ actually induces a homeomorphism between $S(\mathcal{U}, H|_{\mathcal{U}})$ and $S(\mathcal{S}, H(s)) \simeq \Delta(\pi, r) \times \mathbb{R}_{\geq 0}^s$. Hence the map

$$\text{Val} \circ \psi : S(\mathcal{U}, H|_{\mathcal{U}}) \rightarrow \Delta(\pi, r) \times \mathbb{R}_{\geq 0}^s$$

is a homeomorphism.

It is shown in [GRW14], 4.5 that the skeleton $S(\mathcal{U}, H|_{\mathcal{U}})$ only depends on the minimal stratum S contained in the special fiber of \mathcal{U} . Therefore we denote it by Δ_S and call it the *canonical polyhedron of S* . The dimensions of S and of the canonical polyhedron Δ_S are defined in an obvious way and add up to d .

As we have seen, Δ_S is homeomorphic to $\Delta(\pi, r) \times \mathbb{R}_{\geq 0}^s$ for suitable data r, s and π as above. We call $\Delta(\pi, r)$ the *finite part* and $\mathbb{R}_{\geq 0}^s$ the *infinite part* of Δ_S .




FIGURE 4. The finite part $\Delta(\pi, r)$ in the case $r = 0$ and $s = 2$, in the case $r = s = 1$ and in the case $r = 2$ and $s = 0$ (from left to right).

As explained above, $S \mapsto \Delta_S$ is a bijective correspondence between faces Δ_S of the skeleton $S(\mathcal{X}, H)$ and vertical strata S induced by the divisor D in the special fiber \mathcal{X}_s . Recall that D is given by the horizontal divisor H plus all irreducible components of \mathcal{X} (which we also call vertical divisors). Note that a vertical stratum T satisfies $S \subset \bar{T}$ if and only if Δ_T is a closed face of Δ_S .

Now we can glue Δ_S and $\Delta_{S'}$ along the union of all canonical polyhedra associated to vertical strata R such that $\bar{R} \supset S \cup S'$. In this way we define the skeleton of (\mathcal{X}, H) as the union of all Δ_S for strata S in the special fiber \mathcal{X}_s :

$$S(\mathcal{X}, H) = \bigcup_S \Delta_S.$$

We obtain a piecewise linear space $S(\mathcal{X}, H)$ whose charts are integral Γ -affine polyhedra.

The skeleton $S(\mathcal{X}, H)$ is a closed subset of the analytic space $X^{\text{an}} \setminus H_K$, where we write H_K for the generic fiber of the support of H on \mathcal{X} . Using methods from [Ber99], one can prove that it is in fact a strong deformation retract of $X^{\text{an}} \setminus H_K$. In [GRW14], Theorem 4.13 it is shown that the retraction map can be extended to a retraction map from X^{an} to a suitable compactification of the skeleton.

If the horizontal divisor $H = 0$, then the skeleton $S(\mathcal{X}, 0)$ is equal to Berkovich's skeleton of a semistable scheme, see [Ber99].

6. FUNCTIONS ON THE SKELETON

Throughout this section we fix a strictly semistable pair (\mathcal{X}, H) . We use the notation from the previous sections.

We want to show that for every non-zero rational function f on X such that the support of the divisor of f is contained in H_K , the function $-\log |f|$ factors through a piecewise integral Γ -affine map on the skeleton. Moreover, we will define and investigate certain slopes.

Theorem 6.1 ([GRW14], Proposition 5.2). *Let f be a non-zero rational function on X such that the support of $\text{div}(f)$ is contained in H_K . We put $U = X \setminus H_K$ and consider the function*

$$F = -\log |f| : U^{\text{an}} \rightarrow \mathbb{R}.$$

Then F factors over through the retraction map $\tau : U^{\text{an}} \rightarrow S(\mathcal{X}, H)$ to the skeleton:

$$\begin{array}{ccc} U^{\text{an}} & \xrightarrow{F} & \mathbb{R} \\ & \searrow \tau & \nearrow F|_{S(\mathcal{X}, H)} \\ & S(\mathcal{X}, H) & \end{array}$$

Moreover, the restriction of F to $S(\mathcal{X}, H)$ is an integral Γ -affine function on each canonical polyhedron.

The theorem is proved by considering the formal building blocks with distinguished strata and using a result of Gubler [Gu07], Proposition 2.11.

Recall that we denote the dimension of X by d . The slope formula for F is basically a balancing condition around each $(d-1)$ -dimensional canonical polyhedron of the skeleton which involves slopes in the direction of all adjacent d -dimensional polyhedra.

Let Δ_S be a d -dimensional canonical polyhedron of the skeleton $S(\mathcal{X}, H)$ containing the $(d-1)$ -dimensional canonical polyhedron Δ_T . Then the stratum S in the special fiber

\mathcal{X}_s is contained in the closure \overline{T} (which is a smooth projective curve), and it is obtained by intersecting \overline{T} with one additional irreducible component of the divisor D .

This component is either vertical, i.e. a component of the special fiber, or horizontal, i.e. given by some component of H . If it is vertical, then the finite part of $\Delta_S \simeq \Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s$ (i.e. the simplex $\Delta(r, \pi)$) is strictly larger than the finite part of Δ_T . In this case we say that Δ_S extends Δ_T in a bounded direction. If the component is horizontal, then the infinite part of Δ_S (i.e. the product $\mathbb{R}_{\geq 0}^s$) is strictly larger than the infinite part of Δ_T . In this case we say that Δ_S extends Δ_T in an unbounded direction.

We define the *bounded degree* $\deg_b(\Delta_T)$ as the number of canonical polyhedral Δ_S extending Δ_T in a bounded direction. Similarly, we define the *unbounded degree* $\deg_u(\Delta_T)$ as the number of canonical polyhedra Δ_S extending Δ_T in an unbounded direction.

In the case $d = 1$, i.e. if X is a curve, all Δ_T are vertices. A one-dimensional canonical polyhedron Δ_S extending Δ_T in a bounded direction is simply an edge of length $v(\pi) > 0$. In this case, the stratum S corresponds to the intersection of two irreducible components in the special fiber \mathcal{X}_s . A one-dimensional canonical polyhedron Δ_S extending Δ_T in an unbounded direction is a ray of the form $\mathbb{R}_{\geq 0}$. In this case, the stratum S is a component of the intersection of the irreducible component of the special fiber given by T with a horizontal component of the divisor. Hence the degree $\deg_b(\Delta_T)$ is the number of bounded edges in Δ_T , and $\deg_u(\Delta_T)$ is the number of unbounded rays starting in the vertex Δ_T .

Now we want to define multiplicities via intersection theory. Since \mathcal{X} is not noetherian, the standard intersection theory tools from algebraic geometry are not available. However, on admissible formal schemes one can use analytic geometry to associate Weil divisors to Cartier divisors, and there is a refined intersection product with Cartier divisors, see [Gu98], [Gu03] and Appendix A in [GRW14].

Definition 6.2. For every vertex $u \in \Delta_T$ we denote by V_u the associated irreducible component of the special fiber \mathcal{X}_s . We define an integer $\alpha(u, \Delta_T)$ as follows:

If Δ_T has zero-dimensional finite part $\{u\}$, we simply put $\alpha(u, \Delta_T) = \deg_b(\Delta_T)$.

If not, then T lies in at least two irreducible components of the special fiber \mathcal{X}_s , which means that the finite part of T is a simplex $\Delta(r, \pi)$ of dimension at least one. If \mathcal{U} is a formal chart associated to S , it is shown in [GRW14], Proposition 4.17 that there exists a unique effective Cartier divisor C_u on \mathcal{U} such that its Weil divisor is equal to $v(\pi)(V_u \cap \mathcal{U}_s)$. We define $-\alpha(u, \Delta_T)$ as the intersection number (C_u, \overline{T}) , which is equal to the degree of the pullback of the line bundle associated to C_u from \mathcal{U} to \overline{T} .

Note that Cartwright [Ca13] introduced the intersection numbers $\alpha(u, \Delta_T)$ in the (noetherian) situation where K° is discrete valuation ring. He uses them to endow the compact skeleton $S(\mathcal{X})$ with the structure of a tropical complex, see [Ca13], Definition 1.1. This notion also involves a local Hodge condition which plays no role in our slope formula. The

paper [Ca13] develops a theory of divisors on tropical complexes and investigates their relation to algebraic divisors.

Here we also need multiplicities for rays in Δ_T , which are defined as one-dimensional faces of the unbounded part of T .

Definition 6.3. *Let H_r be the horizontal component corresponding to the ray r in Δ_T . We put*

$$\alpha(r, \Delta_T) = -(H_r \cdot \bar{T}),$$

where we take the intersection product of \bar{T} with the Cartier divisor H_r on \mathcal{X} .

Let F be a piecewise integral Γ -affine function on the skeleton $S(\mathcal{X}, H)$. Now we are ready to define outgoing slopes of F on a $(d-1)$ -dimensional canonical polyhedron Δ_T along a d -dimensional polyhedron Δ_S .

Definition 6.4. *Let Δ_T be a $(d-1)$ -dimensional canonical polyhedron and let Δ_S be a d -dimensional canonical polyhedron in $S(\mathcal{X}, H)$ containing Δ_T . We denote by $\Delta(r, \pi)$ the finite part of Δ_S . Let $F : \Delta_S \rightarrow \mathbb{R}$ be an integral Γ -affine function.*

i) If Δ_S extends Δ_T in a bounded direction, then there exists a unique vertex w of Δ_S not contained in Δ_T . We put

$$\text{slope}(F; \Delta_T, \Delta_S) = \frac{1}{v(\pi)} \left(F(w) - \frac{1}{\deg_b(\Delta_T)} \sum_{u \in \Delta_T} \alpha(u, \Delta_T) F(u) \right),$$

where we sum over all vertices u in Δ_T .

ii) If Δ_S extends Δ_T in an unbounded direction, then there exists a unique ray s in Δ_S not contained in Δ_T , and we put

$$\text{slope}(F; \Delta_T, \Delta_S) = d_s F - \frac{1}{\deg_u(\Delta_T)} \sum_{r \in \Delta_T} \alpha(r, \Delta_T) d_r F,$$

where we sum over all rays in Δ_T . For any ray r we denote by $d_r F$ the derivative of F along the primitive vector in the direction of r .

If X is a curve, $\Delta_T = u$ is a vertex and Δ_S is an edge with vertices u and w , then $\text{slope}(F; \Delta_T, \Delta_S) = \frac{1}{v(\pi)}(F(w) - F(u))$. If Δ_S is a ray s starting in u , then we simply have $\text{slope}(F; \Delta_T, \Delta_S) = d_s F$. In higher dimensions, the definition is more involved, since the naive slope $\frac{1}{v(\pi)}(F(w) - F(u))$ depends on the choice of a vertex u in Δ_T . Therefore we define a replacement for a weighted midpoint in Δ_T as $\frac{1}{\deg_b(\Delta_T)} \sum_{u \in \Delta_T} \alpha(u, \Delta_T) u$. Note that this point does not necessarily lie in Δ_T , since $\alpha(u, \Delta_T)$ may be negative.

We can now formulate the slope formula for skeleta.

Theorem 6.5 ([GRW14], Theorem 6.9). *Let $f \in K(X)^*$ be a non-zero rational function such that the support of $\operatorname{div}(f)$ is contained in H_K . Let $F : S(\mathcal{X}, H) \rightarrow \mathbb{R}$ be the restriction of the function $-\log|f|$ to the skeleton. Then F is continuous and integral Γ -affine on each canonical polyhedron of $S(\mathcal{X}, H)$, and for all $(d-1)$ -dimensional canonical polyhedra we have*

$$\sum_{\Delta_S \succ \Delta_T} \operatorname{slope}(F; \Delta_T, \Delta_S) = 0,$$

where the sum runs over all d -dimensional canonical polyhedra Δ_S containing Δ_T .

If X is a curve, the slope formula basically says that the sum of all outgoing slopes along edges or rays in a fixed vertex is zero. In this case, the slope formula is shown in [BPR11, Theorem 5.69]. It is a reformulation of the non-archimedean Poincaré-Lelong formula proven in Thuillier's thesis [Thu05], Proposition 3.3.15. The Poincaré-Lelong formula is an equation of currents in the form $dd^c \log|f| = \delta_{\operatorname{div}(f)}$, where dd^c is a certain distribution-valued operator. This version of the slope formula was generalized to higher dimensions in the ground-breaking paper [Ch-Du12], where a theory of differential forms and currents on Berkovich spaces is developed. The approach of [Ch-Du12] uses tropical charts and does not rely on models or skeleta. In higher dimensions we see no direct relation to our slope formula.

7. FAITHFUL TROPICALIZATIONS

We will now investigate the relation between skeleta, which are polyhedral substructures of analytic varieties, and tropicalizations, which are polyhedral images of algebraic or analytic varieties.

7.1. Finding a faithful tropicalization for a skeleton. We start with a strictly semistable pair (\mathcal{X}, H) with skeleton $S(\mathcal{X}, H)$ and generic fiber X . We consider rational maps $f : X \dashrightarrow \mathbb{G}_{m,K}^n$ from X to a split torus. If $U \subset X$ is a Zariski open subvariety where f is defined, then $f|_U : U \rightarrow \mathbb{G}_{m,K}^n$ induces a tropicalization $\operatorname{Trop}_f(U)$ of U , which is defined as the image of the map $\operatorname{trop} \circ f^{\operatorname{an}} : U^{\operatorname{an}} \rightarrow \mathbb{R}^n$ as in section 2.3.

Note that the skeleton is contained in the analytification of every Zariski open subset of X , since it only contains norms on the function field of X .

Definition 7.1. *A rational map $f : X \dashrightarrow \mathbb{G}_{m,K}^n$ from X to a split torus is called a faithful tropicalization of the skeleton $S(\mathcal{X}, H)$ if the following conditions hold:*

- i) The map $\operatorname{trop} \circ f^{\operatorname{an}}$ is injective on $S(\mathcal{X}, H)$.*
- ii) Each canonical polyhedron Δ_S of $S(\mathcal{X}, H)$ can be covered by finitely many integral Γ -affine polyhedra such that the restriction of $\operatorname{trop} \circ f^{\operatorname{an}}$ to each of those polyhedra is a unimodular integral Γ -affine map.*

For the definition of unimodular integral affine maps see section 5.1

It is easy to see that a rational map which is unimodular on the skeleton stays unimodular if we enlarge it with more rational functions on X , see [GRW14], Lemma 9.3.

If we look at a building block $\mathcal{S} = \text{Spec}K^\circ[x_0, \dots, x_d]/(x_1 \dots x_r - \pi)$ as in Definition 5.1 with its tropicalization $\Delta(r, \pi) \times \mathbb{R}_{\geq 0}^s$, we find that the coordinate functions x_0, \dots, x_{r+s} induce a faithful tropicalization. Collecting these rational functions on X for all canonical polyhedra of the skeleton, we get a rational map on X which is locally unimodular on the skeleton. It is shown in the proof of Theorem 9.5 of [GRW14] how to enlarge this collection of rational function in order to ensure injectivity of the tropicalization map on the skeleton. In this way one can show the following result.

Theorem 7.2 ([GRW14], Theorem 9.5). *Let (\mathcal{X}, H) be a strictly semistable pair. Then there exists a collection of non-zero rational functions f_1, \dots, f_n on X such that the resulting rational map $f = (f_1, \dots, f_n) : X \dashrightarrow \mathbb{G}_{m,K}^n$ is a faithful tropicalization of the skeleton $S(\mathcal{X}, H)$.*

7.2. Finding a topological copy of the tropicalization inside the analytic space.

Let us now start with a given tropicalization of an affine K -variety U , i.e. consider a closed immersion $\varphi : U \hookrightarrow (\mathbb{G}_{m,K}^n)^{\text{an}}$ in a split torus. As in section 2.3 we consider the tropicalization map

$$\text{trop}_\varphi = \text{trop} \circ \varphi^{\text{an}} : U^{\text{an}} \rightarrow \mathbb{R}^n.$$

Recall that for every point $\omega \in \text{Trop}_\varphi(U)$ the preimage $\text{trop}_\varphi^{-1}(\omega) \subset U^{\text{an}}$ of the tropicalization map is the Berkovich spectrum of an affinoid algebra A_ω .

Lemma 7.3. *If the tropical multiplicity of ω is equal to one, then A_ω contains a unique Shilov boundary point.*

This claim follows from [BPR11], Remark after Proposition 4.17, which shows how the preimage of tropicalization is related to initial degenerations. It can also be proved directly, see [GRW14], Corollary 10.4.

Now we can show that on the locus of tropical multiplicity one there exists a natural section of the tropicalization map.

Theorem 7.4 ([GRW14], Theorem 10.6). *Let $Z \subset \text{Trop}_\varphi(U)$ be a subset such that the tropical multiplicity of every point in Z is equal to one. By Lemma 7.3 this implies that for every $\omega \in Z$ the affinoid space $\text{trop}_\varphi^{-1}(\omega)$ has a unique Shilov boundary point which we denote by $s(\omega)$.*

The map $s : Z \rightarrow U^{\text{an}}$, given by $\omega \rightarrow s(\omega)$ is continuous and a partial section of the tropicalization map, i.e. on Z we have $\text{trop}_\varphi \circ s = \text{id}_Z$.

Hence the image $s(Z)$ is a subset of U^{an} which is homeomorphic to Z . Moreover, if Z is contained in the closure of its interior in $\text{Trop}_\varphi(U)$, then s is the unique continuous section of the tropicalization map on Z .

We give a sketch of the proof. It is enough to show that the section s is continuous and uniquely defined. Since everything behaves nicely under base change we may assume that the valuation map $|K^*| \rightarrow \mathbb{R}_{>0}$ is surjective. Let us first consider the case that $U = \mathbb{G}_{m,K}^n$ and φ the identity map. Then the section s is the identification of the tropicalization \mathbb{R}^n (which has multiplicity one everywhere) with the skeleton of $\mathbb{G}_{m,K}^n$ as defined in (4.1). It is clear from the explicit description of s in (4.1) that it is continuous in this case.

Moreover, if s' is a different section of the tropicalization map satisfying $s(\omega) \neq s'(\omega)$, we find a Laurent polynomial f on which those two seminorms differ. Since $s(\omega)$ is the unique Shilov boundary point in the fiber of tropicalization, $s'(\omega)$ applied to f is strictly smaller than $s(\omega)$ applied to f . Since $s(\omega)$ is equal to $s'(\omega)$ on all monomials, it follows that the initial degeneration of f at ω cannot be a monomial. Therefore ω is contained in the tropical hypersurface $\text{Trop}(f)$. A continuity argument shows that the same argument works in a small neighbourhood of ω . This is a contradiction since $\text{Trop}(f)$ has codimension one. Therefore s is indeed uniquely determined if φ is the identity map.

For general $\varphi : U \hookrightarrow \mathbb{G}_{m,K}^n$ one shows that there exists a linear map $\mathbb{Z}^n \rightarrow \mathbb{Z}^d$ with the following property: Let $\alpha : \mathbb{G}_{m,K}^n \rightarrow \mathbb{G}_{m,K}^d$ be the corresponding homomorphism of tori and consider $\psi = \alpha \circ \varphi : U \rightarrow \mathbb{G}_{m,K}^d$. Let $S(\mathbb{G}_{m,K}^d)$ be the skeleton of the torus as in (4.1). Then for all $\omega \in Z$ we prove that $\{s(\omega)\} = \text{trop}_{\varphi}^{-1}(\omega) \cap \psi^{-1}(S(\mathbb{G}_{m,K}^d))$. This implies that $s(Z) = \text{trop}_{\varphi}^{-1}(Z) \cap \psi^{-1}(S(\mathbb{G}_{m,K}^d))$ is closed, from which we can deduce continuity of s . Uniqueness follows from uniqueness in the torus case by composing the sections with ψ .

Note that the preceding theorem does not make any assumption on the existence of specific models. If we assume that (\mathcal{X}, H) is a semistable pair and $U = X \setminus H_K$, then the image of the section s defined in Theorem 7.4 is contained in the skeleton $S(\mathcal{X}, H)$. This is shown in [GRW14], Proposition 10.8.

The preceding theorem treats the case of tropicalizations in tori. Let $\varphi : X \hookrightarrow Y$ be a closed embedding in a toric variety Y associated to the fan Δ . Then we may consider the associated tropicalization $\text{Trop}_{\varphi}(X)$ of X , i.e. the image of $\text{trop} \circ \varphi^{\text{an}} : X^{\text{an}} \rightarrow Y^{\text{an}} \rightarrow N_{\mathbb{R}}^{\Delta}$, where $N_{\mathbb{R}}^{\Delta}$ is the associated compactification of $N_{\mathbb{R}}$, see section 2.3. We can apply Theorem 7.4 to all torus orbits. In this way, we get a section of the tropicalization map on the locus $Z \subset \text{Trop}_{\varphi}(X)$ of tropical multiplicity one which is continuous on the intersection with each toric stratum. It is a natural question if this section is continuous on the whole of Z . Note that a positive answer would give an alternative proof of theorem 4.1 for the tropical Grassmannian.

REFERENCES

- [BPR11] M. Baker, S. Payne, J. Rabinoff: *Nonarchimedean geometry, tropicalization and metrics on curves*. Preprint 2011. <http://arxiv.org/abs/1104.0320v2>.

- [BaRu10] M. Baker, R. Rumely: Potential theory and dynamics on the Berkovich projective line. *Mathematical Surveys and Monographs* 159. AMS 2010.
- [Ber90] V. Berkovich: Spectral theory and analytic geometry over non-Archimedean fields. *Mathematical Surveys and Monographs* 33. AMS 1990.
- [Ber99] V. Berkovich: *Smooth p -adic spaces are locally contractible*. *Invent. Math.* 137 (1999) 1-84.
- [Ber04] V. Berkovich: *Smooth p -adic spaces are locally contractible. II*. In: *Geometric aspects of Dwork theory*. De Gruyter 2004, p. 293-370.
- [Bo14] S. Bosch: *Lectures on Formal and Rigid Geometry*. *Lecture Notes in Mathematics* 2105. Springer 2014.
- [BGR84] S. Bosch, S. Güntzer, R. Remmert: *Non-Archimedean Analysis*. Springer 1984.
- [Ca13] D. Cartwright: *Tropical Complexes*. Preprint 2013. <http://arxiv.org/abs/1308.3813>.
- [CHW14] M. Cueto, M. Häbich, A. Werner: *Faithful tropicalization of the Grassmannian of planes*. *Math. Ann.* 360 (2014) 391-437.
- [Ch-Du12] A. Chambert-Loir, A. Ducros: *Formes différentielles réelles et courants sur les espaces de Berkovich*. Preprint 2012. <http://arxiv.org/abs/1204.6277>.
- [Co08] B. Conrad: *Several approaches to non-archimedean geometry*. In: *p -adic geometry*. University Lecture Series 45. AMS 2008.
- [Du07] A. Ducros: *Espaces analytiques p -adiques au sens de Berkovich*. *Séminaire Bourbaki* 2005/06. Astérisque 311, exposé 958 (2007)
- [dJ96] J.A. de Jong: *Smoothness, semi-stability and alterations*. *Publ. Math. IHES* 83 (1996) 51-93.
- [DP14] J. Draisma, E. Postingshel: *Faithful tropicalization and torus actions*. Preprint 2014. <http://arxiv.org/abs/1404.4715>.
- [Du] A. Ducros: *Espaces de Berkovich, squelettes et théorie des modèles*. *Confluentes Math.* 4 (2012) 1250007.
- [EKL06] M. Einsiedler, M. Kapranov, D. Lind: *Non-Archimedean amoebas and tropical varieties*. *J. Reine Angew. Math.* 601 (2006) 139-157.
- [FGP14] T. Foster, P. Gross, S. Payne: *Limits of tropicalizations*. *Israel J. Math.* 201 (2014) 835-846.
- [Gu98] W. Gubler: *Local heights of subvarieties over non-Archimedean fields*. *J. Reine Angew. Math.* 498 (1998) 61-113.
- [Gu03] W. Gubler: *Local and canonical heights of subvarieties*. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) 2 (2003) 711-760.
- [Gu07] W. Gubler: *Tropical varieties for non-Archimedean analytic spaces*. *Invent Math.* 169 (2007) 321-376.
- [Gu13] W. Gubler: *A guide to tropicalizations*. In: *Algebraic and combinatorial aspects of tropical geometry*. *Contemp. Math.* 589. AMS 2013, p. 125-189.
- [GRW14] W. Gubler, J. Rabinoff, A. Werner: *Skeletons and Tropicalizations*. Preprint 2014. <http://arxiv.org/abs/1404.7044>.
- [Ma-St] D. Maclagan, B. Sturmfels: *Introduction to Tropical Geometry*. To appear 2015.
- [Pay09] S. Payne: *Analytification is the limit of tropicalization*. *Math. Res. Lett.* 16 (2009) 543-556.
- [SS04] D. Speyer, B. Sturmfels: *The tropical Grassmannian*. *Adv. Geom.* 4 (2004) 389-411.
- [Tem] M. Temkin: *Introduction to Berkovich analytic spaces*. In: *Berkovich Spaces and Applications*. *Lecture Notes in Mathematics* 2119. Springer 2015, p. 3-67.
- [Thu05] A. Thuillier: *Théorie du potentiel sur les courbes on géométrie analytique non-archimédienne. Applications à la théorie d'Arakelov*. Thèse Rennes 2005. <http://tel.ccsd.cnrs.fr/documents/archives0/00/01/09/90/index.html>
- [Tyo12] I. Tyomkin: *Tropical geometry and correspondence theorem via toric stacks*. *Math. Ann.* 353 (2012) 945-995.

Institut für Mathematik
Goethe-Universität Frankfurt
Robert-Mayer-Strasse 8

