


6. (a) $x = \sqrt{3}$ and $y = -1 \Rightarrow r = \sqrt{(\sqrt{3})^2 + (-1)^2} = 2$ and $\tan \theta = \frac{-1}{\sqrt{3}} \quad [\theta = -\frac{\pi}{6} + n\pi]$. Since $(\sqrt{3}, -1)$ is in the


fourth quadrant, the polar coordinates are (i) $(2, \frac{11\pi}{6})$ and (ii) $(-2, \frac{5\pi}{6})$.

(b) $x = -6$ and $y = 0 \Rightarrow r = \sqrt{(-6)^2 + 0^2} = 6$ and $\tan \theta = \frac{0}{-6} = 0 \quad [\theta = n\pi]$. Since $(-6, 0)$ is on the negative x -axis, the polar coordinates are (i) $(6, \pi)$ and (ii) $(-6, 0)$.


7. $1 < r \leq 3$. The curves $r = 1$ and $r = 3$ represent circles centered at O with radius 1 and 3, respectively. So $1 < r \leq 3$ represents the region outside the radius 1 circle and on or inside the radius 3 circle. Note that θ can take on any value.




8. $r \geq 2, 0 \leq \theta \leq \pi$. This is the region on or outside the circle $r = 2$ in the first and second quadrants.




9. $0 \leq r \leq 1, -\pi/2 \leq \theta \leq \pi/2$. This is the region on or inside the circle $r = 1$ in the first and fourth quadrants.




10. $3 < r < 5, 2\pi/3 \leq \theta \leq 4\pi/3$



11. $2 \leq r < 4, 3\pi/4 \leq \theta \leq 7\pi/4$



12. $r \geq 0$, $\pi \leq \theta \leq 5\pi/2$. This is the region in the third, fourth, and first quadrants including the origin and points on the negative x -axis and positive y -axis.



13. Converting the polar coordinates $(4, \frac{4\pi}{3})$ and $(6, \frac{5\pi}{3})$ to Cartesian coordinates gives us $(4 \cos \frac{4\pi}{3}, 4 \sin \frac{4\pi}{3}) = (-2, -2\sqrt{3})$ and $(6 \cos \frac{5\pi}{3}, 6 \sin \frac{5\pi}{3}) = (3, -3\sqrt{3})$. Now use the distance formula

$$\begin{aligned} d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{[3 - (-2)]^2 + [-3\sqrt{3} - (-2\sqrt{3})]^2} \\ &= \sqrt{5^2 + (-\sqrt{3})^2} = \sqrt{25 + 3} = \sqrt{28} = 2\sqrt{7} \end{aligned}$$

14. The points (r_1, θ_1) and (r_2, θ_2) in Cartesian coordinates are $(r_1 \cos \theta_1, r_1 \sin \theta_1)$ and $(r_2 \cos \theta_2, r_2 \sin \theta_2)$, respectively. The *square* of the distance between them is

$$\begin{aligned} &(r_2 \cos \theta_2 - r_1 \cos \theta_1)^2 + (r_2 \sin \theta_2 - r_1 \sin \theta_1)^2 \\ &= (r_2^2 \cos^2 \theta_2 - 2r_1 r_2 \cos \theta_1 \cos \theta_2 + r_1^2 \cos^2 \theta_1) + (r_2^2 \sin^2 \theta_2 - 2r_1 r_2 \sin \theta_1 \sin \theta_2 + r_1^2 \sin^2 \theta_1) \\ &= r_1^2 (\sin^2 \theta_1 + \cos^2 \theta_1) + r_2^2 (\sin^2 \theta_2 + \cos^2 \theta_2) - 2r_1 r_2 (\cos \theta_1 \cos \theta_2 + \sin \theta_1 \sin \theta_2) \\ &= r_1^2 - 2r_1 r_2 \cos(\theta_1 - \theta_2) + r_2^2, \end{aligned}$$

so the distance between them is $\sqrt{r_1^2 - 2r_1 r_2 \cos(\theta_1 - \theta_2) + r_2^2}$.

15. $r^2 = 5 \Leftrightarrow x^2 + y^2 = 5$, a circle of radius $\sqrt{5}$ centered at the origin.

16. $r = 4 \sec \theta \Leftrightarrow \frac{r}{\sec \theta} = 4 \Leftrightarrow r \cos \theta = 4 \Leftrightarrow x = 4$, a vertical line.

17. $r = 5 \cos \theta \Rightarrow r^2 = 5r \cos \theta \Leftrightarrow x^2 + y^2 = 5x \Leftrightarrow x^2 - 5x + \frac{25}{4} + y^2 = \frac{25}{4} \Leftrightarrow (x - \frac{5}{2})^2 + y^2 = \frac{25}{4}$,

a circle of radius $\frac{5}{2}$ centered at $(\frac{5}{2}, 0)$. The first two equations are actually equivalent since $r^2 = 5r \cos \theta \Rightarrow$


$r(r - 5 \cos \theta) = 0 \Rightarrow r = 0$ or $r = 5 \cos \theta$. But $r = 5 \cos \theta$ gives the point $r = 0$ (the pole) when $\theta = 0$. Thus, the equation $r = 5 \cos \theta$ is equivalent to the compound condition $(r = 0 \text{ or } r = 5 \cos \theta)$.

18. $\theta = \frac{\pi}{3} \Rightarrow \tan \theta = \tan \frac{\pi}{3} \Rightarrow \frac{y}{x} = \sqrt{3} \Leftrightarrow y = \sqrt{3}x$, a line through the origin.


19. $r^2 \cos 2\theta = 1 \Leftrightarrow r^2 (\cos^2 \theta - \sin^2 \theta) = 1 \Leftrightarrow (r \cos \theta)^2 - (r \sin \theta)^2 = 1 \Leftrightarrow x^2 - y^2 = 1$, a hyperbola centered at the origin with foci on the x -axis.

20. $r^2 \sin 2\theta = 1 \Leftrightarrow r^2 (2 \sin \theta \cos \theta) = 1 \Leftrightarrow 2(r \cos \theta)(r \sin \theta) = 1 \Leftrightarrow 2xy = 1 \Leftrightarrow xy = \frac{1}{2}$, a hyperbola centered at the origin with foci on the line $y = x$.


31. r has a maximum value of approximately 1 slightly before $\theta = \frac{\pi}{4}$ and slightly after $\theta = \frac{7\pi}{4}$. r has a minimum value of -2 when $\theta = \pi$. The graph touches the pole ($r = 0$) when $\theta = 0, \frac{\pi}{2}, \frac{3\pi}{2}$, and 2π . Since r is positive in the θ -intervals $(0, \frac{\pi}{2})$ and $(\frac{3\pi}{2}, 2\pi)$, and negative in the interval $(\frac{\pi}{2}, \frac{3\pi}{2})$, the graph lies entirely in the first and fourth quadrants.




32. r increases from 0 to 1 (local max) in the interval $[0, \frac{\pi}{2}]$. It then decreases slightly, after which r increases to a maximum of 2 at $\theta = \pi$. The graph is symmetric about $\theta = \pi$, so the polar curve is symmetric about the polar axis.




33. $r = -2 \sin \theta$




34. $r = 1 - \cos \theta$




35. $r = 2(1 + \cos \theta)$




36. $r = 1 + 2 \cos \theta$




37. $r = \theta, \theta \geq 0$




38. $r = \theta^2, -2\pi \leq \theta \leq 2\pi$



39. $r = 3 \cos 3\theta$



40. $r = -\sin 5\theta$



41. $r = 2 \cos 4\theta$

