

# Projective (symmetries of) TQFTs

Jackson Van Dyke

December 18, 2023

# Table of contents

- 1 Anomalies
- 2 Anomal(ous symmetr)ies of three-dimensional TQFTs
- 3 Future directions
- 4 (If time allows:) Higher projective symmetries

- **TQFT** = fully-extended symmetric-monoidal functor  $\mathbf{Bord}_d^{\text{fr}} \rightarrow \mathcal{T}$ .
- **Relative/twisted/boundary theory** is a (lax) natural transformation<sup>1</sup>  $F: 1 \rightarrow \beta$  (or  $\beta \rightarrow 1$ ).
- An **anomaly**<sup>2</sup> is an invertible once-categorified  $d$ -dimensional TQFT  $\alpha$ , and an anomalous  $d$ -dimensional TQFT is a relative theory  $F: \alpha \rightarrow 1$ .

## Example

Let  $V$  be a finite-dimensional vector space.

- $V$  classifies a TQFT  $\mathbf{Bord}_1^{\text{fr}} \rightarrow \mathbf{Vect}$ .
- $G \rightarrow \text{GL}(V)$  classifies a TQFT  $\mathbf{Bord}_1^{BG} \rightarrow \mathbf{Vect}$ .
- $G \rightarrow \text{PGL}(V)$  classifies an anomalous 1-d TQFT on  $\mathbf{Bord}_1^{BG}$ .

---

<sup>1</sup>Theo Johnson-Freyd and Claudia Scheimbauer. (Op)lax natural transformations, twisted quantum field theories, and “even higher” Morita categories, 2017

<sup>2</sup>Daniel S. Freed. What is an anomaly?, 2023

# Anomal(ous symmetr)ies of three-dimensional TQFTs

Building on existing results<sup>3,4</sup> I introduce:<sup>5</sup>  $B^2\mu_q \hookrightarrow 3\text{Pin} \twoheadrightarrow \text{O}(L \oplus L^\vee)$ .

## Theorem (VD<sup>5</sup>)

*The framed Dijkgraaf-Witten theory for a finite abelian group  $L$  canonically defines the following, which are equivalent:*

- a symmetric-monoidal functor  $\mathbf{Bord}_3^{B 3\text{Pin}(L \oplus L^*, \text{ev})} \rightarrow \mathbf{Fus}$
- an anomalous theory on  $\mathbf{Bord}_3^{B \text{O}(L \oplus L^*)}$
- $\text{O}$  acts via “twice-categorified integral transforms”<sup>6</sup>.
- We can replace  $\text{O}$  with the 2-group  $\text{Aut}_{\mathbf{EqBr}} \sigma_{BL}^3(S^1)$ , and then a certain “level” controls the non-triviality of the anomaly<sup>5</sup>.

<sup>3</sup> Pavel Etingof, Dmitri Nikshych, and Victor Ostrik. [Fusion categories and homotopy theory \(appendix by E. Meir\)](#), 2010

<sup>4</sup> Jürgen Fuchs, Jan Priel, Christoph Schweigert, and Alessandro Valentino. [On the Brauer groups of symmetries of abelian Dijkgraaf-Witten theories](#), 2015

<sup>5</sup> Jackson Van Dyke. [Projective symmetries of three-dimensional TQFTs](#), 2023.  
arXiv: 2311.01637 [math.QA]

<sup>6</sup> Jackson Van Dyke. [Symmetries of quantization of finite groupoids](#), 2023.  
arXiv: 2312.00117 [math.QA]

1-dimensional	3-dimensional
$(V, q)$	$(A, q)$
$SO(V, q) \subset O(V, q)$	$SO(A, q) \subset O(A, q)$
$\mathbf{k}^\times$	$B^2\mathbf{k}^\times$
$\text{Cliff}(V)$	$\mathcal{A} = (\mathbf{Vect}[A], *, \beta_q)$
$\{x, y\} = b_q(x, y)$	$\beta_q: \mathbf{k}_a * \mathbf{k}_b \xrightarrow{b_q(a,b) \text{ id}} \mathbf{k}_b * \mathbf{k}_a$
$V \rtimes O(V, q)$	$\text{Aut}_{\text{EqBr}}(\mathcal{A})$
$\text{Pin}(V, q)$	$3\text{Pin}(A, q)$
$\text{Spin}(V, q)$	$3\text{Spin}(A, q)$
$V \simeq L \oplus L^*$	$A \simeq L \oplus L^*$
$\wedge^\bullet L^*$	$\mathcal{C} = (\mathbf{Vect}[L^*], *)$
$\text{End}(\wedge^\bullet L^*) \simeq \text{Cliff}$	$\text{Aut}_{\text{Fus}}(\mathcal{C}) \simeq \text{Pic}(\mathcal{A})$

# Future directions

- Analogous results for fusion 2-categories<sup>7</sup>?
- $\mathbb{P}\mathbf{Fus}$  replaced with the projectivization of the  $(\infty, n + m + 2)$ -category  $\mathbf{Alg}_n(m \mathbf{Pr}^L)$  á la JFS<sup>1</sup>?
- Gapped systems, topological phases of matter
- Non-semisimple finite ribbon categories
  - Link and manifold invariants<sup>8</sup>
  - Rozansky-Witten theory and relative Langlands

## Conjecture

*The truncation of  $\mathbf{Aut}(\mathbf{RW}_M(*))$  to a group is  $\mathrm{Sp}(M)$ .*

**Rmk:** The  $k$ -invariant of  $B \mathbf{Aut}$  in  $H^4(B \mathrm{Sp}(M), \mathbb{C}^\times)$  would then be the projectivity/anomaly of the action  $\mathrm{Sp}(M) \curvearrowright \mathbf{RW}_M$ .

<sup>7</sup> Christopher L. Douglas and David J. Reutter. [Fusion 2-categories and a state-sum invariant for 4-manifolds](#), 2018

<sup>1</sup> Theo Johnson-Freyd and Claudia Scheimbauer. [\(Op\)lax natural transformations, twisted quantum field theories, and “even higher” Morita categories](#), 2017

<sup>8</sup> Johannes Berger, Azat M. Gainutdinov, and Ingo Runkel. [Non-semisimple link and manifold invariants for symplectic fermions](#), 2023

Thank You!

# References



Johannes Berger, Azat M. Gainutdinov, and Ingo Runkel.  
Non-semisimple link and manifold invariants for symplectic fermions, 2023.



Christopher L. Douglas and David J. Reutter.  
Fusion 2-categories and a state-sum invariant for 4-manifolds, 2018.



Pavel Etingof, Dmitri Nikshych, and Victor Ostrik.  
Fusion categories and homotopy theory (appendix by E. Meir), 2010.



Daniel S. Freed.  
What is an anomaly?, 2023.



Daniel S. Freed, Gregory W. Moore, and Constantin Teleman.  
Topological symmetry in quantum field theory, 2022.



Jürgen Fuchs, Jan Priel, Christoph Schweigert, and Alessandro Valentino.  
On the Brauer groups of symmetries of abelian Dijkgraaf-Witten theories, 2015.



Theo Johnson-Freyd and Claudia Scheimbauer.  
(Op)lax natural transformations, twisted quantum field theories, and “even higher” Morita categories, 2017.



Jackson Van Dyke.  
Projective symmetries of three-dimensional TQFTs, 2023.  
[arXiv: 2311.01637 \[math.QA\]](https://arxiv.org/abs/2311.01637).



Jackson Van Dyke.  
Symmetries of quantization of finite groupoids, 2023.  
[arXiv: 2312.00117 \[math.QA\]](https://arxiv.org/abs/2312.00117).



# Higher projective symmetries

In Theorem C.16<sup>5</sup>, I relate twisted quantization<sup>9</sup> with anomalies:

## Theorem (VD<sup>5</sup>)

$$\alpha_c \rightarrow 1 \quad \iff \quad 1 \rightarrow \sigma_{X,c}^{d+1}$$

E.g. (projective) group rep. mod. / (twisted) group alg.

Given a **trivialization**, they will reduce to (the same)  $X$ -theories:

$$1 \xrightarrow{\sim} \alpha_c \xrightarrow{F_\alpha} 1$$

$F_X$

$$1 \longrightarrow \sigma_{X,c}^{d+1} \xrightarrow{\quad} \sigma_X^{d+1}$$

$F_X$

<sup>5</sup> Jackson Van Dyke. [Projective symmetries of three-dimensional TQFTs](#), 2023. arXiv: 2311.01637 [math.QA]

<sup>9</sup> Daniel S. Freed, Gregory W. Moore, and Constantin Teleman. [Topological symmetry in quantum field theory](#), 2022