

# DEFINITION OF THE HOMOLOGY COBORDISM INVARIANTS

LECTURE: JACKSON VAN DYKE

The point of today will be to see that  $\text{SWFH}^{\text{Pin}(2)}(Y, \mathbb{F})$  gives rise to some  $\beta : \Theta_3^H \rightarrow \mathbb{Z}$ . Recall  $\Theta_3^H$  is the homology cobordism group consisting of  $Y^3$  oriented  $\mathbb{Z}HS^3$  (integral homology three spheres). The upshot will be that  $\beta$  satisfies:

$$(1) \quad \beta(-Y) = -\beta(Y) \qquad \mu(Y) = \beta(Y) \pmod{2}$$

where  $\mu$  is the Rokhlin homomorphism, so the SES:

$$(2) \quad 0 \rightarrow \ker \mu \rightarrow \Theta_3^H \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 0$$

does not split.

## 1. THE MODULE STRUCTURE ON EQUIVARIANT HOMOLOGY

Let  $G$  be a Lie group acting on a space  $X$ . The Borel homology is

$$(3) \quad H_*^G(X) = H_*(X \times_G EG)$$

where  $EG$  is the universal bundle. This is a module over  $H_G^*(\text{pt}) = H^*(BG)$ . For us  $G = \text{Pin}(2)$ , and we want to get our hands on

$$(4) \quad H^*(B\text{Pin}(2), \mathbb{F}) .$$

Recall  $\text{Pin}(2) = S^1 \cup jS^1 \subset \mathbb{H}$ . Also recall  $\text{SU}(2)$  is isomorphic to the unit elements of  $\mathbb{H}$ , so it is diffeomorphic to  $S^3$ . This means we get a fibration:

$$(5) \quad \begin{array}{ccc} \text{Pin}(2) & \xrightarrow{i} & \text{SU}(2) \\ & & \downarrow \psi \\ & & \mathbb{R}\mathbb{P}^2 \end{array}$$

where  $i$  is the inclusion and  $\psi$  is the Hopf fibration follows by the involution on  $S^2$ .

In such a setting, we get another fibration<sup>1</sup>

$$(6) \quad \begin{array}{ccc} \mathbb{R}\mathbb{P}^2 & \longrightarrow & B\text{Pin}(2) \\ & & \downarrow \\ & & B\text{SU}(2) = \mathbb{H}\mathbb{P}^\infty \end{array}$$

where  $\mathbb{H}\mathbb{P}^\infty$  is the infinite-dimensional quaternionic projective space.

Now we calculate  $H^*(B\text{Pin}(2), \mathbb{F})$  using a Leray-Serre spectral sequence. Recall that this is

$$(7) \quad E_2^{p,q} = H^p(B\text{SU}(2), H^q(\mathbb{R}\mathbb{P}^2)) \Rightarrow H^{p+q}(B\text{Pin}(2)) .$$

---

*Date:* October 18, 2019.

<sup>1</sup>Explicitly this is given by the inclusion of orbits. See [Peter May's response to this post on MathOverflow](#).



in each grading. Furthermore, we can recover the module action on Borel homology from the one on Borel cohomology. The upshot is that  $\text{SWFH}_*^{\text{Pin}(2)}$  is the homology of a complex (the equivariant cellular complex of the Conley index) of the form:

$$(18) \quad \begin{array}{c} \vdots \\ \mathbb{F} \\ \downarrow v \\ 0 \\ \downarrow v \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \\ \downarrow v \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \\ \downarrow v \\ \mathbb{F} \\ \downarrow v \\ 0 \\ \downarrow v \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \\ \downarrow v \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \\ \downarrow v \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \\ \downarrow v \\ \mathbb{F} \end{array} \quad \begin{array}{c} \text{(finite part)} \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \xrightarrow{q} \mathbb{F} \\ \mathbb{F} \xrightarrow{q} \mathbb{F} \end{array} .$$

The finite part can be any finite dimensional vector space, with some  $v$  and  $q$  actions. There can be various other differentials both inside it and relating to the infinite towers. The important part is that there are always three infinite  $v$  towers in the complex, and they produce three infinite towers in homology. These towers correspond to the  $S^1$ -fixed point set of  $\text{SWF}(Y)$ , and the finite part comes from the cells with a free  $\text{Pin}(2)$  action.

The upshot is that since  $(\text{SWF}(Y))^{S^1} = S^{n(Y,g)}$  and  $n(Y,g) \equiv 2\mu \pmod{4}$ , we have:

- All elements in the first tower are in degree  $2\mu \pmod{4}$ ,
- all elements in the second tower are in degree  $2\mu + 1 \pmod{4}$ , and
- all elements in the third tower are in degree  $2\mu + 2 \pmod{4}$ .

### 3. DEFINITION OF THE INVARIANTS

Write  $A, B, C \in \mathbb{Z}$  for the lowest degrees of each infinite  $v$ -tower in homology. Now define the following:

$$(19) \quad \alpha := \frac{A}{2} \quad \beta := \frac{B-1}{2} \quad \gamma := \frac{C-2}{2} .$$

The point is that:

$$(20) \quad A, \quad B-1, \quad C-2, \quad \equiv 2\mu \pmod{4}$$

so therefore

$$(21) \quad \alpha, \beta, \gamma \equiv \mu \pmod{2} .$$

Note that the module structure requires

$$(22) \quad \alpha \geq \beta \geq \gamma .$$

#### 4. DESCENT TO HOMOLOGY COBORDISM

All we need to do now is check that  $\alpha$ ,  $\beta$ , and  $\gamma$  descend to maps  $\Theta_3^H \rightarrow \mathbb{Z}$ . This will use the construction of cobordism maps on Seiber-Witten Floer spectra.

Let  $W^4$  be a smooth oriented Spin (4) cobordism with  $b_1(W) = 0$ , and with  $\partial W = (-Y_0) \cup Y_1$ . (We really just care about when  $W$  is a homology cobordism between homology three spheres.) We can consider the Seiberg-Witten equations on  $W$ , and do a finite-dimensional approximation to the solution space in the same way that we did in three dimensions. This gets more complicated. . .

In the end we get a stable equivariant map between two suspension spectra

$$(23) \quad \Psi_W : \Sigma^{m\mathbb{H}} \text{SWF}(Y_0) \rightarrow \Sigma^{n\tilde{\mathbb{R}}} \text{SWF}(Y_1) .$$

(Recall  $\tilde{\mathbb{R}}$  is the rep where  $S^1$  acts trivially, and  $j$  acts by multiplication by  $-1$ .) In particular:

$$(24) \quad m = \frac{-\sigma(W)}{8} = \text{ind } \not{D} \quad n = b_2^+(W) = \text{ind}(d^+) .$$

In the case that  $W$  is a smooth oriented homology cobordism between homology spheres, there is a unique Spin (4) structure,  $b_1(W) = 0$ , and  $n = m = 0$ . Let  $F_W$  be the homomorphism induced on Pin (2)-equivariant homology by the map  $\Psi_W$ .

It follows from equivariant localization that in degree  $k \gg 0$ , the map  $F_W$  is an isomorphism. Further  $F_W$  is a module map, so we have a commutative diagram

$$(25) \quad \begin{array}{ccc} \mathbb{F} & \xrightarrow{F_W} & \mathbb{F} \\ \downarrow v & & \downarrow v \\ \mathbb{F} & \xrightarrow{F_W} & \mathbb{F} \end{array} .$$

Because of the module structure, we cannot have  $\alpha(Y_1) < \alpha(Y_0)$ , and likewise for  $\beta$  and  $\gamma$ . In conclusion,

$$(26) \quad \alpha(Y_1) \geq \alpha(Y_0) \quad \beta(Y_1) \geq \beta(Y_0) \quad \gamma(Y_1) \geq \gamma(Y_0) .$$

Now we can simply repeat the entire argument with reversed orientation and reversed direction of  $W$ . Therefore we get

$$(27) \quad \alpha(Y_1) \leq \alpha(Y_0) \quad \beta(Y_1) \leq \beta(Y_0) \quad \gamma(Y_1) \leq \gamma(Y_0) ,$$

and therefore we have equalities. Therefore  $\alpha$ ,  $\beta$ , and  $\gamma$  all descend to maps

$$(28) \quad \Theta_3^H \rightarrow \mathbb{Z} .$$